

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 July 2003 (17.07.2003)

PCT

(10) International Publication Number
WO 03/057235 A2

(51) International Patent Classification⁷: **A61K 38/00**

(21) International Application Number: **PCT/GB03/00062**

(22) International Filing Date: 10 January 2003 (10.01.2003)

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:

0200507.2 10 January 2002 (10.01.2002) GB

60/392,109 28 June 2002 (28.06.2002) US

PCT/US02/31944

24 September 2002 (24.09.2002) US

(71) Applicants (for all designated States except US): **IMPERIAL COLLEGE INNOVATIONS LTD** [GB/GB]; 47 Prince's Gate, Exhibition Road, London SW7 2QA (GB). **OREGON HEALTH AND SCIENCE UNIVERSITY** [US/US]; 2525 S.W. First Avenue, Suite AD-120, Portland, OR 97201 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **COWLEY, Michael**

[AU/US]; 6724 S.E. 19th Avenue, Portland, OR 97202 (US). **CONE, Roger** [US/US]; 16563 S. Hattan Road, Oregon City, OR 97045 (US). **LOW, Malcolm** [NZ/US]; 4650 S.W. Upper Drive, Lake Oswego, OR 97035 (US). **BUTLER, Andrew** [NZ/US]; 3730 S.W. 12th Avenue, Apartment 10, Portland, OR 97201 (US). **BLOOM, Stephen, Robert** [GB/GB]; Imperial College Innovations Limited, 47 Prince's Gate, Exhibition Road, London SW7 2QA (GB). **SMALL, Caroline, Jane** [GB/GB]; Imperial College Innovations Limited, 47 Prince's Gate, Exhibition Road, London SW7 2QA (GB). **BATTERHAM, Rachel, Louise** [GB/GB]; Imperial College Innovations Limited, 47 Prince's Gate, Exhibition Road, London SW7 2QA (GB). **GHATEI, Mohammad, Ali** [GB/GB]; Imperial College Innovations Limited, 47 Prince's Gate, Exhibition Road, London SW7 2QA (GB).

(74) Agents: **SILVESTON, Judith et al.**; Abel & Imray, 20 Red Lion Street, London WC1R 4PQ (GB).


(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,

[Continued on next page]

(54) Title: MODIFICATION OF FEEDING BEHAVIOR



WO 03/057235 A2



(57) Abstract: Methods are disclosed for decreasing calorie intake, food intake, and appetite in a subject. The methods include peripherally administering PYY or an agonist thereof and GLP-1 or an agonist thereof to the subject, simultaneously or sequentially, thereby decreasing the calorie intake of the subject.

BEST AVAILABLE COPY



SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

-1-

MODIFICATION OF FEEDING BEHAVIOUR

PRIORITY CLAIM

This application claims the benefit of UK Application No. GB0200507.2
5 filed January 10, 2002, and International Application PCT/ US02/31944
which are both incorporated by reference in their entirety herein.

STATEMENT OF GOVERNMENT SUPPORT

This disclosure was made in part with United States government support
10 pursuant to grants RR00163, DK51730 and DK55819, from the National Institutes of
Health. The United States government has certain rights in the disclosure.

FIELD

This application relates to the use of agents to control appetite, feeding, food
15 intake, energy expenditure and calorie intake, particularly in the field of obesity.

BACKGROUND

According to the National Health and Nutrition Examination Survey
(NHANES III, 1988 to 1994), between one third and one half of men and women in
20 the United States are overweight. In the United States, sixty percent of men and
fifty-one percent of women, of the age of 20 or older, are either overweight or obese.
In addition, a large percentage of children in the United States are overweight or
obese.

The cause of obesity is complex and multi-factorial. Increasing evidence
25 suggests that obesity is not a simple problem of self-control but is a complex
disorder involving appetite regulation and energy metabolism. In addition, obesity
is associated with a variety of conditions associated with increased morbidity and
mortality in a population. Although the etiology of obesity is not definitively
established, genetic, metabolic, biochemical, cultural and psychosocial factors are
30 believed to contribute. In general, obesity has been described as a condition in
which excess body fat puts an individual at a health risk.

-2-

There is strong evidence that obesity is associated with increased morbidity and mortality. Disease risk, such as cardiovascular disease risk and type 2 diabetes disease risk, increases independently with increased body mass index (BMI). Indeed, this risk has been quantified as a five percent increase in the risk of cardiac disease for females, and a seven percent increase in the risk of cardiac disease for males, for each point of a BMI greater than 24.9 (see Kenchaiah et al., *N. Engl. J. Med.* 347:305, 2002; Massie, *N. Engl. J. Med.* 347:358, 2002). In addition, there is substantial evidence that weight loss in obese persons reduces important disease risk factors. Even a small weight loss, such as 10% of the initial body weight in both 10 overweight and obese adults has been associated with a decrease in risk factors such as hypertension, hyperlipidemia, and hyperglycemia.

Although diet and exercise provide a simple process to decrease weight gain, overweight and obese individuals often cannot sufficiently control these factors to effectively lose weight. Pharmacotherapy is available; several weight loss drugs 15 have been approved by the Food and Drug Administration that can be used as part of a comprehensive weight loss program. However, many of these drugs have serious adverse side effects. When less invasive methods have failed, and the patient is at high risk for obesity related morbidity or mortality, weight loss surgery is an option in carefully selected patients with clinically severe obesity. However, these 20 treatments are high-risk, and suitable for use in only a limited number of patients. It is not only obese subjects who wish to lose weight. People with weight within the recommended range, for example, in the upper part of the recommended range, may wish to reduce their weight, to bring it closer to the ideal weight. Thus, a need remains for agents that can be used to effect weight loss in overweight and obese 25 subjects.

SUMMARY

Disclosed herein are findings that peripheral administration of PYY, or an agonist thereof, to a subject results in decreased food intake, caloric intake, and 30 appetite, and an alteration in energy metabolism. The subject can be any subject, including, but not limited to, a human subject. In several embodiments, the subject

desires to lose weight, is obese, overweight, or suffers from a weight-related disorder. PYY₃₋₃₆ can preferably be administered to the subject.

Also disclosed are findings that co-administration of PYY, or an agonist thereof, for example, PYY₃₋₃₆, with GLP-1 or an agonist thereof to a subject results in a greater decrease in food intake, caloric intake, and appetite, than when PYY or an agonist thereof is administered alone. The effect is synergistic.

In one embodiment, a method is disclosed for decreasing calorie intake in a subject. The method includes peripherally administering a therapeutically effective amount of PYY or an agonist thereof and GLP-1 or an agonist thereof to the subject, thereby decreasing the calorie intake of the subject.

In another embodiment, a method is disclosed for decreasing appetite in a subject. The method includes peripherally administering a therapeutically effective amount of PYY or an agonist thereof and GLP-1 or an agonist thereof to the subject, thereby decreasing the appetite of the subject.

In a further embodiment, a method is disclosed for decreasing food intake in a subject. The method includes peripherally administering a therapeutically effective amount of PYY or an agonist thereof, and GLP-1 or an agonist thereof to the subject, thereby decreasing the food intake of the subject.

In yet another embodiment, a method is disclosed herein for increasing energy expenditure in a subject. The method includes peripherally administering a therapeutically effective amount of PYY or an agonist thereof, and GLP-1 or an agonist thereof to the subject, thereby increasing energy expenditure in the subject.

A method is also disclosed for decreasing calorie intake, food intake, or appetite in a human subject. The method includes peripherally injecting a therapeutically effective amount of PYY or an agonist thereof in a pharmaceutically acceptable carrier to the subject in a pulse dose, thereby decreasing the calorie intake, food intake, or appetite of the subject, and also administering GLP-1 or an agonist thereof.

In each method above, the GLP-1 or an agonist thereof may be administered simultaneously or substantially simultaneously as the PYY or agonist thereof, or sequentially, in any order. The PYY or agonist thereof and the GLP-1 or agonist thereof may be administered in a single pharmaceutical composition or in separate

compositions, and they may be administered by the same route or by different routes.

The foregoing and other features and advantages will become more apparent from the following detailed description of several embodiments, which proceeds
5 with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE FIGURES

10 **Fig. 1** is a set of diagrams and digital images showing the generation of transgenic mice expressing EGFP in ARC POMC neurons. Fig. 1a is a schematic diagram of the structure of the POMC-EGFP transgene. Fig. 1b is a digital image showing the identification of a single POMC neuron (arrowhead on recording electrode tip) by EGFP fluorescence (upper) and IR-DIC microscopy (lower) in a living ARC slice prior to electrophysiological recordings. Fig. 1c is a set of digital images showing the co-localization (bright, on right) of EGFP (left) and β -endorphin immunoreactivity (middle) in ARC POMC neurons. Scale bars: b & c, 50 μ m. Fig. 1d is a set of diagrams showing the distribution of EGFP-positive neuronal soma throughout the ARC nucleus. \circ = 5 cells, \bullet = 10 cells.

20 **Fig. 2** is a tracing and graphs showing activation of MOP-Rs hyperpolarizes the EGFP-labeled POMC neurons by opening G protein-coupled inwardly-rectifying potassium channels. Fig. 2a is a tracing showing met-enkephalin hyperpolarizes POMC neurons and inhibits all action potentials. The horizontal bar indicates the time when 30 μ M Met-Enk was bath-applied to the slice. Fig. 2b is a graph showing met-enkephalin current and reversal potential is shifted by extracellular K⁺ concentration. Fig. 2c is a graph showing met-enkephalin activates MOP-Rs on POMC neurons. A Met-Enk (30 μ M) current was observed and the MOP-R specific antagonist CTAP (1 μ M) was applied for 1 minute. Following CTAP Met-Enk
25 elicited no current. The figure is representative of three experiments.

30

-5-

Fig. 3 are tracings and graphs demonstrating that leptin depolarizes POMC neurons via a non-specific cation channel, and decreases GABAergic tone onto POMC cells. Fig. 3a is a tracing demonstrating that leptin depolarizes POMC neurons and increases the frequency of action potentials within 1 to 10 minutes of 5 addition. The figure is a representative example of recordings made from 77 POMC neurons. Fig. 3b is a graph showing that leptin causes a concentration dependent depolarization of POMC cells. The depolarization caused by leptin was determined at 0.1, 1, 10, 50, and 100 nM ($EC_{50} = 5.9$ nM) in (8, 7, 9, 3, 45) cells respectively. Fig. 3c is a graph showing that leptin depolarizes POMC cells by activating a 10 nonspecific cation current. The figure is representative of the response in 10 cells. Fig. 3d is a graph showing that leptin decreases the frequency of IPSCs in POMC cells. The figure is an example of 5 cells in which leptin (100 nM) decreased the frequency of IPSCs. Fig. 3e is a tracing demonstrating that leptin had no effect on 5 adjacent non-fluorescent ARC neurons. Fig. 3f is a tracing showing that leptin 15 hyperpolarized 5 non-fluorescent ARC neurons.

Fig. 4 is a set of images showing that the GABAergic inputs to POMC cells are from NPY neurons that co-express GABA. Fig. 4a is a graph showing that NPY 20 decreases the frequency of mini IPSCs in POMC neurons. Fig. 4b is a graph demonstrating that D-Trp⁸- γ MSH (7nM), a dose that selectively activates MC3-R, increases the frequency of GABAergic IPSCs in POMC neurons. Fig. 4c is a tracing showing that D-Trp⁸- γ MSH hyperpolarizes POMC neurons. Figs. 4a, 4b and 4c are representative. Fig. 4d is a set of digital images demonstrating that expression of NPY in nerve terminals adjacent to POMC neurons in the ARC. NPY nerve 25 terminals (black, arrowheads); POMC neuronal soma (grey). Scale bar, 10 μ m. Fig. 4e is a digital image showing expression of GABA and NPY in nerve terminals synapsing onto POMC neurons in the ARC. GABA immunoreactivity (10 nm gold particles, arrowheads without tail) and NPY immunoreactivity (25 nm gold particles, arrows with tail) are in separate vesicle populations co-localized within synaptic 30 boutons that make direct contact with the soma of POMC neurons (DAB contrasted with uranyl acetate and lead citrate, diffuse black in cytoplasm). Scale bar, 1 μ m. Fig. 4f is a diagram of the model of NPY/GABA and POMC neurons in the ARC.

Fig. 5 is a set of graphs relating to the feeding response to PYY₃₋₃₆ in rats.

Fig. 5a is a bar graph of dark-phase feeding tabulating food intake after intraperitoneal injection of PYY₃₋₃₆. Freely feeding rats were injected with PYY₃₋₃₆ at the doses indicated (μg/100g), or saline, just prior to 'lights off' and 4-hour cumulative food intake was measured. Results are the mean ± s.e.m. (n = 8 per group), * = p < 0.05, ** = p < 0.01, *** = < 0.001 compared to saline. Fig. 5b is a bar graph of food intake after intraperitoneal injection of PYY₃₋₃₆. Fasted rats were injected with PYY₃₋₃₆ at the doses indicated (μg/100g), or saline, and 4-hour cumulative food intake was measured. Results are shown as the mean ± s.e.m. (n = 8 per group), * = p < 0.05, ** = p < 0.01, *** = < 0.001 compared to saline. Fig. 5c is a bar graph of cumulative food intake after intraperitoneal injection of saline or PYY₃₋₃₆. Fasted rats were injected with either saline (closed bars) or PYY₃₋₃₆ 5μg/100g (open bars) and cumulative food intake measured at the time points indicated. Results are expressed as mean ± s.e.m. (n = 12 per group), ** = p < 0.01 compared to saline. Fig. 5d is a line graph of body weight gain during chronic treatment with PYY₃₋₃₆. Rats were injected intraperitoneally with PYY₃₋₃₆ 5μg/100g (open squares) or saline (filled inverted triangles) twice daily for 7 days. Body weight gain was calculated each day. Results are expressed as mean ± s.e.m. (n = 12 per group) ** = p < 0.01 compared to saline.

Fig. 6 is a set of digital images of c-fos expression in *Pomc-EGFP* mice.

Figs. 6a and 6b are digital images of representative sections (bregma -1.4 mm²²) of c-fos expression in the arcuate nucleus of *Pomc-EGFP* mice response to 25 intraperitoneal saline (Fig. 6a) or PYY₃₋₃₆ (5μg/100g) (Fig. 6b). Scale bar 100 μm. 3V, third ventricle; Arc, arcuate nucleus. Figs. 6c and 6d are digital images of representative sections showing POMC-EGFP neurons (Fig. 6c) and c-fos immunoreactivity (Fig. 6d) either co-localizing (bright arrows) or alone (single darker arrow). Scale bar 25 μm.

30

Fig. 7 is a set of bar graphs relating to intra-arcuate PYY₃₋₃₆ in rats and feeding effects of IP PYY₃₋₃₆ in *Y2r*- null mice. Fig. 7a is a bar graph of food intake

following intra-arcuate PYY₃₋₃₆ injection. Fasted rats were injected with saline or PYY₃₋₃₆ into the arcuate nucleus at the doses indicated. Post-injection 2-hour food intake was measured, ** = p < 0.01 compared to saline. Figs. 7b and 7c are bar graphs of feeding response to PYY₃₋₃₆ in Y2r-null mice following IP administration: 5 wild type littermates mice (Fig. 7b) and Y2r-null mice (Fig. 7c), fasted for 24 hours, were injected with PYY₃₋₃₆ at the doses indicated (μ g/100g), or saline, and 4-hour cumulative food intake was measured. Results are the mean \pm s.e.m. (n = 5 per group), * = p < 0.05, ** = p < 0.01 compared to saline.

Fig. 8 is a set of images relating to the electrophysiological and neuropeptide responses to PYY₃₋₃₆ and Y2A. Fig. 8a is a tracing showing the effect of PYY₃₋₃₆ (10 nM) on the frequency of action potentials in POMC neurons (whole-cell configuration recordings; n = 22) * p < 0.05. PYY₃₋₃₈ was administered at time D for 3 minutes; baseline, -3 to 0 minute; PYY₃₋₃₆, 2-5 minutes; and wash-out, 8-11 minutes. Inset shows a representative recording of membrane potential and action potential frequency. Fig. 8b is a graph of the effect if PYY₃₋₃₈ (10nM) on the frequency of action potentials in loose cell-attached patch recordings (n=8). Data from individual cells were normalized to the firing rate for the 200s before PYY₃₋₃₈ addition. Fig. 8c is a tracing and a graph of the effect of PYY₃₋₃₆ (50nM) on spontaneous IPSCs onto POMC neurons (n=13). Inset shows a representative recording of IPSCs before and after PYY₃₋₃₆ (50nM), respectively. Results in Fig. 20 8a-8c are expressed as mean \pm s.e.m. Fig. 8d and 8e are bar graphs showing NPY (Fig. 8d) and %-MSH (Fig. 8e) released from hypothalamic explants in response to Y2A. Hypothalamic slices were incubated with artificial CSF (aCSF), with or without 50nM Y2A, for 45 minutes. Results are expressed as mean \pm s.e.m. (n=40); 25 ** = p < 0.01; *** = p < 0.001 compared to saline.

Fig. 9 is a set of graphs showing the effect of PYY₃₋₃₆ infusion on appetite and food intake in human subjects. Fig. 9a is a graph of the calorie intake from a “free-choice” buffet meal 2 hours after infusion with saline or PYY₃₋₃₆. The thin lines indicate individual changes in calorie intake for each subject between saline 30 and PYY₃₋₃₆ administration. The thick line represents mean change between the two infusions (n = 12). Fig. 9b is a graph of the 24-hour calorie intake following

-8-

infusion with saline or PYY₃₋₃₆. Total calorie intake, as assessed by food diaries, is shown for the 24-hour period following either saline or PYY₃₋₃₆ infusion. Data is given as mean \pm s.e.m. (n = 12), *** = p < 0.0001 compared to saline. Fig. 9c is a graph of the appetite score (relative scale). Visual analogue scores (Raben et al., *Br. J. Nutr.* 73, 517-30, 1995) show perceived hunger during and after infusions. The results are presented as change from baseline scores and are the mean \pm s.e.m. for all 12 subjects.

5 **Fig. 10** shows plasma PYY₃₋₃₆ levels following subcutaneous administration of 10 nmols PYY₃₋₃₆ (broken line) and 20 nmols PYY₃₋₃₆ (solid line).

10 **Fig. 11** shows the effect on 1-hour food intake in non-fasted rats of IP administration of GLP-1 in the presence or absence of concomitant exendin 9-39 in the arcuate nucleus of the rat. S = Saline, G = GLP-1 (25 nmol/kg), and Ex = exendin 9-39 (20 nmoles/kg).

15 **Fig. 12** shows the effects of co-administration of PYY₃₋₃₆ and GLP-1 on 1-hour food intake in non-fasted rats injected (intraperitoneally) prior to the onset of the dark phase. IP GLP-1 (30 μ g/kg = 1xG or 60 μ g/kg = 2xG in 500 μ l saline) or IP PYY₃₋₃₆ (3 μ g/kg = 1xP or 6 μ g/kg = 2xP in 500 μ l saline). Co-administration (P+G group) is GLP-1 30 μ g/kg and PYY₃₋₃₆ 3 μ g/kg combined.

20 **SEQUENCE LISTING**

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each 25 nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand.

DETAILED DESCRIPTION

30 **I. Abbreviations**

α -MSH: alpha melanocortin stimulating hormone

Arc: arcuate nucleus
EPSP: excitatory postsynaptic potential
GABA: γ aminobutyric acid
GFP, EGFP: green fluorescent protein
5 **GLP-1:** glucagons-like peptide-1
ICV: intracerebroventricular
IP: intraperitoneal
IPSC: inhibitory postsynaptic current
kb: kilobase
10 **kg:** kilogram
MOP-R: μ -opioid receptor
MV: millivolts
NPY: neuropeptide Y
Oxm: oxyntomodulin
15 **pmol:** picomole
POMC: proopiomelanocortin
RIA: radioimmunoassay
RPA: RNase protection assay
s.e.m: standard error of the mean
20 **TH:** tyrosine hydroxylase
 μ M: micromolar
V: volts
Y2A: N-acetyl (Leu²⁸, Leu³¹) NPY (24-36)

25 **II. Terms**
Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew et al. (eds.), *The Encyclopedia of Molecular Biology*, published 30 by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

-10-

In order to facilitate review of the various embodiments of this disclosure, the following explanations of specific terms are provided:

Action potential: A rapidly propagated electrical message that speeds along an axon of a neuron and over the surface membrane of many muscle and glandular cells. In axons they are brief, travel at constant velocity, and maintain a constant amplitude. Like all electrical messages of the central nervous system, the action potential is a membrane potential change caused by the flow of ions through ion channels in the membrane. In one embodiment, an action potential is a regenerative wave of sodium permeability.

Animal: Living multi-cellular vertebrate organisms, a category that includes, for example, mammals and birds. The term mammal includes both human and non-human mammals. Similarly, the term "subject" includes both human and veterinary subjects.

Anorexia: A lack or loss of the appetite for food. In one embodiment, anorexia is a result of "anorexia nervosa." This is an eating disorder primarily affecting females, usually with onset in adolescence, characterized by refusal to maintain a normal minimal body weight, intense fear of gaining weight or becoming obese, and a disturbance of body image resulting in a feeling of being fat or having fat in certain areas even when extremely emaciated, undue reliance on body weight or shape for self-evaluation, and amenorrhea. Associated features often include denial of the illness and resistance to psychotherapy, depressive symptoms, markedly decreased libido, and obsessions or peculiar behavior regarding food, such as hoarding. The disorder is divided into two subtypes, a restricting type, in which weight loss is achieved primarily through diet or exercise, and a binge-eating/purging type, in which binge eating or purging behavior also occur regularly.

Antagonist: A substance that tends to nullify the action of another, as an agent that binds to a cell receptor without eliciting a biological response, blocking binding of substances that could elicit such responses.

Appetite: A natural desire, or longing for food. In one embodiment, appetite is measured by a survey to assess the desire for food. Increased appetite generally leads to increased feeding behavior.

Appetite Suppressants: Compounds that decrease the desire for food.

Commercially available appetite suppressants include, but are not limited to, amfepramone (diethylpropion), phentermine, mazindol and phenylpropanolamine, fenfluramine, dexfenfluramine, and fluoxetine.

5 **Binding:** A specific interaction between two molecules, such that the two molecules interact. Binding can be specific and selective, so that one molecule is bound preferentially when compared to another molecule. In one embodiment, specific binding is identified by a disassociation constant (K_d).

10 **Body Mass Index (BMI):** A mathematical formula for measuring body mass, also sometimes called Quetelet's Index. BMI is calculated by dividing weight (in kg) by height² (in meters²). The current standards for both men and women accepted as "normal" are a BMI of 20-24.9 kg/m². In one embodiment, a BMI of greater than 25 kg/m² can be used to identify an obese subject. Grade I obesity corresponds to a BMI of 25-29.9 kg/m². Grade II obesity corresponds to a BMI of 15 30-40 kg/m²; and Grade III obesity corresponds to a BMI greater than 40 kg/m² (Jequier, *Am. J Clin. Nutr.* 45:1035-47, 1987). Ideal body weight will vary among species and individuals based on height, body build, bone structure, and sex.

20 **c-fos:** The cellular homologue of the viral v-fos oncogene found in FBJ (Finkel-Biskis-Jinkins) and FBR murine osteosarcoma viruses (MSV). The human fos gene maps to chromosome 14q21-q31. Human fos has been identified as TIS-28.

25 C-fos is thought to have an important role in signal transduction, cell proliferation, and differentiation. It is a nuclear protein which, in combination with other transcription factors (for example, jun) acts as a trans-activating regulator of gene expression. C-fos is an immediate early response gene, which are believed to play a key role in the early response of cells to growth factors. C-fos is involved also in the control of cell growth and differentiation of embryonic hematopoietic cells and neuronal cells. The human c-fos coding amino acid and nucleic sequences are known (e.g., see Verma et al., *Cold Spring Harb. Symp. Quant. Biol.* 51, 949, 30 1986; GenBank Accession Nos. K00650 and M16287, and is available on the internet).

-12-

Caloric intake or calorie intake: The number of calories (energy) consumed by an individual.

Calorie: A unit of measurement in food. A standard calorie is defined as 4.184 absolute joules, or the amount of energy it takes to raise the temperature of 5 one gram of water from 15 to 16° C (or 1/100th the amount of energy needed to raise the temperature of one gram of water at one atmosphere pressure from 0° C to 100° C), food calories are actually equal to 1,000 standard calories (1 food calorie = 1 kilocalorie).

Conservative variation: The replacement of an amino acid residue by 10 another, biologically similar residue. Examples of conservative variations include the substitution of one hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acid, or glutamine for asparagine, and the like. The term "conservative variation" also includes the use of a 15 substituted amino acid in place of an unsubstituted parent amino acid provided that antibodies raised to the substituted polypeptide also immunoreact with the unsubstituted polypeptide.

Non-limiting examples of conservative amino acid substitutions include those listed below:

20

	Original Residue	Conservative Substitutions
	Ala	Ser
	Arg	Lys
25	Asn	Gln, His
	Asp	Glu
	Cys	Ser
	Gln	Asn
	Glu	Asp
30	His	Asn; Gln
	Ile	Leu, Val
	Leu	Ile; Val
	Lys	Arg; Gln; Glu
	Met	Leu; Ile
35	Phe	Met; Leu; Tyr
	Ser	Thr
	Thr	Ser
	Trp	Tyr

-13-

Tyr
Val

Trp; Phe
Ile; Leu

Depolarization: An increase in the membrane potential of a cell. Certain stimuli reduce the charge across the plasma membrane. These can be electrical stimuli (which open voltage-gated channels), mechanical stimuli (which activate mechanically-gated channels) or certain neurotransmitters (which open ligand-gated channels). In each case, the facilitated diffusion of sodium into the cell increases the resting potential at that spot on the cell creating an excitatory postsynaptic potential (EPSP). Depolarizations can also be generated by decreasing the frequency of inhibitory postsynaptic currents (IPSCs), these are due to inhibitory neurotransmitters facilitating the influx of chloride ions into the cell, creating an IPSC. If the potential is increased to the threshold voltage (about -50 mV in mammalian neurons), an action potential is generated in the cell.

15 Diabetes: A failure of cells to transport endogenous glucose across their membranes either because of an endogenous deficiency of insulin and/or a defect in insulin sensitivity. Diabetes is a chronic syndrome of impaired carbohydrate, protein, and fat metabolism owing to insufficient secretion of insulin or to target tissue insulin resistance. It occurs in two major forms: insulin-dependent diabetes mellitus (IDDM, type I) and non-insulin dependent diabetes mellitus (NIDDM, type II) which differ in etiology, pathology, genetics, age of onset, and treatment.

The two major forms of diabetes are both characterized by an inability to deliver insulin in an amount and with the precise timing that is needed for control of glucose homeostasis. Diabetes type I, or insulin dependent diabetes mellitus (IDDM) is caused by the destruction of β cells, which results in insufficient levels of endogenous insulin. Diabetes type II, or non-insulin dependent diabetes, results from a defect in both the body's sensitivity to insulin, and a relative deficiency in insulin production.

30 Exendin: A 39-amino acid peptide isolated from the salivary glands of the Gila monster (*Heloderma suspectum*) (Eng J et al J Biol Chem 267:7402-7405, 1992), see SEQ ID NO:341. Exendin is an example of an agonist at the GLP-1 receptor. Molecules derived from exendin-4 and that also have GLP-1 agonist activity are further examples of GLP-1 agonists.

Food intake: The amount of food consumed by an individual. Food intake can be measured by volume or by weight. In one embodiment, food intake is the total amount of food consumed by an individual. In another embodiment, food intake is the amount of proteins, fat, carbohydrates, cholesterol, vitamins, minerals, 5 or any other food component, of the individual. "Protein intake" refers to the amount of protein consumed by an individual. Similarly, "fat intake," "carbohydrate intake," "cholesterol intake," "vitamin intake," and "mineral intake" refer to the amount of proteins, fat, carbohydrates, cholesterol, vitamins, or minerals consumed by an individual.

10 **Glucagon-like peptide-1:** Peptides produced by processing of preproglucagon, which is a 160 amino acid polypeptide, in the central nervous system (CNS) and the intestine. GLP-1 (1-37) is the initial product. GLP-1 (1-37) is amidated by post-translational processing to yield GLP-1 (1-36) NH, or is enzymatically processed to give GLP-1 (7-37) (SEQ ID NO: 338). GLP-1 (7-37) 15 can be amidated to give GLP-1 (7-36) amide (SEQ ID NO: 339), which is the most biologically active form. GLP-1 is released into the circulation in response to nutrient intake. Intestinal cells secrete GLP-1 (7-37) and GLP-1 (7-36) amide in a ratio of 1 to 5.

20 GLP-1 receptors are found in the brainstem, arcuate nucleus and paraventricular nucleus. Physiological actions of GLP-1 in man include stimulation of insulin release, suppression of gastric acid secretion and slowing of gastric emptying.

25 A GLP-1 agonist is a peptide, small molecule, or chemical compound that preferentially binds to the GLP-1 receptor and stimulate the same biological activity as GLP-1. In one embodiment, an agonist for the GLP-1 receptor binds to the receptor with an equal or greater affinity than a GLP-1 peptide. In another embodiment, an agonist selectively binds the GLP-1 receptor, as compared to binding to another receptor.

30 One of skill in the art can readily determine the dissociation constant (K_d) value of a given compound. This value is dependent on the selectivity of the compound tested. For example, a compound with a K_d which is less than 10 nM is generally considered an excellent drug candidate. However, a compound that has a

-15-

lower affinity, but is selective for the particular receptor, can also be a good drug candidate. In one specific, non-limiting example, an assay, such as a competition assay, is used to determine if a compound of interest is a GLP-1 receptor agonist.

GLP-1 agonists include GLP-1 related peptides and peptides that result from 5 natural or synthetic enzymatic or chemical processing of a GLP-1 peptide or a related peptide.

Hyperpolarization: A decrease in the membrane potential of a cell. Inhibitory neurotransmitters inhibit the transmission of nerve impulses via 10 hyperpolarization. This hyperpolarization is called an inhibitory postsynaptic potential (IPSP). Although the threshold voltage of the cell is unchanged, a hyperpolarized cell requires a stronger excitatory stimulus to reach threshold.

Inhibitory Postsynaptic Current: A current that inhibits an 15 electrophysiological parameter of a postsynaptic cell. The potential of a postsynaptic cell can be analyzed to determine an effect on a presynaptic cell. In one embodiment, the postsynaptic cell is held in voltage clamp mode, and postsynaptic currents are recorded. If necessary, antagonists of other classes of current can be added. In one specific, non-limiting example, to record GABAergic IPSCs, blockers of excitatory channels or receptors can be added. The instantaneous frequency over time is then determined.

20 In one embodiment, IPSCs give a measure of the frequency of GABA release from an NPY neuron. Thus, as NPY neurons release GABA onto POMC neurons, measurement of IPSC frequency is a gauge of the inhibitory tone that POMC neurons are receiving, and can be used to assess the effect of an agonist of PYY.

Membrane potential: The electrical potential of the interior of the cell with 25 respect to the environment, such as an external bath solution. One of skill in the art can readily assess the membrane potential of a cell, such as by using conventional whole cell techniques. Activation of a cell is associated with less negative membrane potentials (for example shifts from about -50 mV to about -40 mV).- These changes in potential increase the likelihood of action potentials, and thus lead 30 to an increase in the rate of action potentials.

The rate of action potentials can be assessed using many approaches, such as using conventional whole cell access, or using, for example, perforated-patch whole-

cell and cell-attached configurations. In each event the absolute voltage or current is not assessed, rather the frequency of rapid deflections characteristic of action potentials is assessed, as a function of time (therefore this frequency is an instantaneous frequency, reported in "bins"). This time component can be related to 5 the time at which a compound, such as a PYY agonist, is applied to the bath to analyze the effect of the compound, such as the PYY agonist, on action potential firing rate.

10 **Neuropeptide Y (NPY):** A 36-amino acid peptide that is a neuropeptide identified in the mammalian brain. NPY is believed to be an important regulator in both the central and peripheral nervous systems and influences a diverse range of physiological parameters, including effects on psychomotor activity, food intake, central endocrine secretion, and vasoactivity in the cardiovascular system. High concentrations of NPY are found in the sympathetic nerves supplying the coronary, cerebral, and renal vasculature and have contributed to vasoconstriction. NPY 15 binding sites have been identified in a variety of tissues, including spleen, intestinal membranes, brain, aortic smooth muscle, kidney, testis, and placenta. In addition, binding sites have been reported in a number of rat and human cell lines.

20 NPY receptor has structure/activity relationships within the pancreatic polypeptide family. This family includes NPY, which is synthesized primarily in neurons; peptide YY (PYY), which is synthesized primarily by endocrine cells in the gut; and pancreatic polypeptide (PP), which is synthesized primarily by endocrine cells in the pancreas. These 36 amino acid peptides have a compact helical structure involving an amino acid structure, termed a "PP-fold" in the middle of the peptide.

25 NPY binds to several receptors, including the Y1, Y2, Y3, Y4 (PP), Y5, Y6, and Y7 receptors. These receptors are recognized based on binding affinities, pharmacology, and sequence (if known). Most, if not all of these receptors are G protein coupled receptors. The Y1 receptor is generally considered to be postsynaptic and mediates many of the known actions of neuropeptide Y in the 30 periphery. Originally, this receptor was described as having poor affinity for C-terminal fragments of neuropeptide Y, such as the 13-36 fragment, but interacts with

-17-

the full length neuropeptide Y and peptide YY with equal affinity (e.g., see PCT publication WO 93/09227).

Pharmacologically, the Y2 receptor is distinguished from Y1 by exhibiting affinity for C-terminal fragments of neuropeptide Y. The Y2 receptor is most often 5 differentiated by the affinity of neuropeptide Y(13-36), although the 3-36 fragment of neuropeptide Y and peptide YY provides improved affinity and selectivity (see Dumont et al., *Society for Neuroscience Abstracts* 19:726, 1993). Signal transmission through both the Y1 and the Y2 receptors are coupled to the inhibition of adenylate cyclase. Binding to the Y-2 receptor was also found to reduce the 10 intracellular levels of calcium in the synapse by selective inhibition of N-type calcium channels. In addition, the Y-2 receptor, like the Y1 receptors, exhibits differential coupling to second messengers (see U.S. Patent No. 6,355,478). Y2 receptors are found in a variety of brain regions, including the hippocampus, substantia nigra-lateralis, thalamus, hypothalamus, and brainstem. The human, 15 murine, monkey and rat Y2 receptors have been cloned (e.g., see U.S. Patent No. 6,420,352 and U.S. Patent No. 6,355,478).

A Y2 receptor agonist is a peptide, small molecule, or chemical compound that preferentially binds to the Y2 receptor and stimulates intracellular signaling. In one embodiment, an agonist for the Y2 receptor binds to the receptor with an equal 20 or greater affinity than NPY. In another embodiment, an agonist selectively binds the Y2 receptor, as compared to binding to another receptor.

One of skill in the art can readily determine the dissociation constant (K_d) value of a given compound. This value is dependent on the selectivity of the compound tested. For example, a compound with a K_d which is less than 10 nM is 25 generally considered an excellent drug candidate. However, a compound that has a lower affinity, but is selective for the particular receptor, can also be a good drug candidate. In one specific, non-limiting example, an assay, such as a competition assay, is used to determine if a compound of interest is a Y2 receptor agonist.

Assays useful for evaluating neuropeptide Y receptor antagonists are also well 30 known in the art (see U.S. Patent No. 5,284,839, which is herein incorporated by reference, and Walker et al., *Journal of Neurosciences* 8:2438-2446, 1988).

-18-

Normal Daily Diet: The average food intake for an individual of a given species. A normal daily diet can be expressed in terms of caloric intake, protein intake, carbohydrate intake, and/or fat intake. A normal daily diet in humans generally comprises the following: about 2,000, about 2,400, or about 2,800 to 5 significantly more calories. In addition, a normal daily diet in humans generally includes about 12 g to about 45 g of protein, about 120 g to about 610 g of carbohydrate, and about 11 g to about 90 g of fat. A low calorie diet would be no more than about 85%, and preferably no more than about 70%, of the normal caloric intake of a human individual.

10 In animals, the caloric and nutrient requirements vary depending on the species and size of the animal. For example, in cats, the total caloric intake per pound, as well as the percent distribution of protein, carbohydrate and fat varies with the age of the cat and the reproductive state. A general guideline for cats, however, is 40 cal/lb/day (18.2 cal/kg/day). About 30% to about 40% should be protein, 15 about 7% to about 10% should be from carbohydrate, and about 50% to about 62.5% should be derived from fat intake. One of skill in the art can readily identify the normal daily diet of an individual of any species.

Obesity: A condition in which excess body fat may put a person at health risk (see Barlow and Dietz, *Pediatrics* 102:E29, 1998; National Institutes of Health, 20 National Heart, Lung, and Blood Institute (NHLBI), *Obes. Res.* 6 (suppl. 2):51S-209S, 1998). Excess body fat is a result of an imbalance of energy intake and energy expenditure. In one embodiment, the Body Mass Index (BMI) is used to assess obesity. In one embodiment, a BMI of 25.0 kg/m^2 to 29.9 kg/m^2 is overweight, while a BMI of 30 kg/m^2 is obese.

25 In another embodiment, waist circumference is used to assess obesity. In this embodiment, in men a waist circumference of 102 cm or more is considered obese, while in women a waist circumference of 89 cm or more is considered obese. Strong evidence shows that obesity affects both the morbidity and mortality of individuals. For example, an obese individual is at increased risk for heart disease, 30 non-insulin dependent (type 2) diabetes, hypertension, stroke, cancer (e.g. endometrial, breast, prostate, and colon cancer), dyslipidemia, gall bladder disease,

sleep apnea, reduced fertility, and osteoarthritis, amongst others (see Lyznicki et al., *Am. Fam. Phys.* 63:2185, 2001).

Overweight: An individual who weighs more than their ideal body weight. An overweight individual can be obese, but is not necessarily obese. In one embodiment, an overweight individual is any individual who desires to decrease their weight. In another embodiment, an overweight individual is an individual with a BMI of 25.0 kg/m² to 29.9 kg/m²

Oxyntomodulin: A further peptide produced by post-translational processing of preproglucagon in the CNS and intestine, see SEQ ID NO: 340..

10 Agonists of oxyntomodulin may be identified as described above for agonists of GLP-1 and of NPY. include oxyntomodulins of other species and also modified sequences, for example, sequences in which The term OXM used in this text also covers any analogue of the above OXM sequence, wherein the histidine residue at position 1 is maintained or replaced by an aromatic moiety carrying a positive charge or a derivative thereof, preferably wherein the moiety is an amino acid, more preferably wherein it is a histidine derivative, while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 or 22 of the other amino acids in the above OXM sequence can be independently replaced by any other independently chosen amino acid, with the exception of histidine in position 1.

15 20 Any one or more (to 22) other alpha-amino acid residue in the sequence can be independently replaced by any other one alpha-amino acid residue. Preferably, any amino acid residue other than histidine is replaced with a conservative replacement as well known in the art i.e. replacing an amino acid with one of a similar chemical type such as replacing one hydrophobic amino acid with another.

25 30 As discussed above, 1 to 22 of the amino acids can be replaced. In addition to the replacement option above, this may be by a non-essential or modified or isomeric form of an amino acid. For example, 1 to 22 amino acids can be replaced by an isomeric form (for example a D-amino acid), or a modified amino acid, for example a nor-amino acid (such as norleucine or norvaline) or a non-essential amino acid (such as taurine). Furthermore, 1 to 22 amino acids may be replaced by a corresponding or different amino acid linked via its side chain (for example gamma-

-20-

linked glutamic acid). For each of the replacements discussed above, the histidine residue at position 1 is unaltered or defined above.

In addition, 1, 2, 3, 4 or 5 of the amino acid residues can be removed from the OXM sequence with the exception of histidine at the 1 position (or as defined 5 above). The deleted residues may be any 2, 3, 4 or 5 contiguous residues or entirely separate residues.

The C-terminus of the OXM sequence may be modified to add further amino acid residues or other moieties.

10 **Pancreatic Polypeptide:** A 36 amino acid peptide produced by the pancreas that is has homology to PYY and NPY.

15 **Peripheral Administration:** Administration outside of the central nervous system. Peripheral administration does not include direct administration to the brain. Peripheral administration includes, but is not limited to intravascular, intramuscular, subcutaneous, inhalation, oral, rectal, transdermal or intra-nasal administration

20 **Polypeptide:** A polymer in which the monomers are amino acid residues which are joined together through amide bonds. When the amino acids are alpha-amino acids, either the L-optical isomer or the D-optical isomer can be used, the L-isomers being preferred. The terms "polypeptide" or "protein" as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term "polypeptide" is specifically intended to cover naturally occurring proteins, as well as those which are recombinantly or synthetically produced. The term "polypeptide fragment" refers to a portion of a polypeptide, for example such a fragment which exhibits at least one useful 25 sequence in binding a receptor. The term "functional fragments of a polypeptide" refers to all fragments of a polypeptide that retain an activity of the polypeptide. Biologically functional peptides can also include fusion proteins, in which the peptide of interest has been fused to another peptide that does not decrease its desired activity.

30 **PYY:** A peptide YY polypeptide obtained or derived from any species.

Thus, PYY includes the human full length polypeptide (as set forth in SEQ ID NO: 1) and species variations of PYY, including e.g. murine, hamster, chicken, bovine,

-21-

rat, and dog PYY (SEQ ID NOS: 5-12). In one embodiment, PYY agonists do not include NPY. PYY also includes PYY₃₋₃₆. A "PYY agonist" is any compound which binds to a receptor that specifically binds PYY, and elicits an effect of PYY. In one embodiment, a PYY agonist is a compound that affects food intake, caloric intake, or appetite, and/or which binds specifically in a Y receptor assay or competes for binding with PYY, such as in a competitive binding assay with labeled PYY. PYY agonists include, but are not limited to, compounds that bind to the Y2 receptor.

Substantially purified: A polypeptide which is substantially free of other proteins, lipids, carbohydrates or other materials with which it is naturally associated. For example, the polypeptide may be at least 50%, 80% or 90% free of other proteins, lipids, carbohydrates or other materials with which it is naturally associated.

Therapeutically effective amount: A dose sufficient to prevent advancement, or to cause regression of a disorder, or which is capable of relieving a sign or symptom of a disorder, or which is capable of achieving a desired result. In several embodiments, a therapeutically effect of PYY or an agonist thereof is an amount sufficient to inhibit or halt weight gain, or an amount sufficient to decrease appetite, or an amount sufficient to reduce caloric intake or food intake or increase energy expenditure.

Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of this disclosure, suitable methods and materials are described below. The term "comprises" means "includes." All publications, patent applications, patents, and

other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

5

**Methods for Altering Food Intake, Appetite, Caloric Intake
and Energy Expenditure**

A method is disclosed herein for reducing food intake by peripherally administering to a subject a therapeutically effective amount of PYY or an agonist of PYY and GLP-1 or an agonist thereof. In one embodiment, administration of PYY, or an agonist of PYY, and GLP-1 or an agonist thereof, results in a decrease in the amount, either the total weight or the total volume of food. In other embodiment, administration of PYY, or an agonist thereof, and GLP-1 or an agonist thereof, results in a decrease of the intake of a food component, such as a decrease in the 10 ingestion of lipids, carbohydrates, cholesterol, or proteins. In the any of the methods disclosed herein, a preferred compound, PYY₃₋₃₆ can be administered. This disclosure includes the corresponding uses of PYY or an agonist thereof and GLP-1 or an agonist thereof, for the manufacture of a medicament or medicaments for the 15 purposes set herein, and includes the use of PYY₃₋₃₆.

A method is also disclosed herein for reducing caloric intake by peripherally administering to a subject a therapeutically effective amount of PYY or an agonist of PYY and GLP-1 or an agonist thereof. In one embodiment, total caloric intake is 20 reduced by peripheral administration of a therapeutically effective amount of PYY and GLP-1 or an agonist thereof. In other embodiments, the caloric intake from the 25 ingestion of a specific food component, such as, but not limited to, the ingestion of lipids, carbohydrates, cholesterol, or proteins, is reduced.

In an additional embodiment, a method is disclosed herein for reducing appetite by administering a therapeutically effective amount of PYY or an agonist thereof and GLP-1 or an agonist thereof. Appetite can be measured by any means 30 known to one of skill in the art. For example, decreased appetite can be assessed by a psychological assessment. In this embodiment, administration of PYY and the other agent(s) results in a change in perceived hunger, satiety, and/or fullness.

-23-

Hunger can be assessed by any means known to one of skill in the art. In one embodiment, hunger is assessed using psychological assays, such as by an assessment of hunger feelings and sensory perception using a questionnaire, such as, but not limited to, a Visual Analog Score (VAS) questionnaire (see the Examples section). In one specific, non-limiting example, hunger is assessed by answering questions relating to desire for food, drink, prospective food consumption, nausea, and perceptions relating to smell or taste.

In a further embodiment, a method is disclosed herein for altering energy metabolism in a subject. The method includes peripherally administering a therapeutically effective amount of PYY or an agonist thereof and GLP-1 or an agonist thereof, to the subject, thereby altering energy expenditure. Energy is burned in all physiological processes. The body can alter the rate of energy expenditure directly, by modulating the efficiency of those processes, or changing the number and nature of processes that are occurring. For example, during digestion the body expends energy moving food through the bowel, and digesting food, and within cells, the efficiency of cellular metabolism can be altered to produce more or less heat. In a further embodiment a method is disclosed herein for any and all manipulations of the arcuate circuitry described in this application, that alter food intake coordinately and reciprocally alter energy expenditure. Energy expenditure is a result of cellular metabolism, protein synthesis, metabolic rate, and calorie utilization. Thus, in this embodiment, peripheral administration of PYY and administration of GLP-1 or an agonist thereof, results in increased energy expenditure, and decreased efficiency of calorie utilization. In one embodiment, a therapeutically effective amount of PYY or an agonist and GLP-1 or an agonist thereof, thereof is administered to a subject, thereby increasing energy expenditure.

In several embodiments, PYY (e.g., PYY₃₋₃₆) or an agonist thereof and GLP-1 or an agonist thereof, are used for weight control and treatment, reduction or prevention of obesity, in particular any one or more of the following: preventing and reducing weight gain; inducing and promoting weight loss; and reducing obesity as measured by the Body Mass Index. The disclosure further relates to the use of PYY or an agonist and GLP-1 or an agonist thereof, in control of any one or more of appetite, satiety and hunger, in particular any one or more of the following:

-24-

reducing, suppressing and inhibiting appetite; inducing, increasing, enhancing and promoting satiety and sensations of satiety; and reducing, inhibiting and suppressing hunger and sensations of hunger. The disclosure further relates to the use of PYY an agonist thereof and GLP-1 or an agonist thereof, in maintaining any one or more of 5 a desired body weight, a desired Body Mass Index, a desired appearance and good health.

The subject can be any subject, including both human and veterinary mammalian subjects. Thus, the subject can be a human, or can be a non-human primate, a farm animal such as swine, cattle, and poultry, a sport animal or pet such 10 as dogs, cats, horses, hamsters, rodents, or a zoo animal such as lions, tigers, or bears.

Obesity is currently a poorly treatable, chronic, essentially intractable metabolic disorder. A therapeutic drug useful in weight reduction of obese persons could have a profound beneficial effect on their health. Thus, the subject can be, but 15 is not limited to, a subject who is overweight or obese. In one embodiment, the subject has, or is at risk of having, a disorder wherein obesity or being overweight is a risk factor for the disorder. Disorders of interest include, but are not limited to, cardiovascular disease, (including, but not limited to, hypertension, atherosclerosis, congestive heart failure, and dyslipidemia), stroke, gallbladder disease, 20 osteoarthritis, sleep apnea, reproductive disorders such as, but not limited to, polycystic ovarian syndrome, cancers (e.g., breast, prostate, colon, endometrial, kidney, and esophagus cancer), varicose veins, acanthosis nigricans, eczema, exercise intolerance, insulin resistance, hypertension hypercholesterolemia, cholithiasis, osteoarthritis, orthopedic injury, insulin resistance (such as, but not limited to, type 2 25 diabetes and syndrome X) and tromboembolic disease (see Kopelman, *Nature* 404:635-43; Rissanen et al., *British Med. J.* 301, 835, 1990).

Other associated disorders also include depression, anxiety, panic attacks, migraine headaches, PMS, chronic pain states, fibromyalgia, insomnia, impulsivity, obsessive compulsive disorder, and myoclonus. Obesity is a recognized risk factor 30 for increased incidence of complications of general anesthesia. (See e. g., Kopelman, *Nature* 404:635-43, 2000). It reduces life span and carries a serious risk of co-morbidities listed above.

-25-

Other diseases or disorders associated with obesity are birth defects (maternal obesity associated with increased incidence of neural tube defects), carpal tunnel syndrome (CTS), chronic venous insufficiency (CVT), daytime sleepiness, deep vein thrombosis (DVT), end stage renal disease (ESRD), gout, heat disorders, 5 impaired immune response, impaired respiratory function, infertility, liver disease, lower back pain, obstetric and gynecologic complications, pancreatitis, as well as abdominal hernias, acanthosis nigricans, endocrine abnormalities, chronic hypoxia and hypercapnia, dermatological effects, elephantitis, gastroesophageal reflux, heel spurs, lower extremity edema, mammegaly (causing considerable problems such as 10 bra strap pain, skin damage, cervical pain, chronic odors and infections in the skin folds under the breasts, etc.), large anterior abdominal wall masses (abdominal panniculitis with frequent panniculitis, impeding walking, causing frequent infections, odors, clothing difficulties, low back pain), musculoskeletal disease, pseudo tumor cerebri (or benign intracranial hypertension), and sliding hiatal hernia.

15 The present disclosure relates to treating, prevention, ameliorating or alleviating conditions or disorders caused by, complicated by, or aggravated by a relatively high nutrient availability. By "condition or disorder which can be alleviated by reducing caloric (or nutrient) availability," it is meant any condition or disorder in a subject that is either caused by, complicated by, or aggravated by a 20 relatively high nutrient availability, or that can be alleviated by reducing nutrient availability, for example by decreasing food intake. Subjects who are insulin resistant, glucose intolerant, or have any form of diabetes mellitus (e.g., type 1, 2 or gestational diabetes) can also benefit from this disclosure.

Such conditions or disorders are disorders associated with increased caloric 25 intake, insulin resistance, or glucose intolerance and include, but are not limited to, obesity, diabetes, including type 2 diabetes, eating disorders, insulin-resistance syndromes, and Alzheimer's disease.

In another embodiment, the subject is a subject who desires weight loss, such as female and male subject who desire a change in their appearance. In yet a further 30 embodiment, the subject is a subject who desires decreased feelings of hunger, such as, but not limited to, a person involved in a lengthy task that requires a high level of

concentration (e.g., soldiers on active duty, air traffic controllers, or truck drivers on long distance routes, etc.).

A suitable administration format may be best determined by the subject or by a medical practitioner. In one embodiment, the pharmaceutical compositions 5 that include PYY, or an agonist thereof, will preferably be formulated in unit dosage form, suitable for individual administration of precise dosages. An effective amount of PYY or an agonist thereof can be administered in a single dose, or in multiple doses, for example daily, during a course of treatment. In one embodiment, PYY is administered whenever the effect (e.g., appetite suppression, decreased food intake, 10 or decreased caloric intake) is desired. In another embodiment, PYY or an analog thereof is administered slightly prior to whenever the effect is desired, such as, but not limited to about 10 minutes, about 15 minutes, about 30 minutes, about 60 minutes, about 90 minutes, or about 120 minutes, prior to the time the effect is desired. In another embodiment, a time release formulation is utilized.

15 In one embodiment, a therapeutically effective amount of PYY or an agonist thereof is administered as a single pulse dose, as a bolus dose, or as pulse doses administered over time. Thus, in pulse doses, a bolus administration of PYY is provided, followed by a time period wherein no PYY is administered to the subject, followed by a second bolus administration. In specific, non-limiting examples, 20 pulse doses of PYY are administered during the course of a day, during the course of a week, or during the course of a month.

The therapeutically effective amount of PYY or an agonist thereof will be dependent on the molecule utilized, the subject being treated, the severity and type of the affliction, and the manner of administration. For example, a therapeutically 25 effective amount of PYY or an agonist thereof can vary from about 0.01 μ g per kilogram (kg) body weight to about 1 g per kg body weight, such as about 1 μ g to about 5 mg per kg body weight, or about 5 μ g to about 1 mg per kg body weight. In another embodiment, PYY or an agonist thereof is administered to a subject at 0.5 to 135 picomole (pmol) per kg body weight, or about 72 pmol per kg body weight. 30 In one specific, non-limiting example about 5 to about 50 nmol is administered as a subcutaneous injection, such as about 2 to about 20 nmol, or about 10 nmol is administered as a subcutaneous injection. The exact dose is readily determined by

one of skill in the art based on the potency of the specific compound (such as the PYY polypeptide, or agonist) utilized, the age, weight, sex and physiological condition of the subject. The dose of an agonist can be a molar equivalent of the therapeutically effective dose of PYY or PYY₃₋₃₆.

5 The compositions or pharmaceutical compositions can be administered by any route, including intravenous, intraperitoneal, subcutaneous, sublingual, transdermal, intramuscular, oral, topical, transmucosal, or by pulmonary inhalation. Compositions useful in the disclosure may conveniently be provided in the form of formulations suitable for parenteral (including intravenous, intramuscular and subcutaneous), nasal or oral administration. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion. PYY, including PYY₃₋₃₆, an agonist of PYY, or an antagonist of PYY, can be administered subcutaneously. It is well known in the art that subcutaneous injections can be easily self-administered.

10 In some cases, it will be convenient to provide a PYY or a PYY agonist and another food-intake-reducing, plasma glucose-lowering or plasma lipid-altering agent, in a single composition or solution for administration together. In other cases, it may be more advantageous to administer the additional agent separately from said PYY or PYY agonist.

15 A suitable administration format may best be determined by a medical practitioner for each patient individually. Various pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g., *Remington's Pharmaceutical Sciences* by E. W. Martin. See also Wang, Y. J. and Hanson, M. A., *Journal of Parenteral Science and Technology*, Technical Report No. 10, Supp. 42:2S, 1988.

20 PYY and PYY agonists useful in the methods of this disclosure can be provided as parenteral compositions, e.g., for injection or infusion. Preferably, they are suspended in an aqueous carrier, for example, in an isotonic buffer solution at a pH of about 3.0 to about 8.0, preferably at a pH of about 3.5 to about 7.4, 3.5 to 6.0, or 3.5 to about 5.0. Useful buffers include sodium citrate-citric acid and sodium phosphate-phosphoric acid, and sodium acetate/acetic acid buffers. A form of

repository or "depot" slow release preparation may be used so that therapeutically effective amounts of the preparation are delivered into the bloodstream over many hours or days following transdermal injection or delivery.

Since the PYY and agonists are amphoteric, they may be utilized as free bases, as acid addition salts or as metal salts. The salts must, of course, be pharmaceutically acceptable, and these will include metal salts, particularly alkali and alkaline earth metal salts, e.g., potassium or sodium salts. A wide variety of pharmaceutically acceptable acid addition salts are available. Such products are readily prepared by procedures well known to those skilled in the art.

For use by the physician, the compositions can be provided in dosage unit form containing an amount of a PYY or a PYY agonist with or without another active ingredient, e.g., a food intake-reducing, plasma glucose-lowering or plasma lipid-altering agent. Administration may begin whenever the suppression of nutrient availability, food intake, weight, blood glucose or plasma lipid lowering is desired, for example, at the first sign of symptoms of a weight-related disorder or shortly after diagnosis of obesity, diabetes mellitus, or insulin resistance syndrome.

Therapeutically effective amounts of a PYY or a PYY agonist for use in reducing nutrient availability are those that suppress appetite at a desired level. As will be recognized by those in the field, an effective amount of therapeutic agent will vary with many factors including the potency of the particular compound, age and weight of the patient, the patient's physical condition, the blood sugar level, the weight level to be obtained, and other factors. As will be recognized by those in the field, an effective amount of this therapeutic agent will also vary with many factors including the potency of the particular compound, age and weight of the patient, the patient's physical condition, the blood sugar level, the weight level to be obtained, and other factors. Administration may begin whenever the increased of nutrient availability, food intake, weight, blood glucose or plasma lipid lowering is desired, such as, but not limited to, at the first sign of symptoms of a anorexia or at the onset of weight loss due to AIDS.

The optimal formulation and mode of administration of PYY or a PYY agonist to a patient depend on factors known in the art such as the particular disease or disorder, the desired effect, and the type of patient. While the PYY and PYY

agonists will typically be used to treat human subjects they may also be used to treat similar or identical diseases in other vertebrates such as other primates, farm animals such as swine, cattle and poultry, and sport animals and pets such as horses, dogs and cats.

5 As a pharmaceutical medicament the PYY and PYY agonists of the present disclosure may be administered directly by any suitable technique, including parenterally, intranasally, orally, or by absorption through the skin. The specific route of administration of each agent will depend, e.g., on the medical history of the animal.

10 For parenteral administration, in one embodiment, PYY and PYY agonists can be formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the

15 formulation. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to PYY and PYY agonists.

20 Generally, the formulations are prepared by contacting the PYY or PYY agonist uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such

25 as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

PPY and PYY agonists are also suitably administered by sustained-release systems. Suitable examples of sustained-release PYY and PYY agonists include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or mirocapsules), suitable hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt). Sustained-release PPY and PYY agonist compositions may be

-30-

administered orally, rectally, parenterally, intracistemally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), bucally, or as an oral or nasal spray.

Sustained release matrices include polylactides (U.S. Patent No. 3,773,919, 5 EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., *Biopolymers* 22:547-556, 1983, poly(2-hydroxyethyl methacrylate)); (Langer et al., *J. Biomed. Mater. Res.* 15:167-277, 1981; Langer, *Chem. Tech.* 12:98-105, 1982, ethylene vinyl acetate (Langer et al., *Id.*) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

10 Sustained-release PPY and PYY agonists include liposomally PPY and PYY agonists (see generally, Langer, *Science* 249:1527-1533, 1990; Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317-327 and 353-365, 1989). Liposomes containing PPY peptide and peptide analogs are prepared by methods known per se: 15 DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci. U.S.A.* 82:3688-3692, 1985; Hwang et al., *Proc. Natl. Acad. Sci. U.S.A.* 77:4030-4034, 1980; EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Patent Application No. 83-118008; U.S. Patent No. 4,485,045, U.S. Patent No. 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar 20 type in which the lipid content is greater than about 30 mole percent cholesterol, the selected proportion being adjusted for the optimal performance.

Preparations for administration can be suitably formulated to give controlled release of PYY and PYY agonists. For example, the pharmaceutical compositions may be in the form of particles comprising a biodegradable polymer and/or a 25 polysaccharide jellifying and/or bioadhesive polymer, an amphiphilic polymer, an agent modifying the interface properties of the particles and a pharmacologically active substance. These compositions exhibit certain biocompatibility features which allow a controlled release of the active substance. See U.S. Patent No. 5,700,486.

30 In yet an additional embodiment, PPY and PYY agonists are delivered by way of a pump (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201, 1987; Buchwald et al., *Surgery* 88:507, 1980; Saudek et al., *N. Engl. J. Med.*

-31-

321:574, 1989) or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The key factor in selecting an appropriate dose is the result obtained, as measured by decreases in total body weight or ratio of fat to lean mass, or by other criteria for measuring control or 5 prevention of obesity or prevention of obesity-related conditions, as are deemed appropriate by the practitioner. Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533, 1990).

In another aspect of the disclosure, PYY and PYY agonists are delivered by way of an implanted pump, described, for example, in U.S. Patent No. 6,436,091; 10 U.S. Patent No. 5,939,380; U.S. Patent No. 5,993,414.

Implantable drug infusion devices are used to provide patients with a constant and long term dosage or infusion of a drug or any other therapeutic agent. Essentially such device may be categorized as either active or passive.

Active drug or programmable infusion devices feature a pump or a metering 15 system to deliver the drug into the patient's system. An example of such an active drug infusion device currently available is the Medtronic SynchroMed™ programmable pump. Such pumps typically include a drug reservoir, a peristaltic pump to pump out the drug from the reservoir, and a catheter port to transport the pumped out drug from the reservoir via the pump to a patient's anatomy. Such 20 devices also typically include a battery to power the pump as well as an electronic module to control the flow rate of the pump. The Medtronic SynchroMed™ pump further includes an antenna to permit the remote programming of the pump. Passive drug infusion devices, in contrast, do not feature a pump, but rather rely upon a 25 pressurized drug reservoir to deliver the drug. Thus such devices tend to be both smaller as well as cheaper as compared to active devices. An example of such a device includes the Medtronic IsoMed™. This device delivers the drug into the patient through the force provided by a pressurized reservoir applied across a flow control unit.

The implanted pump can be completely implanted under the skin of a patient, 30 thereby negating the need for a percutaneous catheter. These implanted pumps can provide the patient with PYY or a PYY agonist at a constant or a programmed delivery rate, e.g., to give pulsed doses at or around meal time. Constant rate or

programmable rate pumps are based on either phase-change or peristaltic technology. When a constant, unchanging delivery rate is required, a constant-rate pump is well suited for long-term implanted drug delivery. If changes to the infusion rate are expected, a programmable pump may be used in place of the 5 constant rate pump system. Osmotic pumps may be much smaller than other constant rate or programmable pumps, because their infusion rate can be very low. An example of such a pump is described listed in U.S. Patent No. 5,728,396.

For oral administration, the pharmaceutical compositions can take the form of, for example, tablets or capsules prepared by conventional means with 10 pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets can be 15 coated by methods well known in the art. Liquid preparations for oral administration can take the form of, for example, solutions, syrups or suspensions, or they can be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol 20 syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations can also contain buffer salts, flavoring, coloring and sweetening agents as appropriate.

25 For administration by inhalation, the compounds for use according to the present disclosure are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a 30 pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g., gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound

and a suitable powder base such as lactose or starch.

The compounds can also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

5 In addition to the formulations described previously, the compounds can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an
10 acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Pharmaceutical compositions that comprise a PYY, or an agonist thereof, as described herein as an active ingredient will normally be formulated with an appropriate solid or liquid carrier, depending upon the particular mode of
15 administration chosen. The pharmaceutically acceptable carriers and excipients useful in this disclosure are conventional. For instance, parenteral formulations usually comprise injectable fluids that are pharmaceutically and physiologically acceptable fluid vehicles such as water, physiological saline, other balanced salt solutions, aqueous dextrose, glycerol or the like. Excipients that can be included
20 are, for instance, other proteins, such as human serum albumin or plasma preparations. If desired, the pharmaceutical composition to be administered may also contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate. Other medicinal and pharmaceutical
25 agents, for instance other appetite suppressants, or protease inhibitors, also may be included. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in the art.

The dosage form of the pharmaceutical composition will be determined by the mode of administration chosen. For instance, in addition to injectable fluids,
30 inhalation, suppository, and oral formulations can be employed. The pharmaceutical compositions can be produced of conventional mixing, granulating, confectioning, dissolving or lyophilizing processes.

-34-

Oral formulations may be liquid (e.g., syrups, solutions or suspensions), or solid (e.g., powders, pills, tablets, or capsules). For example, pharmaceutical compositions for oral use can be obtained by combining the active ingredient with one or more solid carriers, optionally granulating a resulting mixture, and, if desired, 5 processing the mixture or granules, if appropriate with the addition of additional excipients, to form tablets or dragee cores.

Suitable carriers include fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, also binders, such 10 as starches, for example corn, wheat, rice or potato starch, methylcellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose and/or polyvinylpyrrolidone, and/or, if desired, disintegrators, such as the above-mentioned starches, also carboxymethyl starch, cross-linked polyvinylpyrrolidone, alginic acid or a salt thereof, such as sodium alginate. Additional excipients include flow 15 conditioners and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol, or derivatives thereof.

For parenteral administration compositions include suitable aqueous solutions of an active ingredient in water-soluble form, for example in the form of a 20 water-soluble salt, or aqueous injection suspensions that contain viscosity-altering substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, if desired, stabilizers. The active ingredient, optionally together with excipients, can also be in the form of a lyophilisate and can be made into a solution prior to parenteral administration by the addition of suitable solvents. Solutions 25 such as those that are used, for example, for parenteral administration can also be used as infusion solutions.

For inhalation, PYY or an agonist thereof is administered as an aerosol or a dispersion in a carrier. In one specific, non-limiting example, PYY or an agonist thereof is administered as an aerosol from a conventional valve, such as, but not 30 limited to, a metered dose valve, through an aerosol adapter also known as an actuator. A suitable fluid carrier can be also included in the formulation, such as, but not limited to, air, a hydrocarbon, such as n-butane, propane, isopentane,

amongst others, or a propellant, such as, but not limited to a fluorocarbon. Optionally, a stabilizer is also included, and/or porous particles for deep lung delivery are included (e.g., see U.S. Patent No. 6,447,743).

Compounds with poor solubility in aqueous systems require formulation by 5 using solubilizing agents such as ionic surfactants, cholates, polyethylene glycol (PEG), ethanol, or other agents which may have undesirable effects when used for inhalation. In addition, a treatment requiring successful delivery into alveoli of the lower pulmonary region may preclude from the formulation the use of certain irritants such as chlorofluorocarbons and should involve a minimum number of 10 required doses. Alternatively, to avoid such limitations, liposomes or hydrophobic particles can be used. In one embodiment, an inhalation formulation for a sustained release includes using aerosol droplet particles approximately 1-2.1 μm in size, or of less than 1 μm in size. Small particle aerosol liposomes and liposome-drug combinations for medical use have been previously described (e.g., see EP 15 87309854.5).

According to the invention, a therapeutically effective amount of PYY or an agonist thereof is administered with a therapeutically effective amount of GLP-1 or an agonist thereof.

The PYY or agonist thereof and the GLP-1 or an agonist thereof may be 20 administered simultaneously or substantially simultaneously, or sequentially, in any order. The PYY or agonist thereof and the GLP-1 or an agonist thereof may be administered in a single pharmaceutical composition or in separate compositions, and they may be administered by the same route or by different routes.

If the PYY and the GLP-1 or an agonist thereof are to be administered in a 25 single pharmaceutical composition, that composition may be any of those described above for PYY or an agonist thereof. The composition may enable simultaneous or substantially simultaneous administration of the PYY or agonist thereof and the GLP-1 or an agonist thereof. If desired, the PYY or agonist thereof and the GLP-1 or an agonist thereof may be compartmentalized in the composition, for example, in 30 different layers of a tablet, or in different granules in a capsule. If desired, such compartmentalization may be designed to give different release properties to the two

components to enable delivery of the PYY or agonist component and the GLP-1 or an agonist thereof at different times, for example, sequentially.

Alternatively, the PYY or agonist thereof and the GLP-1 or an agonist thereof may be formulated in separate pharmaceutical compositions, for example, 5 any of the pharmaceutical compositions described above for PYY and agonists thereof. Such separate compositions may be administered simultaneously or substantially simultaneously, or they may be administered sequentially, in any order.

If administered separately, whether sequentially or simultaneously (or substantially simultaneously), the PYY or agonist thereof and the GLP-1 or an 10 agonist thereof may be administered by the same route or by different routes. It is generally more convenient to administer all the active agents in a single composition. However, in some cases it may be necessary or appropriate to administer the active agents by different routes. For example, peptides are generally not stable on oral administration unless modified or formulated in a special way, so 15 must generally be administered via a non-oral route. Some agonists, for example, GLP-1 agonists, are chemical compounds that are stable when administered orally. It may be appropriate to administer OXM non-orally and the other component by a non-oral route.

When PYY or an agonist thereof and GLP-1 or an agonist thereof are used in 20 the manufacture of a medicament for use in a treatment as described herein, the medicament may be a single pharmaceutical composition comprising both components, as described above, or may be a two-component medicament, one component being a pharmaceutical composition comprising PYY or an agonist thereof, the other component being a pharmaceutical composition comprising GLP-1 25 or an agonist thereof, see above. The medicament, whether a one component medicament or a two component medicament as described above, will generally be packaged with instructions relating to its use. Such instructions will refer to the timing, dose and route of administration of the component(s).

If desired, a further agent that influences food intake and/or appetite may 30 also be administered. Such agents include naturally-occurring agents and agonists thereof, for example, amylin and amylin agonists, leptin and leptin agonists, and oxytomodulin.

If desired, one or more other agents, such as, but not limited to, an additional appetite suppressant, may also be administered. Specific, non-limiting example of an additional appetite suppressant include amfepramone (diethylpropion), phentermine, mazindol and phenylpropanolamine, fenfluramine, dexfenfluramine, 5 and fluoxetine.

10 PYY or a PYY agonist, or GLP-1 or a GLP-1 agonist, or both, can be administered simultaneously with the additional agent(s), or the various substances may be administered sequentially, and in any order. Thus, in one embodiment, PYY or an agonist thereof and GLP-1 or an agonist thereof is formulated and administered with an additional agent, for example, oxyntomodulin or an agonist thereof, or an appetite suppressant as a single composition.

15 Additionally, a method of treating obesity is disclosed herein. The method includes administering to an obese subject a therapeutically effective amount of PYY or a PYY agonist, and GLP-1 or an agonist thereof. The PYY agonist can have potency in at least one of food intake or gastric emptying greater than NPY. PYY or a PYY agonist can be administered peripherally, such as in a single or divided dose. Suitable single or divided doses include, but are not limited to, 1 μ g to about 5 mg or about 0.01 μ g/kg to about 500 μ g/kg per dose. The subject can be insulin resistant or glucose intolerant, or both. In addition to being obese, the 20 subject can have diabetes mellitus.

25 A method of reducing food intake is also disclosed herein. The method includes administering to an obese subject a therapeutically effective amount of PYY or a PYY agonist, and GLP-1 or an agonist thereof. The PYY agonist can have potency in at least one of food intake or gastric emptying greater than NPY. PYY or a PYY agonist can be administered peripherally, such as in a single or divided dose. Suitable single or divided doses include, but are not limited to, 1 μ g to about 5 mg or about 0.01 μ g/kg to about 500 μ g/kg per dose. The subject can have Type II diabetes, and/or can be overweight.

30 A method is disclosed herein for improving lipid profile in a subject. The method includes administering to the subject an effective amount of PYY or a PYY agonist, and GLP-1 or an agonist thereof. An improvement in lipid profile includes, but is not limited to, at least one method of reducing cholesterol levels, reducing

triglyceride levels and increasing HDL cholesterol levels. PYY or a PYY agonist can be administered peripherally, such as in a single or divided dose. PYY or a PYY agonist can be administered peripherally, such as in a single or divided dose. Suitable single or divided doses include, but are not limited to, 1 μ g to about 5 mg or 5 about 0.01 μ g/kg to about 500 μ g/kg per dose. The PYY agonist can have potency in at least one of food intake or gastric emptying greater than NPY.

In another embodiment, a method is disclosed herein for alleviating a condition or disorder which can be alleviated by reducing nutrient availability. The method includes administering to a subject a therapeutically effective amount of 10 PYY or a PYY agonist, and GLP-1 or an agonist thereof. Suitable disorders include any of the disorders mentioned above.

The PYY or a PYY agonist and GLP-1 or GLP-1 agonist can be administered peripherally, such as in a single or divided dose. Suitable single or divided doses include, but are not limited to, 1 μ g to about 5 mg or about 0.01 μ g/kg 15 to about 500 μ g/kg of PYY or a PYY agonist per dose. The PYY agonist can have potency in at least one of food intake or gastric emptying greater than NPY. Suitable doses also include those that raise the concentration of PYY and/or the agonist thereof significantly above the basal concentration of PYY, such as, but not limited to, a dose that that mimic postprandial serum concentrations of PYY (or the 20 agonist). Thus, in one embodiment, PYY or an agonist thereof is administered to achieve the level of to effect a reduction in calorie intake, food intake, or appetite equivalent to the reduction in calorie intake, food intake, or appetite, or to increase the energy expenditure, caused by the postprandial level of PYY₃₋₃₆. Specific, non-limiting examples of doses include, but are not limited doses that produce the effect 25 demonstrated when the serum levels of PYY are from about 40 pM/litre to about 60 pM/litre, for example, from about 40 pM/litre to about 50 pM/litre, or from about 40 pM/litre to about 45 pM/litre, or to about 43 pM/litre.

For all methods disclosed herein, the dose of PYY or PYY₃₋₃₆ can be based on the physiological levels observed post-prandially. The normal circulating levels 30 of PYY₃₋₃₆ are about 8 pmol/litre, typically rising to about 40 to 60 pmol/litre after a meal. Agonists of PYY can be used at analogous doses. A single dose may be administered per day, or divided doses can be used (see above).

As indicated above, PYY₃₋₃₆ has a prolonged and sustained action, continuing to act even after it has been cleared from the circulating blood, for example, for up to 12 and even up to 24 hours. Accordingly, PYY or an agonist thereof may be administered twice per day or even just once. If desired, more doses 5 may be administered, for example, three or four per day. For example, it may be desired to administer PYY or an agonist thereof before a meal, for example, about half an hour before a meal. It may be desired to administer PYY or an agonist thereof before each meal. Alternately, it may be desired to administer as few doses as possible, in which case a regime of one or two doses per day may be used.

10 GLP-1 or an agonist may be administered according to the same regime as the PYY or agonist thereof, whether administered simultaneously or substantially simultaneously with the PYY or PYY agonist, or sequentially. Alternatively, the GLP-1 may be administered in a different regime, for example, with more doses of GLP-1 than of PYY, for example, the PYY or PYY agonist may be administered 15 once or twice per day, and the GLP-1 or agonist thereof may be administered three or four times per day. For example, PYY or a PYY agonist may be administered in the morning and evening, for example, before breakfast and before the evening meal, and the GLP-1 agonist thereof may also be administered before the mid-day meal.

20 In one embodiment, when administered peripherally, PYY, including PYY₃₋₃₆ has its effects at physiological levels. No side effects are observed when PYY₃₋₃₆ is used at such levels. Without being bound by theory, PYY₃₋₃₆ does not affect Y2 receptors throughout the brain, which could cause side effects. It should be noted, without being limiting, that a further advantage of PYY₃₋₃₆ is that PYY₃₋₃₆ 25 does not increase blood pressure. The effects of PYY₃₋₃₆ are as long lasting as 24 hours. Recipients claim a decrease in appetite over that period, and a reduction of food intake of about one third has been reported.

- In one specific, non-limiting example, PYY₃₋₃₆ is administered in a dose of about 1 nmol or more, 2 nmol or more, or 5 nmol or more. In this example, 30 the dose of PYY₃₋₃₆ is generally not more than 100 nmol, for example, the dose is 90 nmols or less, 80 nmols or less, 70 nmols or less, 60 nmols or less, 50 nmols or less, 40 nmols or less, 30 nmols or less, 20 nmols or less, 10 nmols. For example, a

-40-

dosage range may comprise any combination of any of the specified lower dose limits with any of the specified upper dose limits. Thus, exemplar non-limiting dose ranges include a dose of PYY₃₋₃₆ may be within the range of from 1 to 100 nmols, from 1 to 90 nmols, from 1 to 80 nmols. Exemplary, non-limiting dose ranges 5 include, from 2 to 100 nmols, from 2 to 90 nmols, for example, from 2 to 80 nmols etc., from 5 nmols to 100 nmols, from 5 nmols to 90 nmols, from 5 nmols to 80 nmols etc. By way of example, a dose of from about 5 to about 50 nmol may be administered such as, but not limited to, from about 2 to about 20 nmol, for example, about 10 nmol. The selected dose may be administered for example, by 10 injection, for example, as a subcutaneous injection. A PYY agonist may be administered at the same dose.

In a further, non-limiting example, PYY₃₋₃₆ may be administered peripherally at a dose of 0.1 nmoles or more per kg body weight of the subject, for example, 0.2 nmoles or more, for example, 0.4 nmoles or more, for example, 0.6 15 nmoles or more, for example, 0.8 nmoles or more, for example, 1.0 nmoles or more, for example, 1.2 nmoles or more, for example, 1.4 nmoles or more, for example, 1.6 nmoles or more, for example, 1.8 nmoles or more, for example, 2.0 nmoles or more, for example, 2.2 nmoles or more, for example, 2.4 nmoles or more, for example, 2.6 nmoles or more, for example, 2.8 nmoles, for example, 3.0 nmoles or more, for 20 example, up to 3.2 nmoles per kg body weight. PYY₃₋₃₆ or a PYY agonist and may be administered peripherally in an amount of up to 3.0 nmoles per kg body weight, for example, up to 2.8 nmoles, for example, up to 2.6 nmoles, for example, up to 2.4 nmoles, for example, up to 2.2 nmoles, for example, up to 2.0 nmoles, for example, up to 1.8 nmoles, for example, up to 1.4 nmoles, for example, up to 1.2 nmoles, for 25 example, up to 1.0 nmoles, for example, up to 0.8 nmoles, for example, up to 0.6 nmoles, for example, up to 0.4 nmoles, for example, up to 0.2 nmoles per kg body weight.

For subcutaneous administration, a dose of PYY for example PYY₃₋₃₆ or a PYY agonist of 10 nmoles or more, for example, 20 nmoles or more, for example, 30 30 nmoles or more, for example, 40 nmoles or more may be used. It is generally preferable to use up to about 70 nmoles, for example, up to about 60 nmol. A dose of about 40 nmol may be used. In one embodiment, a dose of PYY or PYY₃₋

₃₆ at 0.143 n moles (1/7th of a mole) is administered per kilogram, to achieve a dose that is similar to the postprandial level of PYY.

If PYY itself or an agonist thereof is used, the dose is preferably a molar equivalent of a PYY₃₋₃₆ dose, or achieves the same effect as a PYY₃₋₃₆ dose, as described above. The doses can be calculated on the basis of a subject, such as a subject weighing from 70 to 75 kg. The exact dose is readily determined by one of skill in the art based on the potency of the specific compound (such as the PYY polypeptide, or agonist) utilized, and the age, weight, sex and physiological condition of the subject.

10 All the teachings given above in relation to PYY₃₋₃₆ also apply to PYY itself and to PYY agonists. For example, the teachings relating to routes of administration and dosage regimes.

15 The GLP-1 or an agonist thereof should be administered in an therapeutically effective dose, see the section relating to GLP-1 and agonists thereof below. GLP-1 or an agonist thereof may be used in amounts or doses in the ranges described above for PYY or a PYY agonist. The doses of the various agent may be independent of each other or, for example, equimolar doses may be used, for example, equimolar doses of GLP-1 or an agonist thereof and PYY or an agonist thereof.

20 As disclosed herein, a naturally occurring peptide, PYY or PYY₃₋₃₆ can be used to achieve a physiological effect. This results in minimal side effects and enables long term use, if necessary. The dose of PYY or PYY₃₋₃₆ can be based on the physiological levels observed post-prandially. The normal circulating levels of PYY₃₋₃₆ are about 8 pmol/litre, typically rising to about 40 to 60 pmol/litre after a meal. PYY (e.g., PYY₃₋₃₆) and agonists can be used at analogous doses.

25 The considerations set out above in relation to the administration regime also relate to administration using the subcutaneous route.

30 The various uses of PYY, or an agonist thereof, and GLP-1 or an agonist thereof, as set out above may be in a method of treatment of a mammalian subject in need of such treatment, or may be in the manufacture of a medicament for such treatment. PYY (e.g., PYY₃₋₃₆) or an agonist thereof, and GLP-1 or an agonist thereof, should be administered in an amount effective to achieve the stated object. Some of the treatments described above are medical treatments, for example, the

treatment of obesity. Others, however, do not relate to medical treatment, and are part of the maintenance of a healthy lifestyle, or are for cosmetic purposes.

The use of a combination of any of OXM and GLP-1 or an agonist thereof and PYY or an agonist thereof may serve to increase the effectiveness of any of the 5 agents compared with its use alone, for example, as described above. Alternatively or in addition, use of the two or three agents in combination may reduce any tendency for "escape" when using an agent alone. The term "escape" is used to denote a reduction in effect of an agent with time. For example, if any one of the agents above has been used alone, its effect may reduce with time. Use of one or 10 both of the other agents in addition may reduce or prevent the tendency for that reduction in effectiveness. For example, PYY has a sustained effect and may be used for prolonged periods. If the effect of PYY should appear to reduce, or to reduce or prevent any such reduction in effect, OXM may be administered in addition to the PYY. GLP-1 may also be used for the same purpose, with OXM or 15 with OXM and PYY.

In another aspect the present invention provides a pharmaceutical composition that comprises PYY or a PYY agonist as the active ingredient and a pharmaceutically suitable carrier in a form suitable for subcutaneous administration, for example, as an injectable solution. Such a composition may comprise an amount 20 of PYY for example PYY₃₋₃₆ or a PYY agonist of 10 nmoles or more, for example, 20 nmoles or more, for example, 30 nmoles or more, for example, 40 nmoles or more may be used. It is generally preferable to use up to about 70 nmoles, for example, up to about 60 nmnoles. An amount of about 40nmnoles may present.. Suitable formulations for subcutaneous administration are well known, and are 25 described above. One or more further active ingredient(s) may also be present in the composition, in particular GLP-1 or an agonist thereof, which may be used in amounts and doses as described above. Further active ingredients may be present, for example, as described above, for example, oxyntomodulin or an agonist thereof, or an appetite suppressant as described above.

30 A further aspect of the invention relates to methods as disclosed above, including a method for decreasing calorie intake in a subject, a method for decreasing appetite in a subject, a method for decreasing food intake in a subject, a

-43-

method for weight control or treatment in a subject, and a method for reduction or prevention of obesity, in particular any one or more of the following: preventing and reducing weight gain; inducing and promoting weight loss; and reducing obesity as measured by the Body Mass Index. The methods include control of any one or more 5 of appetite, satiety and hunger, in particular any one or more of the following: reducing, suppressing and inhibiting appetite; inducing, increasing, enhancing and promoting satiety and sensations of satiety; and reducing, inhibiting and suppressing hunger and sensations of hunger. The methods further include maintaining any one or more of a desired body weight, a desired Body Mass Index, a desired appearance 10 and good health. In the methods of this aspect of the invention, PYY, for example, PYY₃₋₃₆ or a PYY agonist is administered subcutaneously to the subject.

Preferably the PYY or PYY agonist is administered at dose of 10 nmoles or more, for example, 20 nmoles or more, for example, 30 nmoles or more, for example, 40 nmoles or more may be used. It is generally preferable to use up to 15 about 70 nmoles, for example, up to about 60 nmnoles. An dose of about 40nmoles be used. As indicated above, PYY₃₋₃₆ has a prolonged and sustained action, continuing to act even after it has been cleared from the circulating blood, for example, for up to 12 and even up to 24 hours. Accordingly, PYY or an agonist thereof may be administered twice per day or even just once. If desired, more doses 20 may be administered, for example, three or four per day. For example, it may be desired to administer PYY or an agonist thereof before a meal, for example, about half an hour before a meal. It may be desired to administer PYY or an agonist thereof before each meal. Alternately, it may be desired to administer as few doses as possible, in which case a regime of one or two doses per day may be used.

25 In this aspect of the invention, PYY or an agonist thereof may be administered alone or PYY or an agonist thereof and GLP-1 or an agonist thereof may be administered. If GLP-1 or an agonist thereof is used in addition to the PYY or PYY agonist, the GLP-1 or agonist thereof may be administered simultaneously or substantially simultaneously with the PYY or agonist thereof, or sequentially, in 30 either order. The PYY or agonist thereof and the GLP-1 or agonist thereof may be administered in a single pharmaceutical composition or in separate compositions, and they may be administered by the same route or by different routes. Further

active ingredients may also be administered, for example, as described above for subcutaneous PYY pharmaceutical compositions.

The present invention also includes PYY or an agonist thereof for use in the manufacture of a medicament for subcutaneous administration for any of the

5 methods of treatment described above.

PYY Agonists

A PYY agonist, of use in the methods of the present disclosure, is a molecule
10 that binds to a receptor that specifically binds PYY, and elicits an effect of PYY. Assays for binding to PYY receptors, and eliciting a response in a cell with a PYY receptor, are known in the art. A specific assay for detecting a PYY agonist is also disclosed herein. Thus, in one embodiment, a PYY agonist binds to a NPY neuron in the arcuate nucleus, which results in an electrophysiological effect on an NPY
15 neuron. As disclosed herein, NPY neurons synapse with POMC neurons. Thus, the electrophysiological effect on the NPY neuron can result in a further electrophysiological effect on a POMC neuron. In one specific, non-limiting example, an administration of PYY agonist results in hyperpolarization of the membrane potential of a POMC neuron. In another specific, non-limiting example, 20 administration of a PYY agonist results in an increase in IPSCs in a POMC neuron.

In another embodiment, PYY agonists do not include NPY. Suitable PYY agonists include molecules that bind NPY neurons, but do not cross the blood/brain barrier. The arcuate nucleus neurons upon which PYY exerts its effects are not protected by the blood/brain barrier, and thus are readily accessible to peripherally
25 available molecules. In addition, other brain sites that express the Y2 receptor are protected by the blood/brain barrier. Without being bound by theory, agents able to bind to the arcuate Y2R, but that do not cross the blood/brain barrier following peripheral administration, are likely to be of use.

In one embodiment, a PYY agonist is a compound that affects food intake,
30 caloric intake, or appetite, and/or which binds specifically in a Y receptor assay or competes for binding with PYY, such as in a competitive binding assay with labeled

PYY. PYY agonists include, but are not limited to, compounds that bind to the Y2 receptor.

PYY and agonists useful in the methods disclosed herein include, but are not limited to, polypeptides comprising, or alternatively consisting of, the amino acid sequence for PYY and agonists thereof, e.g., mutants, fragments and/or variants thereof. Variants include deletions, insertions, inversions, repeats and substitutions (e.g., conservative substitutions and non-conservative substitutions; see, e.g., Tables 1 and 2, *infra*). More than one amino acid (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, etc.) can be deleted or inserted or substituted with another amino acid. Typically conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile; interchange of Ser and Thr containing hydroxy residues, interchange of the acidic residues Asp and Glu, interchange between the amide residues Asn and Gln, interchange of the basic residues Lys and Arg, interchange of the aromatic residues Phe and Tyr, and interchange of the small-sized amino acids Ala, Ser, Thr, Met and Gly. Guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., *Science* 247:1306-1310, 1990.

As another example, polypeptide fragments may contain a continuous series of deleted residues from the amino (N)- or the carboxyl (C)- terminus, or both (see, e.g., Tables 1 and 2, *infra*). Any number of amino acids, ranging from 1 to 24, can be deleted from the N-terminus, the C-terminus or both.

Furthermore, the agonist polypeptides may also include, but are not limited to, polypeptides comprising, or alternatively consisting of, internal deletions of the amino acid sequences for PYY and/or agonist thereof (see, e.g., Table 2, *infra*). Such deletions may comprise one or more amino acid residue deletions (e.g., one, two, three, four, five, six, seven, eight, nine, ten, etc.) and may begin at any amino acid position (e.g., two, three, four, five, six, seven, eight, nine, ten, etc.). In addition, the polypeptides of this disclosure may contain one or more such internal deletions. Such deletions are contemplated in PYY, NPY and PP. Also contemplated are agonist peptides that are PYY, NPY and/or PP chimeras having high affinity and/or selectivity for the Y2 receptor. These chimeras may comprise amino acid substitutions of one or more amino acids (e.g., 1, 2, 3, 4,

5, 6, 7, 8, 9, 10, etc.) from PPY, NPY and/or PP, variants, mutants and/or deletions thereof, with one or more amino acids (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, etc.) from a second PPY, NPY, or PP, variants, mutations and/or deletions thereof. These substitutions may begin at any amino acid position (e.g., two, three, four, five, six, 5 seven, eight, nine, ten, etc.).

Preferably, the peptide is selective for the Y2 receptor. That is, it binds with higher affinity to Y2 compared to other receptors, such as Y1, Y2, Y3, Y4, Y5 and Y6. In another embodiment, the peptide is selective for the Y2 and Y5 receptors over the Y1, Y3, Y4 and Y6 receptors.

10 Other polypeptide fragments are fragments comprising structural or functional domain of the polypeptides of this disclosure. Such fragments include amino acid residues that comprise a polyproline-type II helix (residues 1-8), beta-turn (residues 9-14), amphipathic alpha-helix (residues 15-32) and/or a C-terminal turn structure (residues 33-36). See, Kirby et al., *J Med Chem* 36:385-393, 1993.

15 In addition, this disclosure includes the use of a polypeptide or agonist comprising, or alternatively consisting of, the amino acid sequence for PPY, NPY and PP species variants (see Table 1, *infra*) and/or mutants, and fragments thereof.

Also contemplated are fusion proteins, whereby a PYY or PYY agonist will be fused to another protein or polypeptide (the fusion partner) using recombinant 20 methods known in the art. Alternatively, such a fusion protein may be synthetically synthesized by any known method. Any known peptide or protein can be used as the fusion partner (e.g., serum albumin, carbonic anhydrase, glutathione-S-transferase or thioredoxin, etc.). Preferred fusion partners will not have an adverse biological activity *in vivo*. Such fusion proteins may be designed linking the 25 carboxy-terminus of the fusion partner to the amino-terminus of the PYY or agonist peptide, or vice versa. Optionally, a cleavable linker region may be used linking the PYY or PYY agonist to the fusion partner, and may be cleaved *in vivo* thereby resulting in the release of an active form of PYY or a PYY agonist. Examples of 30 such cleavage regions include, but are not limited to, the linker regions D-D-D-D-Y (SEQ ID NO: 330), G-P-R (SEQ ID NO: 331), A-G-G (SEQ ID NO: 332) and H-P-F-H-L (SEQ ID NO 333), which can be cleaved by enterokinase, thrombin, ubiquitin cleaving enzyme and renin, respectfully. See, e.g., U.S. Patent No. 6,410,707.

Also contemplated as useful PYY agonists are Y2 specific NPY peptide agonists as described in U.S. Patent No. 5,026,685; U.S. Patent No. 5,574,010; U.S. Patent No. 5,604,203; U.S. Patent No. 5,696,093; U.S. Patent No. 6,046,167. See below:

5

Preferred PYY agonists are described herein as follows.

TABLE 1 - PYY: Variation Among Species

10	PEPTIDE YY	AA SEQUENCE
	Human	YPIKPEAPGEDASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 1)
	Rat	YPAKPEAPGEDASPEELSRYYASLRHYLNLVTRQRY (SEQ ID NO: 5)
	Pig	YPAKPEAPGEDASPEELSRYYASLRHYLNLVTRQRY (SEQ ID NO: 6)
	Guinea pig	YPSKPEAPGSDASPEELARYYASLRHYLNLVTRQRY (SEQ ID NO: 7)
15	Frog	YPPKPENPGEDASPEEMTKYLTLRHYINLVTRQRY (SEQ ID NO: 8)
	Raja	YPPKPENPGDDAAPEELAKYYASLRHYINLITRQRY (SEQ ID NO: 9)
	Dogfish	YPPKPENPGEDAPPEELAKYYASLRHYINLITRQRY (SEQ ID NO: 10)
	Lampetra	FPPKPDPNGDNASPEQMARYKAAVRHYINLITRQRY (SEQ ID NO: 11)
	Petromyzon	MPPKPDPNPSPDASPEELSKYMLAVRNYINLITRQRY (SEQ ID NO: 12)
20	NEUROPEPTIDE Y	AA SEQUENCE
	Human	YPSKPDNPAGEDAPAEDMARYSALRHYINLITRQRY (SEQ ID NO: 2)
	Rat	YPSKPDNPAGEDAPAEDMARYSALRHYINLITRQRY (SEQ ID NO: 13)
	Rabbit	YPSKPDNPAGEDAPAEDMARYSALRHYINLITRQRY (SEQ ID NO: 14)
25	Dog	YPSKPDNPAGEDAPAEDMARYSALRHYINLITRQRY (SEQ ID NO: 15)
	Pig	YPSKPDNPAGEDAPAEDLARYYSALRHYINLITRQRY (SEQ ID NO: 16)
	Cow	YPSKPDNPAGEDAPAEDLARYYSALRHYINLITRQRY (SEQ ID NO: 17)
	Sheep	YPSKPDNPAGEDAPAEDLARYYSALRHYINLITRQRY (SEQ ID NO: 18)
	Guinea pig	YPSKPDNPAGEDAPAEDMARYSALRHYINLITRQRY (SEQ ID NO: 19)
30	Avian	YPSKPDSPGEDAPAEDMARYSALRHYINLITRQRY (SEQ ID NO: 20)
	Rana	YPSKPDNPAGEDAPAEDMAKYYASLRHYINLITRQRY (SEQ ID NO: 21)
	Goldfish	YPTKPDNPGEGAPAEELAKYYASLRHYINLITRQRY (SEQ ID NO: 22)
	Dogfish	YPSKPDNPGEGAPAEELAKYYASLRHYINLITRQRY (SEQ ID NO: 23)
	Lampetra	PPNKPDPSPGEDAPAEDLARYLSAVRHYINLITRQRY (SEQ ID NO: 24)
35	PANCREATIC POLYPEPTIDE	AA SEQUENCE
	Human	ASLEPEYPGDNATPEQMAQYAAELRRYINMLTRPRY (SEQ ID NO: 3)

	Sheep	APLEPVYPGDNATPEQMAQYAAELRRYINMLTRPRY (SEQ ID NO: 25)
	Pig	APLEPVYPGDDATPEQMAQYAAELRRYINMLTRPRY (SEQ ID NO: 26)
	Dog	APLEPVYPGDDATPEQMAQYAAELRRYINMLTRPRY (SEQ ID NO: 27)
	Cat	APLEPVYPGDNATPEQMAQYAAELRRYINMLTRPRY (SEQ ID NO: 28)
5	Cow	APLEPEVYPGDNATPEQMAQYAAELRRYINMLTRPRY (SEQ ID NO: 29)
	Rat	APLEPMYPGDYATHEQRAQYETQLRRYINTLTRPRY (SEQ ID NO: 30)
	Mouse	APLEPMYPGDYATPEQMAQYETQLRRYINTLTRPRY (SEQ ID NO: 31)
	Guinea pig	APLEPVYPGDNATPEQQMAQYAAEMRRYINMLTRPRY (SEQ ID NO: 32)
	Chicken	GPSQPTYPGDDAPVEDLIRFYNDLQQYLNVVTRHRY (SEQ ID NO: 33)
10	Alligator	TPLQPKYPGDPGAPVEDLIQFYNDLQQYLNVVTRPRF (SEQ ID NO: 34)
	Bullfrog	APSEPHHPGDQATPDQLAQYYSDLYQYITFIRPRF (SEQ ID NO: 35)

Ref: Beck-Sickinger, A.G., Jung, G., *Biopolymers* 37:123-142, 1995.

15 **TABLE 2 – PEPTIDE AGONIST OF PYY**

	PEPTIDE	SEQUENCE
	PPY(3-36)(human)	
		IKPEAPGEDASPEELNRYYASLRHYLNVLVTRQRY (SEQ ID NO: 334)
20	Ref: Eberlein et al., <i>Peptides</i> 10:797-803, 1989; Grandt et al., <i>Peptides</i> 15(5):815-20, 1994.	

Variations of PPY(3-36)

N-Terminal Deletions of PYY, including but not limited to: PYY(26-36), PYY(25-36), PYY(24-36), PYY(23-36), PYY(22-36), PYY(21-36), PYY(20-36), PYY(19-36), PYY(18-36), PYY(17-36), PYY(16-36), PYY(15-36), PYY(14-36), PYY(13-36), PYY(12-36), PYY(11-36), PYY(10-36), PYY(9-36), PYY(8-36), PYY(7-36), PYY(6-36), PYY(5-36), PYY(4-36), PYY(3-36).

Ref: See, e.g., Balasubramaniam et al., *Pept Res* 1(1):32-5, Sep-Oct 1998; Liu et al., *J Gastrointest Surg* 5(2):147-52, Mar-Apr 2001.

	PEPTIDE	SEQUENCE
	NPY (human)	YPSKPDNPGEDAPAEDMARYYSALRHYINLITRQRY (SEQ ID NO: 2)

-49-

Ref: Tatemoto et al., *Proc Natl Acad Sci U.S.A.* 79:5485-9, 1982.

Variations of NPY

N-Terminal Deletions of NPY, including but not limited to: NPY(26-36), NPY(25-36), NPY(24-36), NPY(23-36), NPY(22-36), NPY(21-36), NPY(20-36), NPY(19-36), NPY(18-36), NPY(17-36), NPY(16-36), NPY(15-36), NPY(14-36), NPY(13-36), NPY(12-36), NPY(11-36), NPY(10-36), NPY(9-36), NPY(8-36), NPY(7-36), NPY(6-36), NPY(5-36), NPY(4-36), NPY(3-36).

5 Ref: See e.g., Gehlert et al., *Proc Soc Exp Biol Med* 218:7-22, 1998; Sheikh et al.,
10 *Am J Physiol* 261:G701-15, Nov. 1991.

Internal Deletions, including but not limited to: (1-4)-Aca-(14-36)pNPY, (1-4)-Aca-(15-36)pNPY, (1-4)-Aca-(16-36)pNPY, (1-4)-Aca-(17-36)pNPY, (1-4)-Aca-(18-36)pNPY, (1-4)-(31-36)pNPY11, (1-4)-Aca-(31-36)pNPY, (4-1)-(31-36)pNPY, (4-1)-Aca-(31-36)pNPY, (4-1)_D-(31-36)pNPY, (4-1)_D-Aca-(31-36)pNPY.

15 Ref: Fournier et al., *Mol Pharmacol* 45(1):93-101, Jan 1994.

Additional Internal Deletion Mutants, including but not limited to: des-AA¹⁰⁻¹⁷-NPY, des-AA¹⁰⁻¹⁷, Ac-[D-Lys⁹(ε-Ac-Ala)]NPY, des-AA¹⁰⁻¹⁷, Ac[D-Lys⁹(ε-Ac-Ala)]NPY, des-AA¹⁰⁻¹⁷[Ala^{7,21}]NPY, des-AA¹⁰⁻¹⁷[Cys^{7,21}]NPY, des-AA¹⁰⁻¹⁷[Glu⁷,Lys²¹]NPY, des-AA¹¹⁻¹⁷[D-Lys¹⁰(ε-Ac), Cys^{7,21}]NPY, des-AA¹⁰⁻¹⁷[D-Cys⁷, D-Lys(ε-Ac), Cys²¹]NPY, des-AA¹⁰⁻¹⁷[D-Cys⁷, Lys⁹(ε-Ac), Cys²¹]NPY, des-AA¹⁰⁻¹⁷[Cys^{7,21}, Pro³⁴]NPY, des-AA¹⁰⁻¹⁷[Asp⁷, Dpr²¹, Pro³⁴]NPY, des-AA¹⁰⁻¹⁷[Glu⁷, Lys²¹, Pro³⁴]NPY, des-AA¹⁰⁻¹⁷[Cys^{7,21}, Leu³¹, Pro³⁴]NPY, des-AA¹⁰⁻²⁰[Cys^{7,21}, Pro³⁴]NPY, 25 des-AA¹⁰⁻¹⁷[Cys^{2,27}]NPY, des-AA¹⁰⁻¹⁷[Cys², D-Cys²⁷]NPY.

Ref: Kirby et al., *J Med Chem* 38:4579-86, 1995.

Cyclic agonist of NPY, including but not limited to: [Lys 25-Glu 29]NPY(Ac-25-36), [Glu 25-Lys 29]NPY(Ac-25-36), [Lys 26-Glu31]NPY(Ac-25-36), [Glu 27-Lys 31]NPY(Ac-25-36), [Lys28-Glu 32]NPY(Ac-25-36), [Lys27-Glu34]NPY(Ac-25-36).

30 Ref: Rist et al., *Eur J Biochem* 247:1019-1028, 1997.

-50-

D-amino acid substitutions: [D-Tyr¹]NPY, [D-Pro²]NPY, [D-Ser³]NPY, [D-Lys⁴]NPY, [D-Pro⁵]NPY, [D-Asp⁶]NPY, [D-Asn⁷]NPY, [D-Pro⁸]NPY, [D-Ala⁹]NPY, [D-Glu¹⁰]NPY, [D-Asp¹¹]NPY, [D-Ala¹²]NPY, [D-Pro¹³]NPY, [D-Ala¹⁴]NPY, [D-Glu¹⁵]NPY, [D-Asp¹⁶]NPY, [D-Leu¹⁷]NPY, [D-Ala¹⁸]NPY, [D-Arg¹⁹]NPY, [D-Tyr²⁰]NPY, [D-Tyr²¹]NPY, [D-Ser²²]NPY, [D-Ala²³]NPY, [D-Leu²⁴]NPY, [D-Arg²⁵]NPY, [D-His²⁶]NPY, [D-Tyr²⁷]NPY, [D-Ile²⁸]NPY, [D-Asn²⁹]NPY, [D-Leu³⁰]NPY, [D-Ile³¹]NPY, [D-Thr³²]NPY, [D-Arg³³]NPY, [D-Gln³⁴]NPY, [D-Arg³⁵]NPY, [D-Tyr³⁶]NPY, [D-Tyr¹, D-Pro²]NPY, [D-Ser³, D-Lys⁴]NPY, [D-Pro⁵, D-Asp⁶]NPY, [D-Asn⁷, D-Pro⁸]NPY, [D-Glu¹⁰, D-Asp¹¹]NPY, [D-Asp¹¹, D-Ala¹²]NPY, [D-Pro¹³, D-Ala¹⁴]NPY, [D-Glu¹⁵, D-Asp¹⁶]NPY, [D-Met¹⁷, D-Ala¹⁸]NPY, [D-Arg¹⁹, D-Tyr²⁰]NPY, [D-Tyr²¹, D-Ser²²]NPY, [D-Ala²³, D-Leu²⁴]NPY, [D-Arg²⁵, D-His²⁶]NPY, [D-Tyr²⁷, D-Ile²⁸]NPY, [D-Asn²⁹, D-Leu³⁰]NPY, [D-Ile³¹, D-Thr³²]NPY, [D-Arg³³, D-Gln³⁴]NPY, [D-Arg³⁵, D-Tyr³⁶]NPY.

Ref: Kirby et al., *J Med Chem* 36:3802-08, 1993; Grundemar et al., *Regulatory Peptides* 62:131-136, 1996.

Other NPY Agonist and Analogs

20 PEPTIDE SEQUENCE
NPY(3-36)
SKPDNPAGEDAPADEMARYYSALRHYNLITRQRY (SEQ ID NO: 335)

Ref: Grandt et al., *Regulatory Peptides* 67(1):33-7, 1996.

25 PEPTIDE SEQUENCE
N-Acetyl NPY(24-36)
LRHYINLITRQRY (SEQ ID NO: 213)

Ref: Potter et al., *Eur J Pharmacol* 267(3):253-262, May 17, 1994.

30 PEPTIDE SEQUENCE
N-Acetyl [Leu²⁸, Leu³¹] NPY(24-36)
LRHYLNLLTRQRY (SEQ ID NO: 214)

Ref: Potter et al., *Eur J Pharmacol* 267(3):253-262, May 17, 1994.

-51-

PEPTIDE SEQUENCE
 [Leu²⁸, Leu³¹] NPY(24-36)
 LRHYLNLLTRQRY (SEQ ID NO: 215)

5 Ref: Potter et al., *Eur J Pharmacol* 267(3):253-262, May 17, 1994.

PEPTIDE SEQUENCE
 [Leu¹⁷, Gln¹⁹, Ala²¹, Ala²², Glu²³, Leu²⁸, Leu³¹] NPY(13-36)
 PAEDLAQYAAELRHYNLLTRQRY (SEQ ID NO: 216)

10 Ref: Potter et al., *Eur J Pharmacol* 267(3):253-262, May 17, 1994.

PEPTIDE SEQUENCE
 Cyclo S-S [Cys²⁰,Cys²⁴]pNPY
 SKPDNPGEDAPAEDMARCYSACRHYINLITRQRY (SEQ ID NO: 315)

15 Ref: Soll et al., *Eur J Biochem* 268(10):2828-37, May 2001.

PEPTIDE SEQUENCE
 Cyclo-(28/32)-Ac-[Lys²⁸-Glu³²]- (25-36)-pNPY
 RHYLNLLIGRQRY (SEQ ID NO: 316)

20 Ref: Cabrele et al., *J Pept Sci* 6(3):97-122, Mar 2000.

PEPTIDE SEQUENCE
 Cyclo-(27/31)-Ac-[Glu²⁷-Lys³¹]- (25-36)-pNPY
 RHGLNLLGRQRY (SEQ ID NO: 317)

25 Ref: Cabrele et al., *J Pept Sci* 6(3):97-122, Mar 2000.

PEPTIDE SEQUENCE
 [Tyr³², Leu³⁴]NPY(27-36)
 YINLIYRLRY (SEQ ID NO: 318)

30 Ref: Leban et al., *J Med Chem* 38:1150-57, 1995.

PEPTIDE SEQUENCE
 [Tyr³², Leu³⁴]NPY(26-36)
 HYINLIYRLRY (SEQ ID NO: 319)

35 Ref: Leban et al., *J Med Chem* 38:1150-57, 1995.

-52-

5 PEPTIDE SEQUENCE

[Tyr³², Leu³⁴]NPY(25-36)

RHYINLIYRLRY (SED ID NO: 320)

Ref: Leban et al., *J Med Chem* 38:1150-57, 1995.

10

[Leu³¹]NPY(27-36)

YINLLYRQRY (SEQ ID NO: 321)

Ref: Leban et al., *J Med Chem* 38:1150-57, 1995.

15 PEPTIDE SEQUENCE

[Tyr³², Leu³⁴] (1-4)-Ahr-(27-36)NPY

YPSL-Aha-YINLIYRLRY (SED ID NO: 322)

Ref: Leban et al., *J Med Chem* 38:1150-57, 1995.

20 PEPTIDE SEQUENCE

[Tyr³², Leu³⁴]NPY(28-36)

INLIYRLRY (SEQ ID NO: 323)

Ref: Leban et al., *J Med Chem* 38:1150-57, 1995.

25 PEPTIDE SEQUENCE

PP (human)

ASLEPEYPGDNATPEQMAQYAAELRRYINMLTRPRY (SEQ ID NO: 3)

Ref: Kimmel et al., *Endocrinology* 83:1323-30, 1968.

30 Variations of PP

N-Terminal Deletions including but not limited to: PP(26-36), PP(25-36), PP(24-36), PP(23-36), PP(22-36), PP(21-36), PP(20-36), PP(19-36), PP(18-36), PP(17-36), PP(16-36), PP(15-36), PP(14-36), PP(13-36), PP(12-36), PP(11-36), PP(10-36), PP(9-36), PP(8-36), PP(7-36), PP(6-36), PP(5-36), PP(4-36), PP(3-36).

35

*TABLE 3 - EXAMPLES OF CONSERVATIVE AMINO ACID
SUBSTITUTIONS OF PYY*

Single point mutations of PYY(25-36)

-53-

	PEPTIDE	SEQUENCE
	[Lys ²⁵]PPY(25-36)	KHYLNLVTRQRY (SEQ ID NO: 36)
	[Thr ²⁷]PPY(25-36)	RHTLNLVTRQRY (SEQ ID NO: 37)
	[Phe ²⁷]PPY(25-36)	RHFLNLVTRQRY (SEQ ID NO: 38)
5	[Ile ²⁸]PYY (25-36)	RHYINLVTRQRY (SEQ ID NO: 39)
	[Val ²⁸]PYY (25-36)	RHYVNLVTRQRY (SEQ ID NO: 40)
	[Gln ²⁹]PYY (25-36)	RHYLQLVTRQRY (SEQ ID NO: 41)
	[Ile ³⁰]PYY (25-36)	RHYLNIVTRQRY (SEQ ID NO: 42)
	[Val ³⁰]PYY (25-36)	RHYLNVVTRQRY (SEQ ID NO: 43)
10	[Ile ³¹]PYY (25-36)	RHYLNLLTRQRY (SEQ ID NO: 44)
	[Leu ³¹]PYY (25-36)	RHYLNLLTRQRY (SEQ ID NO: 45)
	[Ser ³²]PYY (25-36)	RHYLNLSRQRY (SEQ ID NO: 46)
	[Lys ³³]PYY (25-36)	RHYLNLVTKQRY (SEQ ID NO: 47)
	[Asn ³⁴]PYY (25-36)	RHYLNLVTRNRY (SEQ ID NO: 48)
15	[Lys ³⁵]PYY (25-36)	RHYLNLVTRQKY (SEQ ID NO: 49)
	[Thr ³⁶]PYY (25-36)	RHYLNLVTRQRT (SEQ ID NO: 50)
	[Phe ³⁶]PYY (25-36)	RHYLNLVTRQRF (SEQ ID NO: 51)

Double point mutations

	PEPTIDE	SEQUENCE
20	[Lys ²⁵ , Thr ²⁷]PPY(25-36)	KHTLNLVTRQRY (SEQ ID NO: 52)
	[Lys ²⁵ , Phe ²⁷]PPY(25-36)	KHFLNLVTRQRY (SEQ ID NO: 53)
	[Lys ²⁵ , Ile ²⁸]PPY(25-36)	KHYINLVTRQRY (SEQ ID NO: 54)
	[Lys ²⁵ , Val ²⁸]PPY(25-36)	KHYVNLVTRQRY (SEQ ID NO: 55)
25	[Lys ²⁵ , Gln ²⁹]PPY(25-36)	KHYLQLVTRQRY (SEQ ID NO: 56)
	[Lys ²⁵ , Ile ³⁰]PPY(25-36)	KHYLNIVTRQRY (SEQ ID NO: 57)
	[Lys ²⁵ , Val ³⁰]PPY(25-36)	KHYLNVVTRQRY (SEQ ID NO: 58)
	[Lys ²⁵ , Ile ³¹]PPY(25-36)	KHYLNLLTRQRY (SEQ ID NO: 59)
	[Lys ²⁵ , Leu ³¹]PPY(25-36)	KHYLNLLTRQRY (SEQ ID NO: 60)
30	[Lys ²⁵ , Ser ³²]PPY(25-36)	KHYLNLSRQRY (SEQ ID NO: 61)
	[Lys ²⁵ , Lys ³³]PPY(25-36)	KHYLNLVTKQRY (SEQ ID NO: 62)
	[Lys ²⁵ , Asn ³⁴]PPY(25-36)	KHYLNLVTRNRY (SEQ ID NO: 63)
	[Lys ²⁵ , Lys ³⁵]PPY(25-36)	KHYLNLVTRQKY (SEQ ID NO: 64)
	[Lys ²⁵ , Thr ³⁶]PPY(25-36)	KHYLNLVTRQRT (SEQ ID NO: 65)
35	[Lys ²⁵ , Phe ³⁶]PPY(25-36)	KHYLNLVTRQRF (SEQ ID NO: 66)
	[Thr ²⁷ , Ile ²⁸]PPY(25-36)	RHTINLVTRQRY (SEQ ID NO: 67)
	[Thr ²⁷ , Val ²⁸]PPY(25-36)	RHTVNLVTRQRY (SEQ ID NO: 68)
	[Thr ²⁷ , Gln ²⁹]PPY(25-36)	RHTLQLVTRQRY (SEQ ID NO: 69)

	[Thr ²⁷ , Ile ³⁰]PPY(25-36)	RHTLNIVTRQRY (SEQ ID NO: 70)
	[Thr ²⁷ , Val ³⁰]PPY(25-36)	RHTLNVVTRQRY (SEQ ID NO: 71)
	[Thr ²⁷ , Ile ³¹]PPY(25-36)	RHTLNLITRQRY (SEQ ID NO: 72)
	[Thr ²⁷ , Leu ³¹]PPY(25-36)	RHTLNLLTRQRY (SEQ ID NO: 73)
5	[Thr ²⁷ , Ser ³²]PPY(25-36)	RHTLNLVSRQRY (SEQ ID NO: 74)
	[Thr ²⁷ , Lys ³³]PPY(25-36)	RHTLNLVTKQRY (SEQ ID NO: 75)
	[Thr ²⁷ , Asn ³⁴]PPY(25-36)	RHTLNLVTRNRY (SEQ ID NO: 76)
	[Thr ²⁷ , Lys ³⁵]PPY(25-36)	RHTLNLVTRQKY (SEQ ID NO: 77)
	[Thr ²⁷ , Thr ³⁶]PPY(25-36)	RHTLNLVTRQRT (SEQ ID NO: 78)
10	[Thr ²⁷ , Phe ³⁶]PPY(25-36)	RHTLNLVTRQRF (SEQ ID NO: 79)
	[Phe ²⁷ , Ile ²⁸]PPY(25-36)	RHFINLVTRQRY (SEQ ID NO: 80)
	[Phe ²⁷ , Val ²⁸]PPY(25-36)	RHFVNLVTRQRY (SEQ ID NO: 81)
	[Phe ²⁷ , Gln ²⁹]PPY(25-36)	RHFLQLVTRQRY (SEQ ID NO: 82)
	[Phe ²⁷ , Ile ³⁰]PPY(25-36)	RHFLNIVTRQRY (SEQ ID NO: 83)
15	[Phe ²⁷ , Val ³⁰]PPY(25-36)	RHFLNVVTRQRY (SEQ ID NO: 84)
	[Phe ²⁷ , Ile ³¹]PPY(25-36)	RHFLNLITRQRY (SEQ ID NO: 85)
	[Phe ²⁷ , Leu ³¹]PPY(25-36)	RHFLNLLTRQRY (SEQ ID NO: 86)
	[Phe ²⁷ , Ser ³²]PPY(25-36)	RHFLNLVSRQRY (SEQ ID NO: 87)
	[Phe ²⁷ , Lys ³³]PPY(25-36)	RHFLNLVTKQRY (SEQ ID NO: 88)
20	[Phe ²⁷ , Asn ³⁴]PPY(25-36)	RHFLNLVTRNRY (SEQ ID NO: 89)
	[Phe ²⁷ , Lys ³⁵]PPY(25-36)	RHFLNLVTRQKY (SEQ ID NO: 90)
	[Phe ²⁷ , Thr ³⁶]PPY(25-36)	RHFLNLVTRQRT (SEQ ID NO: 91)
	[Phe ²⁷ , Phe ³⁶]PPY(25-36)	RHFLNLVTRQRF (SEQ ID NO: 92)
	[Gln ²⁹ , Ile ³⁰]PYY (25-36)	RHYLQIVTRQRY (SEQ ID NO: 93)
25	[Gln ²⁹ , Val ³⁰]PYY (25-36)	RHYLQVVTRQRY (SEQ ID NO: 94)
	[Gln ²⁹ , Ile ³¹]PYY (25-36)	RHYLQLITRQRY (SEQ ID NO: 95)
	[Gln ²⁹ , Leu ³¹]PYY (25-36)	RHYLQLLTRQRY (SEQ ID NO: 96)
	[Gln ²⁹ , Ser ³²]PYY (25-36)	RHYLQLVSRQRY (SEQ ID NO: 97)
	[Gln ²⁹ , Leu ³³]PYY (25-36)	RHYLQLVTKQRY (SEQ ID NO: 98)
30	[Gln ²⁹ , Asn ³⁴]PYY (25-36)	RHYLQLVTRNRY (SEQ ID NO: 99)
	[Gln ²⁹ , Leu ³⁵]PYY (25-36)	RHYLQLVTRQKY (SEQ ID NO: 100)
	[Gln ²⁹ , Thr ³⁶]PYY (25-36)	RHYLQLVTRQRT (SEQ ID NO: 101)
	[Gln ²⁹ , Phe ³⁶]PYY (25-36)	RHYLQLVTRQRF (SEQ ID NO: 102)
	[Ile ³⁰ , Ile ³¹]PYY (25-36)	RHYLNITRQRY (SEQ ID NO: 103)
35	[Ile ³⁰ , Leu ³¹]PYY (25-36)	RHYLNILTRQRY (SEQ ID NO: 104)
	[Ile ³⁰ , Ser ³²]PYY (25-36)	RHYLNIVSRQRY (SEQ ID NO: 105)
	[Ile ³⁰ , Lys ³³]PYY (25-36)	RHYLNIVTKQRY (SEQ ID NO: 106)
	[Ile ³⁰ , Asn ³⁴]PYY (25-36)	RHYLNIVTRNRY (SEQ ID NO: 107)
	[Ile ³⁰ , Lys ³⁵]PYY (25-36)	RHYLNIVTRQKY (SEQ ID NO: 108)

-55-

	[Ile ³⁰ , Thr ³⁶]PYY (25-36)	RHYLNIVTRQRT (SEQ ID NO: 109)
	[Ile ³⁰ , Phe ³⁶]PYY (25-36)	RHYLNIVTRQRF (SEQ ID NO: 110)
	[Val ³⁰ , Ile ³¹]PYY (25-36)	RHYLNVITRQRY (SEQ ID NO: 111)
	[Val ³⁰ , Leu ³¹]PYY (25-36)	RHYLNVLTRQRY (SEQ ID NO: 112)
5	[Val ³⁰ , Ser ³²]PYY (25-36)	RHYLNVVSQRQRY (SEQ ID NO: 113)
	[Val ³⁰ , Lys ³³]PYY (25-36)	RHYLNVVTQKQRY (SEQ ID NO: 114)
	[Val ³⁰ , Asn ³⁴]PYY (25-36)	RHYLNVVTRNRY (SEQ ID NO: 115)
	[Val ³⁰ , Lys ³⁵]PYY (25-36)	RHYLNVVTRQKY (SEQ ID NO: 116)
	[Val ³⁰ , Thr ³⁶]PYY (25-36)	RHYLNVVTRQRT (SEQ ID NO: 117)
10	[Val ³⁰ , Phe ³⁶]PYY (25-36)	RHYLNVVTRQRF (SEQ ID NO: 118)
	[Ile ³¹ , Ser ³²]PYY (25-36)	RHYLNLISRQRY (SEQ ID NO: 119)
	[Ile ³¹ , Lys ³³]PYY (25-36)	RHYLNLITKQRY (SEQ ID NO: 120)
	[Ile ³¹ , Asn ³⁴]PYY (25-36)	RHYLNLITRNRY (SEQ ID NO: 121)
	[Ile ³¹ , Lys ³⁵]PYY (25-36)	RHYLNLITRQKY (SEQ ID NO: 122)
15	[Ile ³¹ , Thr ³⁶]PYY (25-36)	RHYLNLITRQRT (SEQ ID NO: 123)
	[Leu ³¹ , Phe ³⁶]PYY (25-36)	RHYLNLITRQRF (SEQ ID NO: 124)
	[Leu ³¹ , Ser ³²]PYY (25-36)	RHYLNLLSQRQRY (SEQ ID NO: 125)
	[Val ³¹ , Lys ³³]PYY (25-36)	RHYLNLITTKQRY (SEQ ID NO: 126)
	[Leu ³¹ , Asn ³⁴]PYY (25-36)	RHYLNLITRNRY (SEQ ID NO: 127)
20	[Leu ³¹ , Lys ³⁵]PYY (25-36)	RHYLNLITRQKY (SEQ ID NO: 128)
	[Leu ³¹ , Thr ³⁶]PYY (25-36)	RHYLNLITRQRT (SEQ ID NO: 129)
	[Leu ³¹ , Phe ³⁶]PYY (25-36)	RHYLNLITRQRF (SEQ ID NO: 130)
	[Ser ³² , Lys ³³]PYY (25-36)	RHYLNLVSKQRY (SEQ ID NO: 131)
	[Ser ³² , Asn ³⁴]PYY (25-36)	RHYLNLVSRNRY (SEQ ID NO: 132)
25	[Ser ³² , Lys ³⁵]PYY (25-36)	RHYLNLVSRQKY (SEQ ID NO: 133)
	[Ser ³² , Thr ³⁶]PYY (25-36)	RHYLNLVSRQRT (SEQ ID NO: 134)
	[Ser ³² , Phe ³⁶]PYY (25-36)	RHYLNLVSRQRY (SEQ ID NO: 135)
	[Lys ³³ , Asn ³⁴]PYY (25-36)	RHYLNLVTKNRY (SEQ ID NO: 136)
	[Lys ³³ , Lys ³⁵]PYY (25-36)	RHYLNLVTKQKY (SEQ ID NO: 137)
30	[Lys ³³ , Thr ³⁶]PYY (25-36)	RHYLNLVTKQRT (SEQ ID NO: 138)
	[Lys ³³ , Phe ³⁶]PYY (25-36)	RHYLNLVTKQRF (SEQ ID NO: 139)
	[Asn ³⁴ , Lys ³⁵]PYY (25-36)	RHYLNLVTRNRY (SEQ ID NO: 140)
	[Asn ³⁴ , Thr ³⁶]PYY (25-36)	RHYLNLVTRNRT (SEQ ID NO: 141)
	[Asn ³⁴ , Phe ³⁶]PYY (25-36)	RHYLNLVTRNRF (SEQ ID NO: 142)
35	[Lys ³⁵ , Thr ³⁶]PYY (25-36)	RHYLNLVTRQKT (SEQ ID NO: 143)
	[Lys ³⁵ , Phe ³⁶]PYY (25-36)	RHYLNLVTRQKF (SEQ ID NO: 144)

Point Mutations of PYY(24-36)

-56-

PEPTIDE	SEQUENCE
PYY(24-36)	LRHYLNVLVTRQRY (SEQ ID NO: 145)
[Ile ²⁴]PYY(24-36)	IRHYLNVLVTRQRY (SEQ ID NO: 146)
[Val ²⁴]PYY(24-36)	VRHYLNVLVTRQRY (SEQ ID NO: 147)

5

Also included as PYY(24-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of any of these three mutants with any of the above listed mutants for PYY(25-36), e.g., [Lys²⁵]PPY(24-36) (Amino acid sequence=LKHYLNVLVTRQRY (SEQ ID NO: 191)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 145.

10

Point Mutations of PYY(23-36)

PEPTIDE	SEQUENCE
PYY(23-36)	SLRHYNLNVLVTRQRY (SEQ ID NO: 148)
[Thr ²³]PYY(23-36)	TLRHYNLNVLVTRQRY (SEQ ID NO: 149)

Also included as PYY(23-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(24-36), e.g., [Lys²⁵]PPY(23-36) (Amino acid sequence=SLKHYLNVLVTRQRY (SEQ ID NO: 192)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 148.

20

Point Mutations of PYY(22-36)

PEPTIDE	SEQUENCE
PYY(22-36)	ASLRHYLNLNVLVTRQRY (SEQ ID NO: 150)
[Ser ²²]PYY(22-36)	SSLRHYLNLNVLVTRQRY (SEQ ID NO: 151)

Also included as PYY(22-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(23-36), e.g., [Lys²⁵]PPY(22-36) (Amino acid sequence=ASLKHYLNLNVLVTRQRY (SEQ ID NO: 193)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 150.

30

-57-

Point Mutations of PYY(21-36)

PEPTIDE	SEQUENCE
PYY(21-36)	YASLRHYLNLVTRQRY (SEQ ID NO: 152)
5 [Thr ²¹]PYY(21-36)	TASLRHYLNLVTRQRY (SEQ ID NO: 153)
[Phe ²¹]PYY(21-36)	FASLRHYLNLVTRQRY (SEQ ID NO: 154)

Also included as PYY(21-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of any of these three mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(22-36), e.g., [Lys²⁵]PPY(21-36) (Amino acid sequence=YASLKHYLNLVTRQRY (SEQ ID NO: 194)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 152.

15 Point Mutations of PYY(20-36)

PEPTIDE	SEQUENCE
PYY(20-36)	YYASLRHYLNLVTRQRY (SEQ ID NO: 155)
[Thr ²⁰]PYY(20-36)	TYASLRHYLNLVTRQRY (SEQ ID NO: 156)
[Phe ²⁰]PYY(20-36)	FYASLRHYLNLVTRQRY (SEQ ID NO: 157)

20 Also included as PYY(20-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of any of these three mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(21-36), e.g., [Lys²⁵]PPY(20-36) (Amino acid sequence=YYASLKHYLNLVTRQRY (SEQ ID NO: 195)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 155.

Point Mutations of PYY(19-36)

PEPTIDE	SEQUENCE
PYY(19-36)	RYYASLRHYLNLVTRQRY (SEQ ID NO: 158)
[Lys ¹⁹]PYY(19-36)	KYYASLRHYLNLVTRQRY (SEQ ID NO: 159)

Also included as PYY(19-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two

-58-

mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(20-36), e.g., [Lys²⁵]PPY(19-36) (Amino acid sequence=RYYASLKHYNLNLVTRQRY (SEQ ID NO: 196)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 158.

5

Point Mutations of PYY(18-36)

PEPTIDE	SEQUENCE
PYY(18-36)	NRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 160)
[Gln ¹⁸]PYY(18-36)	QRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 161)

10

Also included as PYY(18-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(19-36), e.g., [Lys²⁵]PPY(18-36) (Amino acid sequence=NRYYASLKHYNLNLVTRQRY (SEQ ID NO: 197)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 160.

Point Mutations of PYY(17-36)

PEPTIDE	SEQUENCE
PYY(17-36)	LNRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 162)
[Ile ¹⁷]PYY(17-36)	INRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 163)
[Val ¹⁷]PYY(17-36)	VNRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 164)

Also included as PYY(17-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of any of these three mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(18-36), e.g., [Lys²⁵]PPY(17-36) (Amino acid sequence=LNRYYASLKHYNLNLVTRQRY (SEQ ID NO: 198)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 162.

30

Point Mutations of PYY(16-36)

PEPTIDE	SEQUENCE
PYY(16-36)	ELNRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 165)
[Asp ¹⁶]PYY(16-36)	DLNRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 166)

Also included as PYY(16-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the 5 above listed mutants for PYY(17-36), e.g., [Lys^{25}]PPY(16-36) (Amino acid sequence=ELNRYYASLKHYNLNLVTRQRY (SEQ ID NO: 199)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 165.

Point Mutations of PYY(15-36)

10	PEPTIDE	SEQUENCE
	PYY(15-36)	EELNRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 167)
	[Asp^{15}]PYY(15-36)	DELNRYASLRHYLNLYNLVTRQRY (SEQ ID NO: 168)

Also included as PYY(15-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the 15 above listed mutants for PYY(16-36), e.g., [Lys^{25}]PPY(15-36) (Amino acid sequence=EELNRYYASLKHYNLNLVTRQRY (SEQ ID NO: 200)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 167.

20

Point Mutations of PYY(14-36)

PEPTIDE	SEQUENCE
PYY(14-36)	PEELNRYYASLRHYLNLYNLVTRQRY (SEQ ID NO: 169)

25 Also included as PYY(14-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of this PYY(14-36) mutant with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(15-36), e.g., [Lys^{25}]PPY(23-36) (Amino acid sequence=PEELNRYYASLKHYNLNLVTRQRY (SEQ ID NO: 201)) would result 30 from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 169.

Point Mutations of PYY(13-36)

PEPTIDE	SEQUENCE

-60-

PYY(13-36)	SPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 170)
[Thr ¹³]PYY(13-36)	TPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 171)

Also included as PYY(13-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(14-36), e.g., [Lys²⁵]PPY(13-36) (Amino acid sequence=SEELNRYYASLKHYNLVTRQRY (SEQ ID NO: 202)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 170.

10

Point Mutations of PYY(12-36)

PEPTIDE	SEQUENCE
PYY(12-36)	ASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 172)
[Ser ¹²]PYY(12-36)	SSPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 173)

15

Also included as PYY(12-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(13-36), e.g., [Lys²⁵]PPY(12-36) (Amino acid sequence=ASEELNRYYASLKHYNLVTRQRY (SEQ ID NO: 203)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 172.

Point Mutations of PYY(11-36)

PEPTIDE	SEQUENCE
PYY(11-36)	DASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 174)
[Glu ¹¹]PYY(11-36)	EASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 175)

Also included as PYY(12-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(12-36), e.g., [Lys²⁵]PPY(11-36) (Amino acid sequence=DASEELNRYYASLKHYNLVTRQRY (SEQ ID NO: 204)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 174.

-61-

Point Mutations of PYY(10-36)

PEPTIDE	SEQUENCE
PYY(10-36)	EDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 176)
[Asp ¹⁰]PYY(10-36)	DDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 177)

5

Also included as PYY(10-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(11-36), e.g., [Lys²⁵]PPY(10-36) (Amino acid

10 sequence=EDASEELNRYYASLKHYLNVLTRQRY (SEQ ID NO: 205)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 176.

Point Mutations of PYY(9-36)

PEPTIDE	SEQUENCE
PYY(9-36)	GEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 178)

Also included as PYY(9-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of this PPY(9-36) mutant with any of the above listed mutants for PYY(25-36), and/or any of the above listed 20 mutants for PYY(10-36), e.g., [Lys²⁵]PPY(9-36) (Amino acid sequence=GEDASPEELNRYYASLKHYLNVLTRQRY (SEQ ID NO: 206)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 178.

25 Potin Mutations of PYY(8-36)

PEPTIDE	SEQUENCE
PYY(8-36)	PGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 179)

Also included as PYY(8-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of this PPY(8-36) mutant with any of the above listed mutants for PYY(25-36), and/or any of the above listed 30 mutants for PYY(9-36), e.g., [Lys²⁵]PPY(8-36) (Amino acid sequence= SEQ ID NO: 207)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 179.

-62-

Point Mutations of PYY(7-36)

PEPTIDE	SEQUENCE
PYY(7-36)	APGEDASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 180)
5 [Ser ⁹]PYY(7-36)	SPGEDASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 181)

Also included as PYY(7-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the 10 above listed mutants for PYY(8-36), e.g., [Lys²⁵]PPY(7-36) (Amino acid sequence=APGEDASEELNRYYASLKHYNLNLVTRQRY (SEQ ID NO: 208)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 180.

15 Point Mutations of PYY(6-36)

PEPTIDE	SEQUENCE
PYY(6-36)	EAPGEDASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 182)
[Asp ⁶]PYY(6-36)	DAPGEDASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 183)

20 Also included as PYY(6-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of either of these two mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(7-36), e.g., [Lys²⁵]PPY(6-36) (Amino acid sequence=EAPGEDASEELNRYYASLKHYNLNLVTRQRY (SEQ ID NO: 209)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO: 182.

Point Mutations of PYY(5-36)

PEPTIDE	SEQUENCE
30 PYY(5-36)	PEAPGEDASPEELNRYYASLRHYLNLVTRQRY (SEQ ID NO: 184)

Also included as PYY(5-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of this PPY(5-36) mutant

-63-

with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(6-36), e.g., [Lys²⁵]PPY(5-36) (Amino acid sequence=PEAPGEDASPEELNRYYASLKHYNLNVLTRQRY (SEQ ID NO: 210)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO:

5 184.

Point Mutations of PYY(4-36)

PEPTIDE	SEQUENCE
PYY(4-26)	KPEAPGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 185)
10 [Arg ⁴]PYY(4-36)	RPEAPGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 186)
[Gln ⁴]PYY(4-36)	QPEAPGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 187)
[Asn ⁴]PYY(4-36)	NPEAPGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 188)

Also included as PYY(4-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of any of these four mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(5-36), e.g., [Lys²⁵]PPY(4-36) (Amino acid sequence=KPEAPGEDASEELNRYYASLKHYNLNVLTRQRY (SEQ ID NO: 211)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO:

20 185.

Point Mutations of PYY(3-36)

PEPTIDE	SEQUENCE
PYY(3-36)	IKPEAPGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 1)
25 [Leu ³]PYY(3-36)	LKPEAPGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 189)
[Val ³] PYY(3-36)	VKPEAPGEDASPEELNRYYASLRHYLNVLTRQRY (SEQ ID NO: 190)

Also included as PYY(3-36) mutations are polypeptide variations (amino acid sequence variations) resulting from the combination of any of these three mutants with any of the above listed mutants for PYY(25-36), and/or any of the above listed mutants for PYY(4-36), e.g., [Lys²⁵]PPY(3-36) (Amino acid sequence=IKPEAPGEDASEELNRYYASLKHYNLNVLTRQRY (SEQ ID NO: 212)) would result from combining the mutations from SEQ ID NO: 36 with SEQ ID NO:

1.

Also contemplated are PYY agonists (NPY analogs) having the formula:

5 X-Q-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-Leu-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-Arg-R₃₄-
Arg-R₃₆-Y

wherein X is H or C^a Me or N^a Me or desamino or an acyl group having 7 carbon atoms or less; Q is R₁₇-R₁₈, R₁₈ or desQ; R₁₇ is Met, Arg, Nle, Nva, Leu, Ala or D-Ala; R₁₈ is Ala, Ser, Ile, D-Ala, D-Ser or D-Ile; R₁₉ is Arg, Lys or Gln; R₂₀ is Tyr or Phe; R₂₁ is Tyr, Glu, His or Ala; R₂₂ is Ser, Ala, Thr, Asn or Asp; R₂₃ is Ala, Asp, Glu, Gln, Asn or Ser; R₂₅ is Arg or Gln; R₂₆ is His, Arg or Gln; R₂₇ is Phe or Tyr; R₂₈ is Ile, Leu, Val or Arg; R₂₉ is Asn or Ile; R₃₀ is Leu, Met, Thr or Val; R₃₁ is Ile, Val or Leu; R₃₂ is Thr or Phe; R₃₄ is Gln, Pro or His; R₃₆ is Phe or Tyr; and Y is 10
15 NH₂ or OH; provided that when Q is R₁₈, then at least one of R₂₇ and R₃₆ is Phe.

Analogs of NPY have the following applications: potent postsynaptic treatment of hypertension and cardiogenic shock, the treatment of acute cardiovascular circulatory failure, and the elevation of intracellular calcium. See U.S. Patent No. 5,026,685.

20 Certain preferred NPY analogs have the formula: X-R₁₈-Arg-Tyr-Tyr-R₂₂-R₂₃-Leu-Arg-His-Tyr-R₂₈-Asn-Leu-R₃₁-Thr-Arg-Gln-Arg-Tyr-NH₂, wherein X is H or C^a Me or N^a Me or desamino or an acyl group having 7 carbon atoms or less; R₁₈ is Ala or Ser; R₂₂ is Ser or Ala; R₂₃ is Ala or Ser; R₂₇ is Phe or Tyr; R₂₈ is Ile or Leu; R₃₁ is Ile or Val; and R₃₆ is Phe or Tyr; provided that at least one of R₂₇ and R₃₆ is 25
25 Phe. See U.S. Patent No. 5,026,685.

Other contemplated NPY analogs have the formula:

X-R₁₇-R₁₈-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-R₂₇-Ile-Asn-Leu-Ile-Thr-Arg-Gln-
30 Arg-R₃₆-NH₂,

wherein R₁₇ is Arg or Leu and R₁₈ is Ser or Ala or Ile; and wherein X, R₂₇ and R₃₆ are as previously indicated.

Still other preferred NPY analogs have the formula:

X-R₁₈-Arg-Tyr-Tyr-Ala-Ser-Leu-R₂₅-His-R₂₇-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-

5 R₃₆-NH₂,

wherein X is desamino or C^a Me or N^a Me and wherein R₁₈, R₂₅, R₂₇ and R₃₆ are as previously indicated.

10 Examples of such NPY agonists include:

pNPY (17-36) having the formula:

H-Leu-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 217)

15

The peptide hNPY (17-36) having the formula:

H-Met-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 218)

20

The peptide [Phe²⁷]-NPY (18-36) having the formula:

H-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Phe-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 219)

The peptide [Ac-D-Ala¹⁷]-NPY (17-36) having the formula:

25 Ac-D-Ala-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 220)

- The peptide NPY (19-36) having the formula:

30 H-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 221)

The peptide [Nle¹⁷]-NPY (17-36) having the formula:

-66-

H-Nle-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 222)

The peptide [D-Ser¹⁸]-NPY (18-36) having the formula:

5 H-D-Ser-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 223)

The peptide [Ala¹⁷, His²¹]-NPY (17-36) having the formula:

H-Ala-Ala-Arg-Tyr-His-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-10 Arg-Tyr-NH₂ (SEQ ID NO: 224)

The peptide [D-Ile¹⁸]-NPY (18-36) having the formula:

D-Ile-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 225)

15 The peptide [Ac-Arg¹⁷]-NPY (17-36) having the formula:
Ac-Arg-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 226)

20 The peptide [Gln¹⁹]-NPY (19-36) having the formula:
H-Gln-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 227)

25 The peptide [Phe²⁰]-NPY (18-36) having the formula:
H-Ala-Arg-Phe-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 228)

30 The peptide [C^a MeLeu¹⁷]-pNPY (17-36) having the formula:
H-C^a MeLeu-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 229)

The peptide [N^a MeLeu¹⁷]-pNPY (17-36) having the formula:

H-N⁹ MeLeu-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 230)

The peptide [desamino Ala¹⁸]-NPY (18-36) having the formula:
5 desamino-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 231)

The peptide [For-Ala¹⁸, Glu²³, Arg²⁶]-NPY (18-36) having the formula:
For-Ala-Arg-Tyr-Tyr-Ser-Glu-Leu-Arg-Arg-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-
10 Tyr-NH₂ (SEQ ID NO: 232)

The peptide [Nva¹⁷, Ala²¹, Leu²⁸]-NPY (17-36) having the formula:
H-Nva-Ala-Arg-Tyr-Ala-Ser-Ala-Leu-Arg-His-Tyr-Leu-Asn-Leu-Ile-Thr-Arg-Gln-
Arg-Tyr-NH₂ (SEQ ID NO: 233)
15 The peptide [Thr²², Gln²³]-NPY (18-36) having the formula:
H-Ala-Arg-Tyr-Tyr-Thr-Gln-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-
Tyr-NH₂ (SEQ ID NO: 234)

20 The peptide [desamino Leu¹⁷, Asn²³, Val³⁰]-NPY (17-36) having the
formula:
H-desamino Leu-Ala-Arg-Tyr-Tyr-Ser-Asn-Leu-Arg-His-Tyr-Ile-Asn-Val-Ile-Thr-
Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 235)

25 The peptide [Asp²², Ser²³, Thr³⁰]-NPY (18-36) having the formula:
H-Ala-Arg-Tyr-Tyr-Asp-Ser-Leu-Arg-His-Tyr-Ile-Asn-Thr-Ile-Thr-Arg-Gln-Arg-
Tyr-NH₂ (SEQ ID NO: 236)

30 The peptide [Gln²⁵, Leu³¹, Pro³⁴]-NPY (18-36) having the formula:
H-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Gln-His-Tyr-Ile-Asn-Leu-Leu-Thr-Arg-Pro-Arg-
Tyr-NH₂ (SEQ ID NO: 237)

-68-

The peptide [Gln² Phe³⁶]-NPY (17-36) having the formula:

H-Leu-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-Gln-Tyr-Arg-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Phe-NH₂ (SEQ ID NO: 238)

5 The peptide [Phe³⁶]-pPYY (19-36) having the formula:

H-Arg-Tyr-Tyr-Ala-Ser-Leu-Arg-His-Tyr-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Phe-NH₂ (SEQ ID NO: 239)

The peptide pPYY (18-36) having the formula:

10 H-Ser-Arg-Tyr-Tyr-Ala-Ser-Leu-Arg-His-Tyr-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 240)

The peptide [Ac-Ser¹⁸, Phe²⁷]-pPYY (18-36) having the formula:

15 Ac-Ser-Arg-Tyr-Tyr-Ala-Ser-Leu-Arg-His-Phe-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 241)

The peptide [Nle¹⁷, Asn²², Phe²⁷]-NPY (17-36) having the formula:

H-Nle-Ala-Arg-Tyr-Tyr-Asn-Ala-Leu-Arg-His-Phe-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 242)

20 The peptide [D-Ala¹⁸, Glu²¹, His³⁴]-NPY (18-36) having the formula:

H-D-Ala-Arg-Tyr-Glu-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-His-Arg-Tyr-NH₂ (SEQ ID NO: 243)

25 The peptide [Bz-Leu¹⁷, Pro³⁴, Phe³⁶]-pNPY (17-36) having the formula:

Bz-Leu-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Pro-Arg-Phe-NH₂ (SEQ ID NO: 244)

The peptide [Lys¹⁹, Phe²⁷, Val²⁸]-NpY (18-36) having the formula:

30 H-Ala-Lys-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Phe-Val-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 245)

-69-

The peptide [D-Ala¹⁷, Val²⁸, Phe³²]-NPY (17-36) having the formula:
D-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Val-Asn-Leu-Ile-Phe-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 246)

5 The peptide [C^a MeSer¹⁸, Met³⁰, Phe³⁶]-NPY (18-36) having the formula:
H-C^a MeSer-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Met-Ile-Thr-Arg-Gln-Arg-Phe-NH₂ (SEQ ID NO: 247)

10 The peptide [Arg¹⁷, Ile¹⁸, Phe^{27,36}]-NPY (17-36) having the formula:
H-Arg-Ile-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Phe-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Phe-NH₂ (SEQ ID NO: 248)

15 The peptide [Ser¹⁸, Phe²⁷]-pNPY (17-36) having the formula:
H-Leu-Ser-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Phe-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 249)

20 The peptide [N^a Melle¹⁸, Gln²⁵, Phe²⁷]-NPY (18-36) having the formula:
N^a Melle-Arg-Tyr-Tyr-Ser-Ala-Leu-Gln-His-Phe-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 250)

25 The peptide [D-Ser¹⁸, Phe³⁶]-NPY (18-36) having the formula:
H-D-Ser-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Phe-NH₂ (SEQ ID NO: 251)

30 The peptide [Asp²³, Arg²⁶]hNPY (17-36) having the formula:
H-Met-Ala-Arg-Tyr-Tyr-Ser-Asp-Leu-Arg-Arg-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 252)


The peptide [Glu²³, Ile²⁹]-NPY (18-36) having the formula:
30 H-Ala-Arg-Tyr-Tyr-Ser-Glu-Leu-Arg-His-Tyr-Ile-Ile-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ ID NO: 253)

-70-

The peptide [D-Ala¹⁷]-NPY(17-36)-OH having the formula:

D-Ala-Ala-Arg-Tyr-Tyr-Ser-Ala-Leu-Arg-His-Tyr-Ile-Asn-Leu-Ile-Thr-Arg-Gln-Arg-Tyr-OH (SEQ ID NO: 254).

5 Other peptide YY agonists have the formula:



10

wherein:

X is a chain of 0-5 amino acids, inclusive, the N-terminal one of which is bonded to R₁ and R₂

15

Y is a chain of 0-4 amino acids, inclusive, the C-terminal one of which is bonded to R₃ and R₄

20

R₁ is H, C₁-C₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl);

R₂ is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl);

25

A²² is an aromatic amino acid, Ala, Aib, Anb, N-Me-Ala, or is deleted;

A²³ is Ser, Thr, Ala, N-Me-Ser, N-Me-Thr, N-Me-Ala, or is deleted;

A²⁴ is Leu, Ile, Vat, Trp, Gly, Aib, Anb, N-Me-Leu, or is deleted;

30

A²⁵ is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys-ε-NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or an aryl group), Orn, or is deleted;

A²⁶ is His, Thr, 3-Me-His, 1-Me-His, β-pyrozolylalanine, N-Me-His, Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys-ε-NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or an aryl group), Orn, or is deleted;

35

A²⁷ is an aromatic amino acid other than Tyr;

A²⁸ is Leu, Ile, Vat, Trp, Aib, Aib, Anb, or N-Me-Leu;

A²⁹ is Asn, Ala, Gln, Gly, Trp, or N-Me-Asn;

A³⁰ is Leu, Ile, Val, Trp, Aib, Anb, or N-Me-Leu;

A³¹ is Vat, Ile, Trp, Aib, Anb, or N-Me-Val;

-71-

A^{32} is Thr, Ser, N-Me-Ser, or N-Me-Thr;

R_3 is H, C_1 - C_{12} alkyl (e.g., methyl), C_6 - C_{18} aryl (e.g., phenyl,


naphthaleneacetyl), C_1 - C_{12} acyl (e.g., formyl, acetyl, and myristoyl), C_7 - C_{18} aralkyl (e.g., benzyl), or C_7 - C_{18} alkaryl (e.g., p-methylphenyl);

5 R_4 is H, C_1 - C_{12} alkyl (e.g., methyl), C_6 - C_{18} aryl (e.g., phenyl, naphthaleneacetyl), C_1 - C_{12} acyl (e.g., formyl, acetyl, and myristoyl), C_7 - C_{18} aralkyl (e.g., benzyl), or C_7 - C_{18} alkaryl (e.g., p-methylphenyl), or a pharmaceutically acceptable salt thereof. See U.S. Patent No. 5,574,010.

10 Particularly preferred agonists of this formula to be used in the method of the disclosure include:

$N\text{-}\alpha\text{-Ala-Ser-Leu-Arg-His-Trp-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Tyr-NH}_2$ (SEQ ID NO: 255).

15 Other peptide YY agonists have the formula:



20 wherein:

the N-terminal amino acid bonds to R_1 and R_2 ;

Y is a chain of 0-4 amino acids, inclusive the C-terminal one of which bonds to R_3 and R_4 ;

25 R_1 is H, C_1 - C_{12} alkyl, C_6 - C_{18} aryl, C_1 - C_{12} acyl, C_7 - C_{18} aralkyl, or C_7 - C_{18} alkaryl;

R_2 is H, C_1 - C_{12} alkyl, C_6 - C_{18} aryl, C_1 - C_{12} acyl, C_7 - C_{18} aralkyl, or C_7 - C_{18} alkaryl;

25 A^{25} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C_1 - C_{10} alkyl group, or an aryl group), Orn, or is deleted;

30 A^{26} is Ala, His, Thr, 3-Me-His, 1-Me-His, β -pyrozolylalanine, N-Me-His, Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C_1 - C_{10} alkyl group, or an aryl group), Orn or is deleted;

A^{27} is an aromatic amino acid;

-72-

A^{28} is Leu, Ile, Val, Trp, Aib, Anb, or N-Me-Leu:

A^{29} is Asn, Ala, Gln, Gly, Trp, or N-Me-Asn:

A^{30} is Leu, Ile, Val, Trp, Aib, α -NBu, or N -Me-Leu.

A^{31} is Val, Ile, Thr, Aib, Aph, or N-Me-Val:


5 A^{32} is Thr, Ser, N-Me-Ser, or N-Me-Thr or D-Tyr:

R₃ is H, C₁-C₁₂ alkyl, C₆-C₁₈ aryl, C₁-C₁₂ acyl, C₇-C₁₈ aralkyl, or C₇-C₁₈ alkaryl; and

R₄ is H, C₁ -C₁₂ alkyl, C₆ -C₁₈ aryl, C₁ -C₁₂ acyl, C₇ -C₁₈ aralkyl, or C₇ -C₁₈ alkaryl, or a pharmaceutically acceptable salt thereof. Note that, unless indicated otherwise, for all peptide YY agonists described herein, each amino acid residue, e.g., Leu and A¹, represents the structure of NH--C(R)H--CO--, in which R is the side chain. Lines between amino acid residues represent peptide bonds which join the amino acids. Also, where the amino acid residue is optically active, it is the L-form configuration that is intended unless D-form is expressly designated.

15

Other PYY agonists have the formula:



wherein:

X is a chain of 0-5 amino acids, inclusive, the N-terminal one of which is bonded to R_1 and R_2 :

Y is a chain of 0-4 amino acids, inclusive, the C-terminal one of which is bonded to R₃ and R₄;

25 R_1 is H, C₁-C₁₂ alkyl (e.g. methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl);

R_2 is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl);

A²² is an aromatic amino acid, Ala, Aib, Anb, N-Me-Ala, or is deleted:

-73-

A^{23} is Ser, Thr, Ala, Aib, N-Me-Ser, N-Me-Thr, N Me-Ala, or is deleted;
 A^{24} is Ieu, Ile, Val, Trp, Gly, Nle, Nva, Aib, Anb, N-Me-Leu, or is deleted;
 A^{25} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or an aryl group), Orn, or is deleted;
5 A^{26} is Ala, His, Thr, 3-Me-His, 1-Me-His, β -pyrozolylalanine, N-Me-His, Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl groups or an aryl group), Orn, or is deleted;
 A^{27} is an aromatic amino acid other than Tyr;
 A^{28} is Leu, Ile, Val, Trp, Nle, Nva, Aib, Anb, or N-Me-Leu;
10 A^{29} is Asn, Ala, Gin, Gly, Trp, or N-Me-Asn;
 A^{30} is Leu, Ile, Val, Trp, Nle, Nva, Aib, Anb, or N-Me-Leu;
 A^{31} is Val, Leu, Ile, Trp, Nle, Nva, Aib, Anb, or N-Me-Val;
 A^{32} is Thr, Ser, N-Me-Ser, N-Me-Thr, or D-Trp;
 R_3 is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl,
15 naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl); and
 R_4 is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl), or a pharmaceutically
20 acceptable salt thereof.
In preferred embodiments, A^{27} is Phe, Nal, Bip, Pcp, Tic, Trp, Bth, Thi, or Dip.
In preferred embodiments X is A^{17} - A^{18} - A^{19} - A^{20} - A^{21} wherein
 A^{17} is Cys, Leu, Ile, Val, Nle, Nva, Aib, Anb, or N-Me-Leu;
25 A^{18} is Cys, Ser, Thr, N-Me-Ser, or N-Me-Thr;
 A^{19} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or C₆-C₁₈ aryl group), Cys, or Orn;
 A^{20} is an aromatic amino acid, or Cys; and
 A^{21} is an aromatic amino acid, Cys, or a pharmaceutically acceptable salt
30 thereof. In yet other preferred embodiments, Y is A^{33} - A^{34} - A^{35} - A^{36} wherein
 A^{33} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or an aryl group), Cys, or Orn;

-74-

A^{34} is Cys, Gln, Asn, Ala, Gly, N-Me-Cln, Aib, or Anb;


A^{35} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or C₆-C₁₈ aryl group), Cys, or Orn; and

5 A³⁶ is an aromatic amino acid, Cys or a pharmaceutically acceptable salt thereof. See U.S. Patent No. 5,604,203.

Particular embodiments include compounds having the formula: N- α -Ac-Ala-Ser-Leu-Arg-His-Phe-Leu-Asn-Leu-Val-Thr-Arg-Gin-Arg-Tyr-NH₂ (SEQ. ID. NO: 325), H-Ala-Ser-Leu-Arg-His-Phe-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ. ID. NO: 326), N- α -Ac-Ala-Ser-Leu-Arg-Thr-Arg-Gin-Arg-Tyr-NH₂ (SEQ. ID. NO: 327), N- α -Ac-Ala-Ser-Leu-Arg-His-Thi-Leu-Asn-Leu-Val-Thr-Arg-Gin-Arg-Tyr-NH₂ (SEQ. ID. NO: 328), N- α -Ac-Tyr-Ser-Leu-Arg-His-Phe-Leu-Asn-Leu-Val-Thr-Arg-Gin-Arg-Tyr-NH₂ (SEQ. ID. NO: 329) or a pharmaceutically acceptable salt thereof.

15

Other PYY agonists have the formula:



wherein the N-terminal amino acid is bounded to R_1 and R_2 ; Y is a chain of 0-4 amino acids, inclusive the C-terminal one of which is bonded to R_3 and R_4 ;

R₁ is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl);

25 R₂ is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl);

A^{25} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or an aryl group), Orn, or is deleted;

-75-

A^{26} is Ala, His, Thr, 3-Me-His, 1-Me-His, β -pyrozolylalanine, N-Me-His, Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl groups or an aryl group), Orn, or is deleted;

A^{27} is an aromatic amino acid;

5 A^{28} is Leu, Ile, Val, Trp, Nle, Nva, Aib, Anb, or N-Me-Leu;

A^{29} is Asn, Ala, Gln, Gly, Trp, or N-Me-Asn;

A^{30} is Leu, Ile, Val, Trp, Nle, Nva, Aib, Anb, or N-Me-Leu;

A^{31} is Val, Ile, Trp, Nle, Nva, Aib, Anb, or N-Me-Val;

A^{32} is Thr, Ser, N-Me-Ser, N-Me-Thr, or D-Trp;

10 R_3 is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl); and

R_4 is H, C₁-C₁₂ alkyl (e.g., methyl), C₆-C₁₈ aryl (e.g., phenyl, naphthaleneacetyl), C₁-C₁₂ acyl (e.g., formyl, acetyl, and myristoyl), C₇-C₁₈ aralkyl (e.g., benzyl), or C₇-C₁₈ alkaryl (e.g., p-methylphenyl), or a pharmaceutically acceptable salt thereof. See U.S. Patent No. 5,604,203.

In particular embodiments, A^{27} is Phe, Nal, Bip, Pcp, Tic, Trp, Bth, Thi, or Dip.

In particular embodiments X is A^{33} - A^{34} - A^{35} - A^{36} wherein

20 A^{33} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or C₆-C₁₈ aryl group), Cys, or Orn;

A^{34} is Gln, Asn, Ala, Gly, N-Me-Gin, Aib, Cys, or Anb;

A^{35} is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R (where R is H, a branched or straight chain C₁-C₁₀ alkyl group, or C₆-C₁₈ aryl group), Cys, or Orn;

25 and

A^{36} is an aromatic amino acid, Cys, or a pharmaceutically acceptable salt thereof.

Preferably, the compound has the formula: N- α -Ac-Arg-His-Phe-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Tyr-NH₂ (SEQ. ID. NO: 324).

30

Exemplary PYY agonists include:

YPAKEAPGEDASPEELSTYYASLR [im-DNP-His²⁶]

(SEQ ID NO: 256)

YLNLVTRZRY-NH ₂	
PYY(22-36)	
ASLRHYLNLVTRQRY-NH ₂	(SEQ ID NO: 257)
[Ala ³²]PYY	
ASLRHYLNVL[Ala]RQRY-NH ₂	(SEQ ID NO: 258)
[Ala ^{23,32}]PYY	
A[Ala]LRHYLNVL[Ala]RQRY-NN ₂	(SEQ ID NO: 259)
[Glu ²⁸]PYY(22-36)	
ASLRHY[Glu]NLVTRQRY-NH ₂	(SEQ ID NO: 260)
N- α -Ac-PYY(22-36)	
N- α -Ac-ASLRHYLNLVTRORY-NH ₂	(SEQ ID NO: 261)
N- α -Ac[p.CL.Phe ²⁶]PYY	
N- α -Ac-ASLR[p.CL.Phe ²⁶]YLNLVTRQRY-NH ₂	(SEQ ID NO: 262)
N- α -Ac[Glu ²⁸]PYY	
N- α -Ac-ASLRHY[Glu]NLVTRQRY-NH ₂	(SEQ ID NO: 263)
N- α -Ac[Phe ²⁷]PYY	
N- α -Ac-ASLRH[Phe]ENLVTRQR[N-Me-Tyr]-NH ₂	(SEQ ID NO: 264)
N- α -Ac[8N-Me-Tyr ³⁶]PYY	
N- α -Ac-ASLRHYENLVTR0R[N-Me-Tyr]-NH ₂	(SEQ ID NO: 265)
N- α -myristoyl-PYY(2214 36)	
N- α -myristoyl-ASLRHYLNLVTRQRY-NH ₂	(SEQ ID NO: 266)
N- α -naphthateneacetyl-PYY(22-36)	
N- α -naphthateneacetyl-ASLRHYLNLVTRQRY-NH ₂	(SEQ ID NO: 267)
N- α -Ac[Phe ²⁷]PYY	
N- α -Ac-ASLRH[Phe]ENLVTR0R[N-Me-Tyr]-NH ₂	(SEQ ID NO: 268)
N- α -Ac-PYY (22-36)	
N- α -Ac-ASLRHYLNLVTRQRY-NH ₂	(SEQ ID NO: 269)
N- α -Ac-[Bth ²⁷]PYY (22-36)	
N- α -Ac-ASLRH[Bth]NLVTRQRY-NH ₂	(SEQ ID NO: 270)
N- α -Ac-[Bip ²⁷]PYY (22-36)	(SEQ ID NO: 271)
N- α -Ac-ASLRH[Bth]NLVTRQRY-NH ₂	(SEQ ID NO: 272)
N- α -Ac-[Nal ²⁷]PYY (22-36)	
N- α -Ac-ASLRH[Bth]NLVTRQRY-NH ₂	(SEQ ID NO: 273)
N- α -Ac-[Trp ²⁷]PYY (22-36)	(SEQ ID NO: 274)
N- α -Ac-ASLRH[Trp]NLVTRQRY-NH ₂	(SEQ ID NO: 275)
N- α -Ac-[Thi ²⁷]PYY (22-36)	
N- α -Ac-ASLRN[Thi]NLVTRQRY-NH ₂	(SEQ ID NO: 276)
N- α -Ac-[Tic ²⁷]PYY (22-36)	
N- α -Ac-ASLRH[Tic]NLVTRQRY-NH ₂	(SEQ ID NO: 277)
N- α -Ac-[Phe ²⁷]PYY (25-36)	
N- α -Ac-H[Phe]NLVTRQRY-NH ₂	(SEQ ID NO: 279)
N- α -Ac-[Phe ²⁷ ,Thi ²⁷]PYY (22-36)	
N- α -Ac-ASLRH[Phe]NLVTRQR[Thi]-NH ₂	(SEQ ID NO: 280)
N- α -Ac-[Thz ²⁶ ,Phe ²⁷]PYY (22-36)	
N- α -Ac-ASLRH[Thz][Phe]NLVTRQRY-NH ₂	(SEQ ID NO: 281)

N- α -Ac-[Phe ²⁷]PYY (22-36)	(SEQ ID NO: 282)
N- α -Ac-ASLRH[Thz][Phe]LNLVTRQRY-NH ₂	
N- α -Ac-[Phe ²⁷]PYY (22-36)	(SEQ ID NO: 289)
N- α -Ac-[Phe]SLRN[Phe]LNLVTRQRY-NH ₂	
N- α -Ac-[Tyr ²² ,Phe ²⁷]PYY (22-36)	(SEQ ID NO: 290)
N- α -Ac-[Tyr]SLRH[Phe]LNLVTRQRY-NH ₂	
N- α -Ac-[Trp ²⁸]PYY (22-36)	(SEQ ID NO: 291)
N- α -Ac-ASLRHY[Trp]NLVTRQRY-NH ₂	
N- α -Ac-[Trp ²⁸]PYY (22-36)	(SEQ ID NO: 292)
N- α -Ac-ASLRHYLN[Trp]VTRQRY-NH ₂	
N- α -Ac-[Ala ²⁶ ,Phe ²⁷]PYY (22-36)	(SEQ ID NO: 293)
N- α -Ac-ASLR[Ala][Phe]LNLVTRQRY-NH ₂	
N- α -Ac-[Bth ²⁷]PYY (22-36)	(SEQ ID NO: 294)
N- α -Ac-ASLR[Bth]LNLVTRQRY-NH ₂	
N- α -Ac-[Phe ²⁷]PYY (22-36)	(SEQ ID NO: 295)
N- α -Ac-ASLRH[Phe]LNLVTRQRY-NH ₂	
N- α -Ac-[Phe ^{27,36}]PYY (22-36)	(SEQ ID NO: 296)
N- α -Ac-ASLRH[Phe]LNLVTRQR[Phe]-NH ₂	
N- α -Ac-[Phe ²⁷ , D-Trp ³²]PYY (22-36)	(SEQ ID NO: 297)
N- α -Ac-ASLRH[Phe]LNLV[D-Trp]RQRY-NH ₂	

Other PYY agonists include neurophilic Y Y2 receptor specific peptides having the formula:

X1(-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14)_n-X15

5 wherein

X1 is NH, CH₃CO or one or two naturally occurring amino acids.

X2 is Leu, Ile or Val.

X3 is Arg, Lys or His.

X4 is His, Lys or Arg.

10 X5 is Tyr or Phe.

X6 is Leu, Ile or Val.

X7 is Asn or Gln.

X8 is Leu, Ile or Val.

X9 is Leu, Ile or Val.

15 X10 is Thr or Ser.

X11 is Arg, His or Lys.

X12 is Gln or Asn.

-78-

X13 is Arg, His or Lys.

X14 is Tyr or Phe.

X15 is COOH, NH₂ or one or two naturally occurring amino acids with the terminal amino acid being in the normal or carboxamide form; and

5 n is 1 to 5. See U.S. Patent No. 5,696,093.

Exemplary agonists include:

CH₃CO-L-R-H-Y-L-N-L-L-T-R-Q-R-Y-NH₂ (SEQ ID NO: 298)

CH₃CO-L-R-H-Y-I-N-L-I-T-R-Q-R-Y-NH₂ (SEQ ID NO: 299)

10 NH₂-L-R-H-Y-L-N-L-L-T-R-Q-R-Y-NH₂ (SEQ ID NO: 300)

NH₂-L-R-H-Y-I-N-L-I-T-R-Q-R-Y-NH₂ (SEQ ID NO: 301)

Other PYY agonists have the formula:

N- α -R¹-[Nle^{24,28,30}, Trp²⁷, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂,

15 N- α -R¹-[Nle^{24,28}, Trp^{27,30}, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂,

N- α -R¹-[Nle^{24,28,30}, Phe²⁷, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂,

N- α -R¹-[Nle^{24,28}, Phe²⁷, Trp³⁰, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂,

N- α -R¹-[Trp³⁰, ψ ^{35/36}]PYY(25-36)-NH₂,

N- α -R¹-[Trp³⁰]PYY(25-36)-NH₂,

20 N- α -R¹-[Nle^{24,28}, Trp³⁰, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂ and

N- α -R¹-[Nle²⁸, Trp³⁰, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂ or a pharmaceutically-acceptable salt thereof,

wherein R¹ is H, (C₁-C₁₂)alkyl or (C₁-C₁₂)acyl; and

ψ is a pseudopeptide bond selected from the group consisting of --CH₂--NH--

25 -, --CH₂--S--, --CH₂--CH₂--, --CH₂--O-- and --CH₂--CO--. See U.S. Patent No. 6,046,162.

Particular compounds of the immediately foregoing group of compounds are where R¹ is acetyl and ψ is --CH₂--NH--.

30 A particular group of compounds is selected from a group consisting of

N- α -Ac-[Nle^{24,28,30}, Trp²⁷, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂, (SEQ ID NO: 302)

-79-

N- α -Ac-[Nle^{24,28}, Trp^{27,30}, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂, (SEQ ID NO: 303)

N- α -Ac-[Nle^{24,28,30}, Phe²⁷, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂, (SEQ ID NO: 304)

5 N- α -Ac-[Nle^{24,28}, Phe²⁷, Trp³⁰, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂, (SEQ ID NO: 305)

N- α -Ac-[Trp³⁰, ψ ^{35/36}]PYY(25-36)-NH₂, (SEQ ID NO: 306)

N- α -Ac-[Trp³⁰]PYY(25-36)-NH₂ (SEQ ID NO: 307) and


N- α -Ac-[Nle²⁸, Trp³⁰, Nva³¹, ψ ^{35/36}]PYY(22-36)-NH₂, (SEQ ID NO: 308) or

10 a pharmaceutically acceptable salt thereof.

Another particular compound has the formula N- α -Ac-[Nle^{24,28}, Trp³⁰, Nva.sup.³¹, ψ ^{35/36}]PYY(22-36)-NH₂ (SEQ. ID. NO: 309) or a pharmaceutically acceptable salt thereof.

Another PYY agonist has the formula (A),

15



20

having one or two pseudopeptide bonds where each pseudopeptide bond is independently selected from the group consisting of --CH₂--NH--, --CH₂--S--, --CH₂--CH₂--, --CH₂--O-- and -CH₂--CO--; wherein:

25

R¹⁰ is a chain of 0-5 amino acids, inclusive, where the N-terminal amino acid is bonded to R¹ and R² by the side chain of the N-terminal amino acid or by the nitrogen of the amino group of the N-terminal amino acid;

R²⁰ is a chain of 0-4 amino acids, inclusive, where the C-terminal amino acid is bonded to R³ and R⁴ by the side chain of the C-terminal amino acid or by the carbon of the carboxyl group of the C-terminal amino acid;

30

R¹, R², R³ and R⁴ are each independently selected from the group consisting of H, (C₁-C₁₂)alkyl, (C₆-C₁₈)aryl, (C₁-C₁₂)acyl, phenyl(C₁-C₁₂)alkyl and ((C₁-C₁₂)alkyl)₁₋₅-phenyl;

A²² is an aromatic amino acid, Ala, Aib, Anb, N-Me-Ala or is deleted;

A²³ is Ser, Thr, Ala, N-Me-Ser, N-Me-Thr, N-Me-Ala or is deleted;

-80-

A²⁴ is Leu, Ile, Nle, Val, Trp, Gly, Aib, Anb, N-Me-Leu or is deleted;
A²⁵ is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys-p.epsilon.-NH-Z, Orn or is deleted;
A²⁶ is His, Thr, 3-Me-His, 1-Me-His, β -pyrazolylalanine, N-Me-His, Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-Z, Orn or is deleted;
A²⁸ is Leu, Ile, Nle, Val, Trp, Aib, Anb or N-Me-Leu;
A²⁹ is Asn, Ala, Gln, Gly, Trp or N-Me-Asn;
A³⁰ is Leu, Ile, Nle, Fla, Val, Trp, Aib, Anb or N-Me-Leu;
A³¹ is Val, Nva, Ile, Trp, Aib, Anb or N-Me-Val; and
A³² is Thr, Ser, N-Me-Ser or N-Me-Thr;
where Z for each occurrence is independently selected from the group consisting of H, (C₁ -C₁₀)alkyl and (C₆ -C₁₈)aryl; or a pharmaceutically acceptable salt thereof. See U.S. Patent No. 6,046,167.
A particular group of compounds of the immediately foregoing group of compounds is where R¹⁰ is A¹⁷ -A¹⁸ -A¹⁹ -A²⁰ -A²¹;
where A¹⁷ is Cys, Leu, Ile, Val, Nle, Nva, Aib, Anb or N-Me-Leu;
A¹⁸ is Cys, Ser, Thr, N-Me-Ser or N-Me-Thr;
A¹⁹ is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R⁵, Cys or Orn;
A²⁰ is an aromatic amino acid or Cys;
A²¹ is an aromatic amino acid or Cys;
R²⁰ is A³³ -A³⁴ -A³⁵ -A³⁶,
A³³ is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R⁵, Cys or Orn;
A³⁴ is Cys, Gin, Asn, Ala, Gly, N-Me-Gln, Aib or Anb;
A³⁵ is Arg, Lys, homo-Arg, diethyl-homo-Arg, Lys- ϵ -NH-R⁵, Cys or Orn; and
A³⁶ is an aromatic amino acid or Cys;
where R⁵ for each occurrence is independently selected from the group consisting of H₁ (C₁ -C₁₀)alkyl and (C₆ -C₁₈) aryl.

A particular group of compounds of the foregoing group of compounds are the compounds of the formula $N\text{-}\alpha\text{-Ac-[Fla}^{27}\text{]} \text{PYY(25-36)-NH}_2$ and $N\text{-}\alpha\text{-Ac-[Fla}^{27}\text{]} \text{PYY(22-36)-NH}_2$ or a pharmaceutically acceptable salt thereof.

5 Another group of PYY agonist has the formula:

$$(I) \quad (R^1 R^2) - A^1 - A^2 - A^3 - A^4 - A^5 - A^6 - A^7 - A^8 - A^9 - A^{10} - R^{30}$$

$$(III) \quad (R^1 R^2) \cdot [A^5 - A^6 - A^7 - A^8 - A^9 - A^{10}]_m R^{30},$$

20 or a pharmaceutically acceptable salt thereof wherein

-----represents an optional bond between the amino acids shown connected where each bond is independently selected from the group consisting of --S--S-- only when the amino acids connected are Cys-Cys, -CO-NH-, -CH₂-NH- and

provided that when the optional bond is

35
$$\begin{array}{c}
 -\text{HN}-\text{CH}-\text{Co}- \\
 | \\
 (\text{CH}_2)_4 \\
 | \\
 -\text{HN}-\text{CH}-\text{CO}- \\
 | \\
 \text{R}
 \end{array}$$

40 it replaces the two amino acids that the optional bond is attached to; q is 1-4; m is 1 to 4;

R^{30} is OH or -O-R¹, provided that when A¹ to A⁷ are deleted then R^{30} is also NH-R¹, where R³⁰ is attached to the carbon atom of the carboxyl of the C-terminal amino acid;

5 R¹ and R² for each occurrence are each independently selected from the group consisting of H, (C₁ -C₁₂)alkyl, (C₆ -C₁₈)aryl, (C₁ -C₁₂)acyl, phenyl(C₁ -C₁₂)alkyl and ((C₁ -C₁₂)alkyl)₁₋₅-phenyl where R¹ and R² are attached to the nitrogen of the amine of the N-terminal amino acid;

10 A¹ is deleted or D- or L- of the following amino acids: Trp, Tyr, Fla, Bth, Nal, Tic, Tic-OH, Dip, Bip or optionally substituted Phe where the Phe is optionally substituted with one to five substituents selected from the group consisting of (C₁ -C₄)alkyl, halo, (C₁ -C₄)alkoxy, amino and nitro;

15 A² is deleted or D- or L- of the following amino acids: Ile, Val, Leu, Nle, Anb, Aib, Pro, Gln or Asn;

15 A³ is deleted or D- or L- of the following amino acids: Asn, Gln, Glu, Asp, Orn, Lys, Dpr or Cys;

20 A⁴ is deleted or D- or L- of the following amino acids: Ile, Val, Leu, Nle, Anb, Aib or Pro;

20 A⁵ is deleted or D- or L- of the following amino acids: Ile, Val, Leu, Nle, Anb, Aib, Pro, Glu, Asp, Orn, Lys, Dpr or Cys;

20 A⁶ is deleted or D- or L- of the following amino acids: Thr, Ser, Trp, Tyr, Fla, Bth, Nal, Tic, Tic-OH, Dip, Bip or optionally substituted Phe where the Phe is optionally substituted with one to five substituents selected from the group consisting of (C₁ -C₄)alkyl, halo, (C₁ -C₄)alkoxy, amino and nitro;

25 A⁷ is deleted or D- or L- of the following amino acids: Arg, Lys, homo-Arg, dialkyl-homo-Arg, Lys- ϵ -NH-R⁷ or Orn;

25 A⁸ is deleted or D- or L- of the following amino acids: Nva, Val, Ile, Leu, Nle, Anb, Aib, Pro, Gln, Asn, Glu, Asp, Orn, Lys, Dpr or Cys;

25 A⁹ is deleted or D- or L- of the following amino acids: Arg, Lys, homo-Arg, dialkyl-homo-Arg, Lys- ϵ -NH-R⁷ or Orn; and

30 A¹⁰ is deleted or D- or L- of the following amino acids: Tyr, Trp, Fla, Bth, Nal, Tic, Tic-OH, Dip, Bip, tyramine or optionally substituted Phe where the Phe is optionally substituted with one to five substituents selected from the group

-83-

consisting of (C₁ -C₄)alkyl, halo, (C₁ -C₄)alkoxy, amino and nitro, or the corresponding decarboxylated optionally substituted Phe;

where R⁷ for each occurrence is independently selected from the group consisting of H.sub.1 (C₁ -C₁₀)alkyl and (C₆ -C₁₈) aryl, provided that not all of A₁ to 5 A₁₀ are deleted at the same time. See U.S. Patent No. 6,046,167.

A particular group of compounds of the immediately foregoing group of compounds is

(SEQ ID NO: 310)

H--Ile--Asn--Pro--Ile--Tyr--Arg--Leu--Arg--Tyr--OMe

10

(SEQ ID NO: 311)

H--Ile--Asn--Pro--Cys--Tyr--Arg--Leu--Arg--Tyr--Ome

15

H--Ile--Asn--Pro--Cys--Tyr--Arg--Leu--Arg--Tyr--Ome,

(SEQ ID NO: 312)

H--Cys--Tyr--Arg--Leu--Arg--Tyr--Ome

20

H--Cys--Tyr--Arg--Leu--Arg--Tyr--OMe,

25

(SEQ ID NO: 313)

H--Ile--Asn--Pro--NH--CH--CO--Tyr--Arg--Leu--Arg--Tyr--OMe

30

H--Ile--Asn--Pro--NH--CH--CO--Tyr--Arg--Leu--Arg--Tyr--OMe,

(SEQ ID NO: 314)

H-[Tyr-Arg-Leu-Arg-Tyr]₂ -Ome

or a pharmaceutically acceptable salt thereof.

35

PYY and PYY agonists may be produced by recombinant DNA technology by chemical synthesis, from natural sources, or by any combination thereof. PYY and PYY agonists may be modified by well known processes such as amidation,

-84-

glycosylation, acylation (e.g. acetylation), sulfation, phosphorylation, cyclization, lipidization and pegylation. Methods for lipidization with fatty acid derivatives of sulfhydryl-containing compounds are disclosed in U.S. Patent No. 5,936,092; U.S. Patent No. 6,093,692; and U.S. Patent No. 6,225,445. Fatty acid derivatives of

5 sulfhydryl-containing PYY and PYY agonists comprising fatty acid-conjugated products with a disulfide linkage are employed for delivery of the PYY and PYY agonists to neuronal cells and tissues. This modification markedly increases the absorption of the compounds relative to the rate of absorption of the unconjugated compounds, as well as prolonging blood and tissue retention of the compounds.

10 Moreover, the disulfide linkage in the conjugate is quite labile in the cells and thus facilitates intracellular release of the intact compounds from the fatty acid moieties.


Fatty acids, as constituents of phospholipids, make up the bulk of cell membranes. Due to their lipidic nature, fatty acids can easily partition into and interact with the cell membrane in a non-toxic way. Therefore, fatty acids represent

15 potentially a useful carrier ligand for the delivery of proteins and peptides. Strategies that may use fatty acids in the delivery of proteins and peptides include the covalent modification of proteins and peptides and the use of fatty acid emulsions.

To prepare such conjugates, a sulfhydryl-containing PYY and PYY agonist

20 is attached to a fatty acid derivative via a reversible, biodegradable disulfide bond. Such a conjugate is expected to bind to the apical side of a cell membrane, reach the basolateral membrane of the GI-epithelium as a result of membrane transport and turnover, and become released into interstitial fluid as the result of disulfide bond reduction.

25 Such lipidized PYY and PYY agonist compounds have the general formula



in which P is a residue derived from a PYY or PYY agonist; R¹ is hydrogen, lower alkyl or aryl; R² is a lipid-containing moiety and R³ is --OH, a lipid-containing

30 moiety or an amino acid chain comprising one or 2 amino acids and terminating in --


CO₂H or -COR². See U.S. Patent No. 5,936,092. These conjugates are particularly useful for increasing the absorption and prolonging blood and tissue retention of PYY and PYY agonists.

Typical alkyl groups include C₁₋₆ alkyl groups including methyl, ethyl, 5 propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, 3-pentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, and the like.

Preferred aryl groups are C₆₋₁₄ aryl groups and typically include phenyl, naphthyl, fluorenlyl, phenanthryl, and anthracetyl groups.

10 The term "lipid-containing moiety" refers to either a lipid group per se or a hydrocarbon-based group (in particular, one or more amino acids) comprising a lipid group. By the term "lipid group" is meant a hydrophobic substituent consisting of 4 to 26 carbon atoms, preferably 5 to 19 carbon atoms. Suitable lipid groups include, but are not limited to, the following: palmityl (C₁₅H₃₁), oleyl (C₁₈H₃₇), stearyl (C₁₈H₃₅), cholate; and deoxycholate.

15 PCT Application No. WO 00/34236 describes drug-carrier conjugates and synthetic strategies for their production, as well as synthetic methods, intermediates, and final products useful for the uptake and release of biologically-active amino group containing compounds. Such lipidized PYY and PYY agonist compounds 20 have general Formula I



in which R² is selected from the group consisting of hydrogen, halo, alkyl, or aryl, wherein the alkyl or aryl groups are optionally substituted with one or more alkoxy, alkoxyalkyl, alkanoyl, nitro, cycloalkyl, alkenyl, alkynyl, alkanoyloxy, alkyl or 25 halogen atoms;

R³ is a lipophilic group; one of R⁴ and R⁵ is a PYY or a PYY agonist and the other of R⁴ and R⁵ is OR⁶ where R⁶ is hydrogen, an alkali metal or a negative charge;

X is oxygen or sulfur;

Y is a bridging natural or unnatural amino acid; n is zero or 1; and m is an integer from zero to 10.

Typical alkyl groups include C₁₋₆ alkyl groups including methyl, ethyl, 5 propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, 3-pentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, and the like.

Typical alkoxy groups include oxygen substituted by any of the alkyl groups mentioned above.

10 Typical alkoxyalkyl groups include any of the above alkyl groups substituted by an alkoxy group, such as methoxymethyl, ethoxymethyl, propoxymethyl, butoxymethyl, pentoxymethyl, hexoxymethyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, methoxyhexyl, and the like.

15 Preferred aryl groups are C₆₋₁₄ aryl groups and typically include phenyl, naphthyl, fluorenlyl, phenanthryl, and anthracyl groups.

Typical alkoxy substituted aryl groups include the above aryl groups substituted by one or more of the above alkoxy groups, e.g., 3-methoxyphenyl, 2-ethoxyphenyl, and the like.

20 Typical alkyl substituted aryl groups include any of the above aryl groups substituted by any of the C₁₋₆ alkyl groups, including the group Ph(CH₂)_n, where n is 1-6, for example, toyl, o-, m-, and p-xylyl, ethylphenyl, 1-propylphenyl, 2-propylphenyl, 1-butylphenyl, 2-butylphenyl, t-butylphenyl, 1-pentylphenyl, 2-pentylphenyl, 3-pentylphenyl.

25 Typical alkenyl groups include C₂₋₆ alkenyl groups, e.g. ethenyl, 2-propenyl, isopropenyl, 2-butenyl, 3-butenyl, 4-pentenyl, 3-pentenyl, 2-pentenyl, 5-hexenyl, 4-hexenyl, 3-hexenyl, and 2-hexenyl groups.

Typical alkynyl groups include C₂₋₆ alkynyl groups e.g. enthynyl, 2-propenyl, 2-butynyl, 3-butynyl, 4-pentynyl, 3-pentynyl, 2-pentynyl, 5-hexynyl, 4hexynyl, 3-hexynyl, and 2-hexynyl groups.

30 Typical alkenyl or alkynyl substituted aryl groups include any of the above C₆₋₁₄ aryl groups substituted by any of the above C₂₋₆ alkenyl or C₂₋₆ alkynyl groups, e.g., ethenylphenyl, 1-propenylphenyl, 2-propenylphenyl, 1butenylphenyl, 2-

butenylphenyl, 1-pentenylphenyl, 2-pentenylphenyl, 3-pentenylphenyl, 1-hexenylphenyl, 2-hexenylphenyl, 3-hexenylphenyl, ethynylphenyl, 1-propynylphenyl, 2-propynylphenyl, 1-butynylphenyl, 2-butynylphenyl, 1-pentynylphenyl, 2-pentynylphenyl, 3-pentynylphenyl, 1-hexynylphenyl, 2-hexynylphenyl, 3-hexynylphenyl groups.

Typical halo groups include fluorine, chlorine, bromine, and iodine.

Typical halo substituted alkyl groups include C₁₋₆ alkyl groups substituted by one or more fluorine, chlorine, bromine, or iodine atoms, e.g., fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, 1,1-difluoroethyl, and

10 trichloromethyl groups.

Typical alkanoyl groups include C₁₋₅C(=O)– alkanoyl groups, e.g., acetyl, propionyl, butanoyl, pentanoyl, and hexanoyl groups, or by an arylalkanoyl group, e.g., a C₁₋₅C(=O) – alkanoyl group substituted by any of the above aryl groups.

15 Typical cycloalkyl groups include C₃₋₈ cycloalkyl groups including cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl groups.

20 The term "lipophilic group" as used herein refers to either a naturally occurring lipid per se, a hydrophobic branched or unbranched hydrocarbon comprising about 4 to about 26 carbon atoms, preferably about 5 to about 19 carbon atoms, a fatty acid or ester thereof, or a surfactant. Suitable lipophilic groups include, but are not limited to, long chain alkanoyl groups including: palmityl (C₁₅H₃₁), oleyl (C₁₅H₂₉), stearyl (C₁₇H₃₅), lauryl (C₁₁H₂₃), cholyl, and myristyl (C₁₃H₂₇)

25 The term "natural or unnatural amino acid" as used herein refers to any of the 21 naturally occurring amino acids as well as D-form amino acids, blocked L-and D-form amino acids such as those blocked by amidation or acylation, substituted amino acids (e.g., those substituted with a sterically hindered alkyl group or a cycloalkyl group such as cyclopropyl or cyclobutyl) in which the substitution introduces a conformational restraint in the amino acid. The preferred naturally occurring amino acids for use in the present disclosure as amino acids or components of a peptide or protein are alanine, arginine, asparagine, aspartic acid, citrulline, cysteine, cystine, γ -glutamic acid, glutamine, glycine, histidine, isoleucine, norleucine, leucine, lysine,

methionine, ornithine, phenylalanine, proline, hydroxyproline, serine, threonine, tryptophan, tyrosine, valine, γ -carboxyglutamate, or O-phosphoserine. The preferred non-naturally occurring amino acids for use in the present disclosure as amino acids or components of peptides or proteins are any of the β -amino acids, e.g.,

5 α -alanine, γ -amino butyric acid, γ -amino butyric acid, γ (aminophenyl)butyric acid, α -amino isobutyric acid, ϵ -amino caproic acid, 7-amino heptanoic acid, amino benzoic acid, aminophenyl acetic acid, aminophenyl butyric acid, cysteine (ACM), methionine sulfone, phenylglycine, norvaline, ornithine, δ -ornithine, α nitro-phenylalanine, 1,2,3,4-terahydroisoquinoline-3-carboxylic acid and thioproline.

10 The present disclosure is also directed to methods of preparing lipidized conjugates of PYY and PYY agonists, pharmaceutical compositions comprising lipidized conjugates of PYY and PYY agonists, and methods of increasing the delivery of amino group-containing PYY and PYY agonists into a cell.

15 Also provided by the disclosure are chemically modified derivatives of PYY and PYY agonists which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent No. 4,179,337). Such modified derivatives include PYY and PYY agonists modified by pegylation. The terms "pegylated" and "pegylation" refer to the process of reacting a poly(alkylene glycol), preferably an

20 activated poly(alkylene glycol), with a facilitator such as an amino acid, e.g. lysine, to form a covalent bond. Although "pegylation" is often carried out using poly(ethylene glycol) or derivatives thereof, such as methoxy poly(ethylene glycol), the term is not intended to be so limited here, but is intended to include any other useful poly(alkylene glycol), such as, for example poly(propylene glycol).

25 The chemical moieties for derivitization may also be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or

30 more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between

about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release

5 desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog). For example, the polyethylene glycol may have an average molecular weight of about 200, 500, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10,000,

10 10,500, 11,000, 11,500, 12,000, 12,500, 13,000, 13,500, 14,000, 14,500, 15,000, 15,500, 16,000, 16,500, 17,000, 17,500, 18,000, 18,500, 19,000, 19,500, 20,000, 25,000, 30,000, 35,000, 40,000, 50,000, 55,000, 60,000, 65,000, 70,000, 75,000, 80,000, 85,000, 90,000, 95,000, or 100,000 kDa.

As noted above, the polyethylene glycol may have a branched structure.

15 Branched polyethylene glycols are described, for example, in U.S. Patent No. 5,643,575; Morpurgo et al., *Appl. Biochem. Biotechnol.* 56:59-72, 1996; Vorobjev et al., *Nucleosides Nucleotides* 18:2745-2750, 1999; and Caliceti et al., *Bioconjug. Chem.* 10:638-646, 1999.

20 The polyethylene glycol molecules (or other chemical moieties) should be attached to the polypeptides or proteins with consideration of effects on functional or antigenic domains of the polypeptides or proteins. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384 (coupling PEG to G-CSF), see also Malik et al., *Exp. Hematol.* 20:1028-1035, 1992 (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol

25 may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid

30 residues and the C-terminal amino acid residue. Sulphydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for

therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

As suggested above, polyethylene glycol may be attached to proteins and polypeptides via linkage to any of a number of amino acid residues. For example,

5 polyethylene glycol can be linked to proteins and polypeptides via covalent bonds to lysine, histidine, aspartic acid, glutamic acid, or cysteine residues. One or more reaction chemistries may be employed to attach polyethylene glycol to specific amino acid residues (e.g., lysine, histidine, aspartic acid, glutamic acid, or cysteine) of the polypeptide or protein or to more than one type of amino acid residue (e.g.,

10 lysine, histidine, aspartic acid, glutamic acid, cysteine and combinations thereof) of the protein or polypeptide.

One may specifically desire proteins and polypeptides chemically modified at the N-terminus. Using polyethylene glycol as an illustration, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.),

15 the proportion of polyethylene glycol molecules to protein (or peptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally

20 pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective

25 derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

As indicated above, pegylation of the proteins and polypeptides may be accomplished by any number of means. For example, polyethylene glycol may be attached to the protein or polypeptide either directly or by an intervening linker.

30 Linkerless systems for attaching polyethylene glycol to proteins and polypeptides are described in Delgado et al., *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-304,

1992; Francis et al., *Intern. J. of Hematol.* 68:1-18, 1998; U.S. Patent No. 4,002,531; U.S. Patent No. 5,349,052; WO 95/06058; and WO 98/32466.

One system for attaching polyethylene glycol directly to amino acid residues of proteins and polypeptides without an intervening linker employs tresylated

5 MPEG, which is produced by the modification of monmethoxy polyethylene glycol (MPEG) using tresylchloride ($\text{ClSO}_2\text{CH}_2\text{CF}_3$). Upon reaction of the protein or polypeptide with tresylated MPEG, polyethylene glycol is directly attached to amine groups of the protein or polypeptide. Thus, the disclosure includes protein-polyethylene glycol conjugates produced by reacting proteins and polypeptides with

10 a polyethylene glycol molecule having a 2,2,2-trifluoroethane sulphonyl group.

Polyethylene glycol can also be attached to proteins and polypeptides using a number of different intervening linkers. For example, U.S. Patent No. 5,612,460 discloses urethane linkers for connecting polyethylene glycol to proteins. Protein-polyethylene glycol conjugates wherein the polyethylene glycol is attached to the 15 protein or polypeptide by a linker can also be produced by reaction of proteins or polypeptides with compounds such as MPEG-succinimidylsuccinate, MPEG activated with 1,1'-carbonyldiimidazole, MPEG-2,4,5-trichloropenylcarbonate, MPEG- ρ -nitrophenolcarbonate, and various MPEG-succinate derivatives. A number of additional polyethylene glycol derivatives and reaction chemistries for 20 attaching polyethylene glycol to proteins and polypeptides are described in WO 98/32466.

The number of polyethylene glycol moieties attached to each protein or polypeptide (i.e., the degree of substitution) may also vary. For example, the pegylated proteins and polypeptides may be linked, on average, to 1, 2, 3, 4, 5, 6, 7, 25 8, 9, 10, 12, 15, 17, 20, or more polyethylene glycol molecules. Similarly, the average degree of substitution within ranges such as 1-3, 2-4, 3-5, 4-6, 5-7, 6-8, 7-9, 8-10, 9-11, 10-12, 11-13, 12-14, 13-15, 14-16, 15-17, 16-18, 17-19, or 18-20 polyethylene glycol moieties per protein or polypeptide molecule. Methods for determining the degree of substitution are discussed, for example, in Delgado et al., 30 *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-304, 1992.

The proteins and polypeptides containing substantially non-antigenic polymers, preferably poly(alkylene glycols) may be prepared, for example, as

described in U.S. Patent No. 5,428,128; U.S. Patent No. 6,127,355; and U.S. Patent No. 5,880,131.

To effect covalent attachment of poly(ethylene glycol) (PEG) to a protein or polypeptide, the hydroxyl end groups of the PEG must first be converted into reactive functional groups. This process is frequently referred to as "activation" and the product is called "activated PEG." Methoxy poly(ethylene glycol) (mPEG), distally capped with a reactive functional group is often used. One such activated PEG is succinimidyl succinate derivative of PEG (SS-PEG). See also Abuchowski et al., *Cancer Biochem. Biophys.* 7:175-186, 1984; and U.S. Patent No. 5,122,614 which discloses poly(ethylene glycol)-N-succinimide carbonate and its preparation.

Alternative substantially non-antigenic polymers that may be employed in the practice of the present disclosure include materials such as dextran, polyvinyl pyrrolidones, polysaccharides, starches, polyvinyl alcohols, polyacrylamides, or other similar non-immunogenic polymers. Those of ordinary skill in the art will realize that the foregoing are merely illustrative and not intended to restrict the type of polymeric substances suitable for use herein.

In one aspect of the disclosure, the polymer is introduced into the peptide or protein molecule after being functionalized or activated for reaction and attachment to one or more amino acids. By activation, it is understood by those of ordinary skill in the art that the polymer is functionalized to include a desired reactive group. See, for example, U.S. Patent No. 4,179,337 and U.S. Patent No. 5,122,614. In this embodiment, the hydroxyl end groups of poly(alkylene glycols) are converted and activated into reactive functional groups.


In another aspect of the disclosure, the polymer is conjugated to a facilitator moiety prior to being introduced into the polypeptide or protein molecule. The facilitator moiety is preferably an amino acid such as lysine, however, non-amino acid moieties are also contemplated. Within the aspect, there are included multifunctionalized organic moieties such as alkyls or substituted alkyls. Such moieties can be prepared to have a nucleophilic functional group such as an amine and an electrophilic group such as an acid as well as a suitably functionalized region for conjugating with the desired polymer or polymers.

The facilitator moieties allow easier inclusion of a polymer into the peptide

or protein molecule during synthesis. For example, poly(alkylene glycols) coupled to facilitator amino acids or amino acid residues in polypeptides or proteins by means of suitable coupling agents are illustrative. A useful review of a number of coupling agents known in the art appears in Dreborg et al., *Critical Reviews in*

5 *Therapeutic Drug Carrier Systems* 6(4):315-165, 1990, see especially, pp. 317-320.

Pegylated PYY peptides and agonists can also be of the general formula



wherein:

D is a residue of a PYY peptide or agonist;

10 X is an electron withdrawing group;


Y and Y' are independently O or S;

(n) is zero (0) or a positive integer; preferably from 1 to about 12;

R₁ and R₂ are independently selected from the group consisting of H, C₁₋₆ alkyls, aryls, substituted aryls, aralkyls, heteroalkyls, substituted heteroalkyls, and

15 substituted C₁₋₆ alkyls;

R₃ is a substantially non-antigenic polymer, C₁₋₁₂ straight or branched alkyl or substituted alkyl, C₅₋₈ cycloalkyl or substituted cycloalkyl, carboxyalkyl, carboalkoxy alkyl, dialkylaminoalkyl, phenylalkyl, phenylaryl or



20 R₄ and R₅ are independently selected from the group consisting of H, C₁₋₆ alkyls, aryls, substituted aryls, aralkyls, heteroalkyls, substituted heteroalkyls and substituted C₁₋₆ alkyls or jointly form a cyclic C_{5-C} ring. See U.S. Patent No. 6,127,355.

Typical alkyl groups include C₁₋₆ alkyl groups including methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, 3-pentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, and the like.

-94-

Preferred aryl groups are C₆₋₁₄ aryl groups and typically include phenyl, naphthyl, fluorenyl, phenanthryl, and anthracyl groups.

Typical alkyl substituted aryl groups include any of the above aryl groups substituted by any of the C₁₋₆ alkyl groups, including the group Ph(CH₂)_n, where n is 5 1-6, for example, toyl, o-, m-, and p-xylyl, ethylphenyl, 1-propylphenyl, 2-propylphenyl, 1-butylphenyl, 2-butylphenyl, t-butylphenyl, 1-pentylphenyl, 2-pentylphenyl, 3-pentylphenyl.

Typical cycloalkyl groups include C₃₋₈ cycloalkyl groups including cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl 10 groups.

Typical electron withdrawing groups include O, NR₁, S, SO and SO₂, wherein R₁ is defined above.

GLP-1 and Agonists Thereof

15

GLP-1 is produced from preproglucagon, which is a 160 amino acid polypeptide, in the central nervous system (CNS) and the intestine. It is released into the circulation in response to nutrient intake. Physiological actions of GLP-1 in man include stimulation of insulin release, suppression of gastric acid secretion and 20 slowing of gastric emptying.

GLP-1 (1-37) (SEQ ID NO: 336) is the initial product of the processing of preproglucagon. GLP-1 (1-37) is amidated by post-translational processing to yield GLP-1 (1-36) NH (SEQ ID NO 337), or is enzymatically processed to give GLP-1 (7-37) (SEQ ID NO: 338). GLP-1 (7-37) can be amidated to give GLP-1 (7-36) 25 amide (SEQ ID NO: 339). The sequences of human GLP-1 are given below:

GLP-1 (1-37): His Asp Glu Phe Glu Arg His Ala Glu Gly Thr Phe The Ser Asp Val Ser Ser Tyr Leu Glu Gly Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ ID NO: 336).

30

-95-

GLP-1 (1-36) amide: His Asp Glu Phe Glu Arg His Ala Glu Gly Thr Phe The Ser Asp Val Ser Ser Tyr Leu Glu Gly Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg NH₂ (SEQ ID NO: 337),

5 GLP-1 (7-37): His Ala Glu Gly Thr Phe The Ser Asp Val Ser Ser Tyr Leu Glu Gly Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ ID NO: 338).

GLP-1 (7-36) amide: His Ala Glu Gly Thr Phe The Ser Asp Val Ser Ser Tyr Leu Glu Gly Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg NH₂
10 (SEQ ID NO: 339).

A GLP-1 agonist is a peptide, small molecule, or chemical compound that preferentially binds to the GLP-1 receptor and stimulates the same biological activity as does GLP-1. In one embodiment, an agonist for the GLP-1 receptor
15 binds to the receptor with an equal or greater affinity than GLP-1. In another embodiment, an agonist selectively binds the GLP-1 receptor, as compared to binding to another receptor. Exendin-4, which is a 39-amino acid peptide isolated from the salivary glands of the Gila monster (*Heloderma suspectum*) (Eng J et al J Biol Chem 267:7402-7405, 1992) is an example of an agonist at the GLP-1 receptor.
20 Molecules derived from exendin-4 and that also have GLP-1 agonist activity are further examples of GLP-1 agonists. GLP-1 agonists include GLP-1 related peptides and peptides that result from natural or synthetic enzymatic or chemical processing of preproglucagon or of a GLP-1 peptide or a related peptide.

Any compound that is described as being a GLP-1 agonist may be used in
25 the present invention, as may any compound that is tested for GLP-1 agonist activity, for example, as described above, and found to function as a GLP-1 agonist. A recombinant GLP-1 receptor suitable for use in screening is disclosed in WO93/19175. Many GLP-1 agonists are known and are described in the art. Examples of published patent specifications that disclose GLP-1 agonists are the
30 following: WO2002/67918, WO2002/66479, WO2002/03978, WO2001/89554, WO2001/14386, WO2001/66135, WO2001/35988, WO2001/14368, WO2001/04156, WO2000/78333, WO2000/59887, WO2000/42026, EP 0955314,

and WO99/43707. Examples of GLP-1 agonists are Arg34, Lys26(N-epsilon-(gamma-Glu(N-alpha-hexadecanoyl))-GLP-1 (7-37), IP7-GLP-1 (7-37)OH.

GLP-1 or an agonist thereof may be administered according to the present invention peripherally at a dose of, for example, 0.1 nmoles or more per kg body weight of the subject, for example, 0.2 nmoles or more, for example, 0.4 nmoles or more, for example, 0.6 nmoles or more, for example, 0.8 nmoles or more, for example, 1.0 nmole or more, for example, 1.2 nmoles or more, for example, 1.4 nmoles or more, for example, 1.6 nmoles or more, for example, 1.8 nmoles or more, for example, 2.0 nmoles or more, for example, 2.2 nmoles or more, for example, 2.4 nmoles or more, for example, 2.6 nmoles or more, for example, 2.8 nmoles, for example, 3.0 nmoles or more, for example, up to 3.2 nmoles per kg body weight. The amount used may be up to 3.0 nmoles per kg body weight, for example, up to 2.8 nmoles, for example, up to 2.6 nmoles, for example, up to 2.4 nmoles, for example, up to 2.2 nmoles, for example, up to 2.0 nmoles, for example, up to 1.8 nmoles, for example, up to 1.4 nmoles, for example, up to 1.2 nmoles, for example, up to 1.0 nmoles, for example, up to 0.8 nmoles, for example, up to 0.6 nmoles, for example, up to 0.4 nmoles, for example, up to 0.2 nmoles per kg body weight. The dose is generally in the range of from 0.1 to 3.2 nmoles per kg body weight, for example, within any combination of upper and lower ranges given above.

We have shown that intracerebroventricular (ICV) administration of GLP-1 to rats potently inhibits food intake. This effect is blocked by the specific GLP-1 receptor antagonist, exendin 9-39. GLP-1 receptors are found in the brainstem, arcuate nucleus and PVN. Following ICV GLP-1 administration, c-fos is expressed in the arcuate nucleus and PVN. However, peripheral administration of GLP-1 in man and rat also inhibits food intake and results in c-fos expression in the brainstem of rats. As is disclosed herein, the effect on 1-hour food intake in non-fasted rats of IP administration of GLP-1 is not influenced by the presence of concomitant exendin 9-39, a GLP-1 receptor agonist, in the arcuate nucleus of the rat. This indicates that circulating GLP-1 acts via the brainstem rather than the arcuate nucleus.

These and other findings suggest that the main site for appetite inhibition by peripheral GLP-1 is the dorsal vagal complex, acting in part directly through the

area postrema. However, the physiological importance of this in man is currently unclear.

As disclosed herein, when administered to humans, PYY was found to reduce appetite. When infused into humans at physiological post-prandial levels, 5 PYY₃₋₃₆ significantly decreased appetite and reduced food intake by a third over 12 hours, and even by a third over 24 hours. Both the effect itself and the duration of the effect are surprising and unpredictable, as they occurred for many hours after the hormone had been cleared from the circulation. The effects, which are produced at physiological levels of the peptide, are strong indications that PYY acts in vivo to 10 regulate feeding behavior.

As disclosed herein, peripheral administration of PYY₃₋₃₆ in the rat caused an increase of c-fos immunoreactivity in the arcuate nucleus of the hypothalamus and a decrease in hypothalamic neuropeptide Y (NPY) mRNA. Further, electro-physiological studies demonstrated that PYY₃₋₃₆ inhibits synaptic activity of the 15 NPY nerve terminals and thus activates POMC neurons, which are known to receive inhibitory NPY synaptic inputs.

Without being bound by theory, these results demonstrate that the gut hormone PYY₃₋₃₆ can act via the neuropeptide Y Y2 receptor. This hypothesis is supported by the observation that when PYY₃₋₃₆ was administered to neuropeptide 20 Y Y2 receptor null mice (Y2R gene knock out mice), no inhibition of feeding was observed. Administration of PYY₃₋₃₆ to wild type littermates of the Y2R null mice was fully effective in inhibiting feeding.

Thus, a novel gut-brain pathway that inhibits feeding after meals is described. Without being bound by theory, the natural pathway involves release of 25 PYY from the gut, its conversion to PYY₃₋₃₆, which acts as an agonist on the neuropeptide Y Y2 receptor (NPY Y2 receptor) in the brain. The NPY Y2 receptor acts as a inhibitory pre-synaptic receptor reducing release of neuropeptide Y, which is a most potent stimulator of feeding, and also acting on the anorexigenic melanocortin systems, the result of the NPY Y2 receptor activity being to suppress 30 appetite and decrease food intake. The action of PYY₃₋₃₆ may occur in the arcuate nucleus of the hypothalamus, but other areas may be also be involved.

-98-

The results obtained show that PYY₃₋₃₆, a gut hormone that circulates in the blood, inhibits appetite at physiological concentrations, and that the inhibitory effect is observed even for some hours after the hormone has been cleared from the blood. This effect has been observed in all species tested, i.e. in mouse, rat and human. The 5 circulating gut hormone appears to act via hypothalamic circuits. The reduction of messenger RNA, necessary for the synthesis of brain appetite regulating hormones, in particular of hypothalamic NPY mRNA may be a possible mechanism for the long action of PYY₃₋₃₆.

When PYY₃₋₃₆ was co-administered parenterally, either IP or IV with GLP-1, 10 a clear synergistic reduction in food intake was produced. This contrasts with the results obtained with oxyntomodulin, where there was no synergistic effect, only a sum of the individual effects. As indicated above, PYY₃₋₃₆ is considered to act via the arcuate nucleus. Oxyntomodulin is also considered to act via that region of the brain. GLP-1, however, is believed to act via the brainstem, as explained above. 15 While not being limited to the following, it appears that, when agents that act via different areas of the brain and/or by different neurological routes are administered parenterally, they produce a synergistic effect, whereas when they act by the same area and/or route they do not.

These results indicate the importance *in vivo* of the interaction of gut 20 hormones on the regulation of appetite and food intake. The present invention demonstrates that, surprisingly, it is possible to regulate appetite and food intake more effectively by informed choice of agents.

The disclosure is illustrated by the following non-limiting Examples.

25

EXAMPLES

Example 1

Material and Methods

30 *Generation of POMC-EGFP mice:* The *EGFP* cassette contains its own Kozak consensus translation initiation site along with *SV40* polyadenylation signals downstream of the *EGFP* coding sequences directing proper processing of the 3' end

-99-

of the EGFP mRNA. The *EGFP* cassette was introduced by standard techniques into the 5' untranslated region of exon 2 of a mouse *Pomc* genomic clone containing 13 kb of 5' and 2 kb of 3' flanking sequences (Young et al., *J Neurosci* 18, 6631-40, 1998). The transgene was microinjected into pronuclei of one-cell stage embryos of 5 C57BL/6J mice (Jackson Laboratories) as described (Young et al., *J Neurosci* 18, 6631-40, 1998). One founder was generated and bred to wildtype C57BL/6J to produce N₁ hemizygous mice. In addition, N₂ and subsequent generations of mice homozygous for the transgene were also generated. The mice are fertile and have normal growth and development.

10 *Immunofluorescence and GFP co-localization:* Anesthetized mice were perfused transcardially with 4% paraformaldehyde and free-floating brain sections prepared with a vibratome. Sections were processed for immunofluorescence and colocalization of GFP fluorescence using standard techniques. Primary antisera and their final dilutions were rabbit anti- β -endorphin, 1:2500 v/v; rabbit anti-NPY, 15 1:25,000 v/v (Alanex Corp.); rabbit anti-ACTH, 1:2000 v/v; and mouse anti-TH, 1:1000 v/v (Incstar). After rinsing, sections were incubated with 10mg/ml biotinylated horse anti-mouse/rabbit IgG (Vector Laboratories) followed by Cy-3 conjugated streptavidin, 1:500 v/v (Jackson Immunoresearch Laboratories). Photomicrographs were taken on a Zeiss Axioscop using FITC and RITC filter sets 20 (Chroma Technology Corp.).

25 *Electrophysiology (Example 2):* 200 μ m thick coronal slices were cut from the ARC of four-week old male POMC-EGFP mice. Slices were maintained in (in mM) [NaCl, 126; KCl, 2.5; MgCl₂, 1.2; CaCl₂.2H₂O, 2.4; NaH₂PO₄.H₂O, 1.2; NaHCO₃, 21.4; Glucose, 11.1] (Krebs) at 35°C and saturated with 95% O₂ 5% CO₂ for 1 hour(hr) prior to recordings. Recordings were made in Krebs at 35° C. Slices were visualized on an Axioskop FS2 (Zeiss) through standard infra red optics and using epifluorescence through a FITC filter set (see Fig. 1c). Whole cell recordings were made from fluorescent neurons using an Axopatch 1D amplifier (Axon 30 Instruments) and Clampex 7 (Axon Instruments). Resting membrane potentials were determined using an event detection protocol on a PowerLab system (AD Instruments, Mountain View, CA) to average expanded traces of the membrane

-100-

potential. Drugs were applied to the bath over the times indicated. The resting membrane potential was stable for up to an hour in cells treated with Krebs alone. I-V relationships for the Met-Enk currents were established using a step protocol; (-60 mV holding potential, sequentially pulsed (40 ms) from -120 to -50 mV, cells 5 were returned to -60 mV for 2 s between voltage steps). The protocol was repeated after Met Enk addition. The net current was the difference between the two I-V relationships. This protocol was repeated in Krebs with 6.5 mM K⁺. I-V relationships to identify the postsynaptic leptin current were performed similarly with slow voltage ramps (5 mV/s from -100 to -20 mV) before and 10 minutes 10 after the addition of leptin (100 nM). GABAergic IPSCs were recorded using a CsCl internal electrode solution (in mM) [CsCl, 140; Hepes, 10; MgCl₂, 5; Bapta, 1; (Mg)-ATP, 5; (Na)GTP, 0.3]. Both mini IPSCs and large amplitude (presumably multisynaptic) IPSCs were observed in the untreated slices. TTX (1 μM) abolished large IPSCs. Data were acquired before and after addition of drug for the times 15 indicated on the figures at a -50 mV holding potential in 2 s. sweeps every 4 s. Mini postsynaptic currents were analyzed using Axograph 4 (Axon Instruments). IPSCs and excitatory postsynaptic currents (EPSCs) were distinguished on the basis of their decay constants; additionally picrotoxin (100 μM) blocked all IPSCs. POMC neurons receive a low EPSC tone and the frequency was not modulated by any of 20 the treatments described here.

Immunostaining for light and electron microscopy: Double immunocytochemistry for NPY and POMC using different color diaminobenzidine(DAB) chromogens was carried out on fixed mouse hypothalami 25 according to published protocols (Horvath et al., *Neuroscience* 51, 391-9, 1992). For electron microscopy, preembedding immunostaining for β-endorphin was using an ABC Elite kit (Vector Laboratories) and a DAB reaction followed by post-embedding labeling of GABA and NPY using rabbit anti-GABA, 1:1000 v/v and gold conjugated (10 nm) goat anti-rabbit IgG or sheep anti-NPY and gold 30 conjugated (25 nm) goat anti-sheep IgG. Finally, sections were contrasted with saturated uranyl acetate (10 minutes) and lead citrate (20-30 s) and examined using a Philips CM-10 electron microscope.

-101-

Animals: Male Wistar rats (200-250g), 7-8 weeks old (Charles River Laboratories, United Kingdom) were maintained under controlled temperature (21-23° C) and light conditions (lights on 07:00-19:00) with *ad libitum* access to water and food (RM1 diet; SDS Ltd., Witham, United Kingdom) except where stated.

5 Arcuate and paraventricular nuclei cannulations and injections were performed as previously described (Glaum et al., *Mol. Pharmacol.* 50, 230-5, 1996; Lee et al., *J. Physiol (Lond)* 515, 439-52; 1999; Shiraishi et al., *Nutrition* 15, 576-9, 1999). Correct intranuclear cannula placement was confirmed histologically at the end of each study period (Glaum et al., *Mol. Pharmacol* 50, 230-5, 1996; Lee et al., *J. Physiol (Lond)* 515, 439-52, 1999; Shiraishi et al., *Nutrition* 15, 576-9, 1999). All animal procedures were approved under the British Home Office Animals (Scientific Procedures) Act, 1986. All injection studies on fasting animals were performed in the early light-phase (0800-0900). All dark-phase feeding studies injections were performed just prior to lights off.

10

15

Male *Pomc-EGFP* mice were studied at 5-6 weeks of age and were generated as described above. *Y2r*-null mice were generated using *Cre-lox P* mediated recombination, which results in the germline deletion of the entire coding region of the *Y2* receptor. All *Y2r*-null mice were maintained on a mixed C57/B16-129SvJ background. Male mice aged 8-12 weeks and between 20-30 g bodyweight were kept under controlled temperature (21-23° C) and light conditions (lights on 06:00-18:00) with *ad libitum* access to water and food (Gordon's Speciality Stock feeds) except where stated. All studies were performed in the early light-phase (0700-0800).

20

25

Intraperitoneal injections: Rats were accustomed to IP injection by injections of 0.5 ml saline on the two days prior to study. For all studies, animals received an IP injection of PYY₃₋₃₆, GLP-1 or saline in 500 µl (for rats) or 100 µl (for mice).

30

Electrophysiology: Whole cell patch clamp recordings were made from POMC neurons in the hypothalamus of 180 μ m thick coronal slices from *Pomc*-EGFP mice, as previously reported (Cowley et al., *Nature* 411, 480-484, 2001). “Loose cell-attached” recordings were made using extracellular buffer in the 5 electrode solution, and maintaining seal resistance between 3-5Mohm throughout the recording. Firing rates were analysed using mini-analysis protocols (MiniAnalysis, Jaejin Software, NJ). Vehicle controls were used in this system, previously validated for the electrophysiological actions of neuropeptides (Cowley et al., *Nature* 411, 480-484, 2001). Data were analysed by ANOVA, Neuman-Keuls 10 posthoc comparison, and Wilcoxon Signed Rank Test.

Hypothalamic explants: Male Wistar rats were killed by decapitation and the whole brain immediately removed, mounted with the ventral surface uppermost and placed in a vibrating microtome (Biorad, Microfield Scientific Ltd., Devon, UK). A 15 1.7 mm slice was taken from the base of the brain to include the PVN and the ARC and immediately transferred to 1ml of artificial CSF (aCSF) (Kim et al., *J. Clin. Invest.* 105, 1005-11, 2000) equilibrated with 95% O₂ and 5% CO₂ and maintained at 37° C. After an initial 2-hour equilibration period, with aCSF replaced every 60 minutes, the hypothalami were then incubated for 45 minutes in 600 μ l aCSF (basal 20 period) before being exposed to the Y2A (50nM) in 600 μ l aCSF. Finally, the viability of the tissue was verified by a 45 minute exposure to 56 mM KCL; isotonicity was maintained by substituting K⁺ for Na⁺. At the end of each period, the aCSF was removed and frozen at -20° C until assayed for NPY and α MSH by radioimmunoassay.

25 *C-fos expression:* C-fos expression was measured in adult Wistar rats and *Pomc*-EGFP mice 2 hours after IP administration of saline or PYY₃₋₃₆ (5 μ g/100g) using standard immunohistochemical techniques (Hoffman et al., *Front. Neuroendocrinol.* 14, 173-213, 1993). Data were obtained from 3 rats and 5 mice in 30 each group. For the *Pomc*-EGFP mice 5 anatomically matched arcuate nucleus sections (Franklin et al., *The Mouse Brain in Stereotaxic Coordinates*, Academic

-103-

Press, San Diego, 1997) were counted from each animal, and images acquired using a Leica TSC confocal microscope (Grove et al., *Neuroscience* 100, 731-40, 2000).

RNase protection assay (RPA): Total RNA was extracted from hypothalamus (Trizol, Gibco). RPAs were performed (RPAIII kit, Ambion) using 5 µg RNA and probes specific for NPY, αMSH and β actin (internal standard). For each neuropeptide, the ratio of the optical density of the neuropeptide mRNA band to that of β actin was calculated. Neuropeptide mRNA expression levels are expressed relative to saline control (mean ± s.e.m. n = 4 per group). The statistical analysis used was ANOVA, with Bonferroni post hoc analysis.

Plasma assays: Human leptin was measured using a commercially available radioimmunoassay (RIA) (Linco Research, USA). All other plasma hormone levels were measured using established in-house RIAs (Tarling et al., *Intensive Care Med.* 23, 256-260, 1997). Glucose concentrations were measured using a YSI 2300STAT analyser (Yellow Springs Instruments Inc., Ohio, USA). Plasma paracetamol levels were measured using an enzymatic colorimetric assay (Olympus AU600 analyzer).

Human Studies: PYY₃₋₃₆ was purchased from Bachem (California, USA). The Limulus Amoebocyte Lysate assay test for pyrogen was negative and the peptide was sterile on culture. Ethical approval was obtained from the Local Research Ethics Committee (project registration 2001/6094) and the study was performed in accordance with the principles of the Declaration of Helsinki. Subjects gave informed written consent.

Each subject was studied on two occasions with at least 1 week between each study. Volunteers filled out a food diary for three days prior to each infusion, and for the following 24 hours. All subjects fasted and drank only water from 20:00 on the evening prior to each study. Subjects arrived at 08:30 on each study day, were cannulated and then allowed to relax for 30 minutes prior to the onset of the study protocol. Blood samples were collected every 30 minutes into heparinised tubes containing 5,000 Kallikrein Inhibitor Units (0.2 ml) of aprotinin (Bayer) and centrifuged. Plasma was separated and then stored at -70° C until analysis. Subjects

-104-

were infused with either saline or $0.8 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ PYY₃₋₃₆ for 90 minutes (about 72 pmol total infusion), in a double blind randomized crossover design.

Two hours after the termination of the infusion, subjects were offered an excess free-choice buffet meal (Edwards et al., *Am. J. Physiol. Endocrinol. Metab.* 281, E155-E166, 2001), such that all appetites could be satisfied. Food and water were weighed pre- and postprandially and caloric intake calculated. Appetite ratings were made on 100 mm visual analogue scores (VAS) with the text expressing the most positive and the negative rating anchored at each end (Raben et al., *Br. J. Nutr.* 73, 517-30, 1995). VAS was used to assess hunger, satiety, fullness, prospective food consumption and nausea. Caloric intake following saline and PYY₃₋₃₆ were compared using a paired t test. The postprandial response curves were compared by ANOVA using repeated paired measures, with time and treatment as factors.

Measurements of Energy Expenditure: To determine the actions of PYY on energy expenditure the OXYMAX system is utilized with rodents following PYY injection into a treatment cohort. This system is also utilized with rodents following a saline injection (control cohort). The equipment measures O₂ consumption and CO₂ production; the efficiency with which the body produces CO₂ from O₂ gives a reliable index of caloric or metabolic efficiency. A similar system is used with human volunteers.

-105-

Example 2

Neural Network in the Arcuate Nucleus

A strain of transgenic mice was generated expressing green fluorescent protein (EGFP Clontech), under the transcriptional control of mouse *Pomc* genomic sequences that include a region located between -13 kb and -2 kb required for accurate neuronal expression (Young et al., *J Neurosci* 18, 6631-40, 1998) (Fig. 1a). Bright green fluorescence (509 nm) was seen in the two CNS regions where POMC is produced: the ARC and the nucleus of the solitary tract. Under ultraviolet (450–480 nm) excitation POMC neurons were clearly distinguished from adjacent, non-fluorescent neurons (Fig. 1b) visualized under infrared optics. Double immunofluorescence revealed >99% cellular co-localization of EGFP and POMC peptides within the ARC (Fig. 1c). There was close apposition of both tyrosine hydroxylase (TH)- and NPY-stained terminals on EGFP-expressing POMC neurons, but no evidence of co-localization of the TH or NPY immunoreactivity with EGFP.

15 Total fluorescent cell counts performed on coronal hypothalamic sections revealed 3148 ± 62 (mean \pm SEM: n=3) POMC-EGFP neurons distributed through the entire ARC (Franklin et al., *The Mouse Brain in Stereotaxic Coordinates*, Academic Press, San Diego, 1997) (Fig. 1d). POMC neurons in the mouse are located both medially and ventrally within the ARC, in contrast to a predominantly lateral position in the

20 rat ARC.

POMC-EGFP neurons in hypothalamic slices had a resting membrane potential of -40 to -45 mV and exhibited frequent spontaneous action potentials. The non-selective opioid agonist met-enkephalin (Met-Enk: 30 μ M; Sigma) caused a rapid (35- 40 s), reversible hyperpolarization (10-20 mV) of the membrane potential of POMC cells (n=10) and prevented spontaneous action potential generation (Fig. 2a). In normal (2.5 mM K⁺) Krebs buffer, the reversal-potential of the inwardly-rectifying opioid current was approximately -90mV, while in 6.5 mM K⁺ Krebs the reversal-potential was shifted to approximately -60 mV (n=3: Fig. 2b). The μ opioid receptor (MOP-R) antagonist CTAP (1 μ M; Phoenix Pharmaceuticals) 25 completely prevented the current induced by Met-Enk in POMC cells (n=3: Fig. 2c). These characteristics indicate the opioid current was due to activation of MOP-R and increased ion conductance through G protein coupled, inwardly-rectifying

potassium channels (GIRK) (Kelly et al., *Neuroendocrinology* 52, 268-75, 1990). The similar opioid responses in EGFP-labeled POMC neurons to that of a guinea pig (Kelly et al., *Neuroendocrinology* 52, 268-75, 1990) or mouse (Slugg et al., *Neuroendocrinology* 72, 208-17, 2000). POMC cells, identified by post-recording immunohistochemistry, suggests that expression of the EGFP transgene does not compromise either expression of receptors nor their coupling to second messenger systems in POMC neurons.

Next, the direct effects of leptin on identified POMC cells in slice preparations were investigated. Leptin (0.1 – 100 nM) depolarized 72 of 77 POMC cells by 3-30 mV (Fig. 3a; mean \pm SEM depolarization at 100 nM leptin = 9.7 \pm 1.2 mV, n= 45) within 2-10 minutes, in a concentration responsive manner (Fig. 3b). There were two components to the depolarization and neither were fully reversible within 40 minutes. Firstly, the depolarization was due to a small inward current which reversed at approximately -20 mV (Fig. 3c), suggesting the involvement of a non-specific cation channel (Powis et al., *Am J Physiol* 274, R1468-72, 1998). Secondly, leptin treatment decreased the GABAergic tone onto POMC cells. GABAergic inhibitory postsynaptic currents (IPSCs) were observed in POMC cells and leptin (100 nM) decreased their frequency by 25% (Fig. 3d) in 5 out of 15 cells suggesting that it acted presynaptically to reduce GABA release (leptin had no effect on IPSCs in 10 out of 15 POMC neurons). The effect on IPSC frequency occurred with a similar lag to the effect on membrane potential. Thus, leptin not only directly depolarizes POMC neurons but also acts at GABAergic nerve terminals to reduce the release of GABA onto POMC neurons, allowing them to adopt a more depolarized resting potential. The consistent depolarization of POMC cells by leptin was specific because leptin had no effect on 5 of 13 adjacent non-fluorescent cells tested (Fig. 3e), while it hyperpolarized 5 (Fig. 3f) and depolarized 3 other non-POMC neurons in the ARC. The electrophysiological effects of leptin reported here are consistent with leptin's biological actions; leptin rapidly causes release of α -MSH from rat hypothalami (Kim et al., *J Clin Invest* 105, 1005-11, 2000), presumably by activating POMC neurons.

Previous reports of neuronal hyperpolarization by leptin (Glaum et al., *Mol Pharmacol* 50, 230-5, 1996; Spanswick et al., *Nature* 390, 521-5, 1997), and the

demonstrated co-localization of GABA and NPY (Horvath et al., *Brain Res* 756, 283-6, 1997) within subpopulations of ARC neurons, led us to speculate that leptin hyperpolarizes NPY/GABA cells that directly innervate POMC neurons, and thus reduces GABAergic drive onto POMC cells. Both the leptin and NPY Y2 receptors 5 are expressed on NPY neurons in the ARC (Hakansson et al., *J Neurosci* 18, 559-72, 1998; Broberger et al., *Neuroendocrinology* 66, 393-408, 1997). Furthermore, activation of Y2 receptors inhibits NPY release from NPY neurons (King et al., *J Neurochem* 73, 641-6, 1999), and presumably would also diminish GABA release 10 from NPY/GABA terminals. This is an alternative pharmacological approach, independent of leptin, to test the hypothesized innervation of POMC neurons by GABAergic NPY neurons. Indeed, NPY (100 nM; Bachem) decreased the frequency of GABAergic IPSCs by 55% within 3 minutes, in all 12 POMC cells tested (Fig. 4a). Both NPY and leptin still inhibited IPSCs in the presence of 15 tetrodotoxin (TTX) (6 of 6 and 3 of 5 cells respectively), indicating that some of the inhibition of IPSCs was occurring through direct effects at presynaptic nerve terminals. POMC neurons express the NPY Y1 receptor (Broberger et al., *Neuroendocrinology* 66, 393-408, 1997) and NPY also hyperpolarized all POMC 20 neurons tested, by an average of 9±6 mV (n=3).

Another pharmacological test to confirm the origin of GABAergic 25 innervation on POMC neurons from NPY/GABA terminals was to test the effect of the recently characterized and highly selective MC3-R agonist D-Trp⁸- γ MSH (Grieco et al., *J Med Chem* 43, 4998-5002, 2000) on local GABA release. D-Trp⁸- γ MSH (7 nM) increased the frequency of GABAergic IPSCs (280 ± 90%) recorded from 3 of 4 POMC neurons (Fig. 4b). It had no effect on one cell. The positive 30 effect of MC3-R activation, together with the negative effects of NPY and leptin, demonstrate the dynamic range of the NPY/GABA synapse onto POMC neurons and point to the important role of this synapse in modulating signal flow within the ARC. D-Trp⁸- γ MSH (7 nM) also hyperpolarized (-5.5 ± 2.4 mV) 9 of 15 POMC neurons tested and decreased the frequency of action potentials (Fig 4c); the remaining cells showed no significant response to D-Trp⁸- γ MSH. These effects could be due entirely to increased GABA release onto the POMC cells, or could be due to an additional postsynaptic action of D-Trp⁸- γ MSH on POMC neurons,

-108-

approximately half of which also express the MC3-R (Bagnol et al., *J Neurosci (Online)* 19, RC26, 1999). Thus, MC3-R acts in a similar autoreceptor manner to MOP-Rs on POMC neurons, diminishing POMC neuronal activity in response to elevated POMC peptides.

5 To further determine that the IPSCs in POMC neurons were due to local innervation by NPY/GABA cells, multi-label immunohistochemistry was performed using light and electron microscopy. Although independent NPY (Csiffary et al., *Brain Res* 506, 215-22, 1990) and GABA (Horvath et al., *Neuroscience* 51, 391-9, 1992) innervation of POMC cells has been reported, co-localization of NPY and 10 GABA in nerve terminals forming synapses onto POMC cells has not been shown. Similar to the rat (Csiffary et al., *Brain Res* 506, 215-22, 1990), a dense innervation of POMC cells by NPY axon terminals was detected in the mouse (Fig. 4d). Electron microscopy confirmed the coexpression of NPY and GABA in axon 15 terminals and revealed that these boutons established synapses on the perikarya of all 15 ARC POMC neurons analyzed (representative example, Fig. 4e).

A detailed model of regulation of this circuit shows dual mechanisms of 20 leptin action in the ARC, interactions between NPY/GABA and POMC neurons, and autoregulatory feedback from opioid and melanocortin peptides as well as NPY (Fig. 4f). In this model, leptin directly depolarizes the POMC neurons and simultaneously hyperpolarizes the somata of NPY/GABA neurons, and diminishes 25 release from NPY/GABA terminals. This diminished GABA release disinhibits the POMC neurons, and result in an activation of POMC neurons and an increased frequency of action potentials.

25

Example 3

Administration of PYY Inhibits Food Intake

The orexigenic NPY and the anorectic alpha melanocortin stimulating 30 hormone (α -MSH) systems of the hypothalamic arcuate nucleus are involved in the central regulation of appetite (Schwartz et al., *Nature* 404, 661-671, 2000). However the potential mechanisms signaling meal ingestion directly to these hypothalamic-feeding circuits are unclear. PYY₃₋₃₆ is a gut-derived hormone that is released postprandially in proportion to the calories ingested (Pedersen-Bjergaard et

-109-

al., *Scand. J. Clin. Lab. Invest.* 56, 497-503, 1996). The effects of peripheral administration of PYY₃₋₃₆ on feeding were investigated.

An intraperitoneal injection (IP) of PYY₃₋₃₆ to freely feeding rats, prior to the onset of the dark-phase, significantly decreased subsequent food intake (Fig. 5a). A 5 similar inhibition of feeding was seen following IP injection in rats fasted for 24 hours (Fig. 5b). A time course of the plasma PYY₃₋₃₆ levels achieved following IP injection of PYY₃₋₃₆ demonstrated a peak level at 15 minutes post injection, which was within the normal postprandial range (peak PYY₃₋₃₆ levels 15 minutes post IP injection of 0.3 μ g/100g = 99.3 ± 10.4 pmol/l vs. peak postprandial level = 112.1 ± 10 7.8 pmol/l, n = 8-10 per group), suggesting that physiological concentrations of PYY₃₋₃₆ inhibit feeding. PYY₃₋₃₆ did not affect gastric emptying (percentage of food ingested remaining in the stomach at 3 hours: PYY₃₋₃₆ = 36 ± 1.9 %, saline = 37.4 ± 1.0 % n = 12) (Barrachina et al., *Am. J. Physiol.* 272, R1007-11, 1997). PYY₃₋₃₆ administered IP twice daily for 7 days reduced cumulative food intake (7-day 15 cumulative food intake: PYY₃₋₃₆ = 187.6 ± 2.7g vs. saline = 206.8 ± 2.3, n = 8 per group, P < 0.0001) and decreased body weight gain (Fig. 5d) (PYY₃₋₃₆ = 48.2 ± 1.3g vs. saline = 58.7 ± 1.9, n = 8 per group, P < 0.002).

Example 4

20 PYY Administration Affects c-fos Expression

To investigate whether this inhibition of food intake involved a hypothalamic pathway, c-fos expression was examined in the arcuate nucleus, an important center of feeding control (Schwartz et al., *Nature* 404, 661-671, 2000; Cowley et al., *Nature* 411, 480-484, 2001), following a single IP injection of PYY₃₋₃₆. There was a 25 2-fold increase in the number of cells positive for c-fos in the lateral arcuate of the rat (PYY₃₋₃₆ = 168 ± 2, saline = 82.7 ± 5, n = 3, P < 0.0001). Likewise in *Pomc-EGFP*-transgenic mice (Cowley et al., *Nature* 411, 480-484, 2001) IP administration of PYY₃₋₃₆ resulted in a 1.8-fold increase in the number of arcuate cells positive for c-fos (Fig. 6b), compared with saline control animals (Fig. 6a) (PYY₃₋₃₆ = 250 ± 40, 30 saline = 137 ± 15, n = 5, P < 0.05). IP PYY₃₋₃₆ caused a 2.6 fold increase in the proportion of POMC neurons that express c-fos (PYY₃₋₃₆ = 20.4 ± 2.9%, saline = 8 ± 1.4%, n = 5, P < 0.006) (Figs. 6c and d).

-110-

These observations suggested that PYY₃₋₃₆ may act via the arcuate nucleus. Thus, the actions of PYY₃₋₃₆, and its effects upon NPY and POMC circuits in the hypothalamus, were studied. In view of the sustained inhibition of food intake and the effects on weight gain following peripheral administration of PYY₃₋₃₆ both *Pomc* and *Npy* hypothalamic messenger RNA (mRNA) were measured using RNase protection assays. A significant decrease in *Npy* mRNA in response to PYY₃₋₃₆ was observed 6 hours post IP injection, compared with saline treated animals (saline = 17.3 ± 2.0, PYY₃₋₃₆ = 8.8 ± 1.0, relative optical density units, P < 0.02). A non-significant increase occurred in *Pomc* mRNA levels.

10

Example 5

Y2 receptors

PYY₃₋₃₆ shows a 70% amino acid sequence identity to NPY and acts through NPY receptors (Soderberg et al., *J. Neurochem.* 75, 908-18, 2000). The Y2R is a putative inhibitory presynaptic receptor and is highly expressed on the arcuate NPY neurons (Broberger et al., *Neuroendocrinology* 66, 393-408, 1997), though not on the neighboring POMC neurons. PYY₃₋₃₆ is a high affinity agonist at the Y2 receptor (Grandt et al., *Regul. Pept.* 51, 151-159, 1994). It was hypothesized that peripheral PYY₃₋₃₆ inhibits food intake via the Y2R in the arcuate nucleus, an area known to be directly accessible to circulating hormones (Kalra et al., *Endocr. Rev.* 20, 68-100, 1999).

To investigate this hypothesis, PYY₃₋₃₆ was injected directly into the arcuate nucleus (Kim et al., *Diabetes* 49, 177-82, 2000). In rats fasted for 24 hours, food intake was significantly decreased by doses as low as 100 fmol (Fig. 7a), resulting in a similar inhibition to that seen following IP administration. To establish whether these effects were via the Y2R, aY2R selective agonist was used (Potter et al., *Eur. J. Pharmacol.* 267, 253-262, 1994), N-acetyl (Leu²⁸, Leu³¹) NPY (24-36) [Y2A]. Its affinity was confirmed using receptor-binding studies (Small et al., *Proc. Natl. Acad. Sci. U.S.A.* 94, 11686-91, 1997) on cell lines expressing the NPY Y1, Y2 and Y5 receptors (Y2 IC₅₀ = 1.3 ± 0.2 nM, Y1 IC₅₀ > 5000 nM, Y5 IC₅₀ > 5000 nM). Intra-arcuate nucleus injection of Y2A in rats previously fasted for 24 hours dose-dependently (100 fmol – 1 nmol) inhibited food intake (chow ingested 2 hours post-

-111-

injection, 0.1 nmol Y2A = 6.2 ± 0.5 g, saline = 8.2 ± 0.6 g, n = 8 per group, P < 0.05).

To confirm the anatomical specificity of this effect Y2A (100 fmol - 1 nmol) was injected into the paraventricular nucleus (PVN) (Kim et al., *J. Clin. Invest.* 105, 1005-11, 2000) of rats fasted for 24 hours and found no alteration of food intake (2 hour post-injection saline = 8.3 ± 0.4 g, 0.1nmol Y2A = 8.0 ± 0.6 g, n = 8 per group). To further determine the role of the Y2R in the feeding inhibition caused by peripheral PYY₃₋₃₆, the effect of PYY₃₋₃₆ on Y2r-null mice and littermate controls was examined. PYY₃₋₃₆ inhibited daytime feeding in a dose responsive manner in fasted male wild-type mice but did not inhibit food intake in fasted male Y2r-null mice (Figs. 7b and 7c). Food intake measured in response to a fast demonstrated that male Y2r-null mice eat significantly more at 2, 4 and 24 hours compared with their littermate controls (24-hour cumulative food intake; Y2r-null mice = 7.1 ± 0.48 g vs. wild-type = 5.3 ± 0.7 g, n = 8 per group, P < 0.05).

The electrophysiological response of hypothalamic POMC neurons to administration of both PYY₃₋₃₆ and Y2A was examined. These neurons were identified using mice with targeted expression of green fluorescent protein in POMC neurons (Cowley et al., *Nature* 411, 480-484, 2001). PYY₃₋₃₆ disinhibited the POMC neurons, resulting in a significant depolarization of 19 of the 22 POMC neurons tested (Fig. 8a inset) (10.3 ± 2.1 mV depolarization, n = 22, P < 0.0003). A similar depolarization was seen with Y2A (8.7 ± 1.8 mV depolarization, n = 9, P < 0.002). The depolarization caused by PYY₃₋₃₆ stimulated a significant increase in the frequency of action potentials in POMC neurons (Fig 8a) (93% increase over control, P < 0.05, n = 22). In the whole cell mode the effect of PYY₃₋₃₆ was sometimes reversed upon washout, but only after a long latency (30 minutes). A similar washout of leptin effects upon these neurons was observed.

To exclude effects of cellular rundown, or seal deterioration, the effects of PYY₃₋₃₆ in the “loose cell-attached” (or extracellular) configuration was examined. PYY₃₋₃₆ caused a reversible 5-fold increase in the frequency of action potentials in loose cell-attached recordings of POMC neurons (Fig. 8b). This increase in firing rate occurred with the same latency as PYY₃₋₃₆ reduced the frequency of inhibitory postsynaptic currents (IPSCs) onto all 13 POMC neurons tested (Fig. 8c) (51.9 ± 9.2

% reduction, n = 13, P < 0.0001), indicating a reduced frequency of GABA release onto POMC neurons. Interestingly, the firing rate of POMC neurons returned to basal, in spite of continued inhibition of IPSCs. A similar effect upon IPSC frequency was seen with Y2A (44.4 ± 9.3% reduction, n = 8, P < 0.004) suggesting this effect to be via Y2R. PYY₃₋₃₆ (25 nM) caused a hyperpolarization (5.2 ± 1.16 mV, P < 0.004, n = 5) of unidentified, but presumably NPY-containing, non-POMC, neurons in the arcuate nucleus. There is a tonic GABAergic inhibition of POMC neurons by NPY neurons (Cowley et al., *Nature* 411, 480-484, 2001) and these results suggest that PYY₃₋₃₆ acts by inhibiting NPY neurons, thus decreasing this GABAergic tone and consequentially disinhibiting POMC neurons. The effect of Y2A on peptide secretion was also examined using hypothalamic explants (Kim et al., *J. Clin. Invest.* 105, 1005-11, 2000). Y2A significantly decreased NPY release, with a concomitant increase in α-MSH release from hypothalamic explants (Figs. 8d and 4e). Taken together, these observations suggest that PYY₃₋₃₆ modulates both the NPY and melanocortin systems in the arcuate nucleus.

Example 6

Human Studies

Because of the importance of the melanocortin system in man (Barsh et al., *Nature* 404, 644-651, 2000) and the profound effects of PYY₃₋₃₆ on both feeding and weight change seen in rodents, the effects of PYY₃₋₃₆ on appetite and food intake were investigated in human subjects. Twelve healthy fasted, non-obese volunteers (six men and six women, mean age 26.7 ± 0.7 years, BMI = 24.6 ± 0.94 kg.m⁻²) were infused with PYY₃₋₃₆ (0.8 pmol.kg⁻¹.min⁻¹) or saline for 90 minutes in a double-blind placebo controlled crossover study.

PYY₃₋₃₆ plasma concentrations increased from mean basal concentration of 8.3 ± 1.0 pM to 43.5 ± 3 pM during the PYY₃₋₃₆ infusion and mimicked postprandial levels (Pedersen-Bjergaard et al., *Scand. J. Clin. Lab. Invest.* 56, 497-503, 1996; Adrian et al., *Gastroenterology* 89, 1070-1077, 1985). Post-infusion, PYY₃₋₃₆ concentrations returned to basal within 30 minutes. PYY₃₋₃₆ infusion resulted in a significant decrease in hunger scores (Raben et al., *Br. J. Nutr.* 73, 517-30, 1995) (Fig. 9c), but not in the scores for sleepiness or sickness. Calorie intake during a

-113-

free-choice buffet meal (Tarling et al., *Intensive Care Med.* 23, 256-260, 1997) two hours after the termination of the infusion was reduced by over a third compared to saline ($36 \pm 7.4\%$, $p < 0.0001$) (Fig. 9a). There was no effect upon fluid intake and no difference in sensations of fullness or nausea reported by the volunteers. PYY₃₋₃₆ 5 administration had no effect on gastric emptying, as estimated by the paracetamol absorption method (Edwards et al., *Am. J. Physiol. Endocrinol. Metab.* 281, E155-E166, 2001; Tarling et al., *Intensive Care Med.* 23, 256-260, 1997), or on plasma glucose, plasma leptin, GLP-1, or insulin. Analysis of the food diaries revealed a significant inhibition of food intake in the 12-hour period following the PYY₃₋₃₆ 10 infusion (saline = 2205 ± 243 kcal, PYY₃₋₃₆ = 1474 ± 207 kcal). However, food intake during a 12 to 24 hour period between the two groups was virtually identical. Overall there was a 33% decrease in cumulative total calorie consumption in the 24-hour period following the PYY₃₋₃₆ infusion (Fig. 9b). These findings demonstrate 15 that infusion of PYY₃₋₃₆, matching postprandial levels, caused a marked inhibition of both appetite and food intake in man.

In an additional study, two groups of healthy subjects ($n = 12$ per group, 6 males and 6 females), one with increased Body Mass Index (BMI) (mean = 32.73 ± 0.93 kg/m²) and another group with low BMI (mean = 20.49 ± 2.05 kg/m²), were studied on two occasions with at least 1 week between each study. All subjects 20 fasted and drank only water from 20:00 hours on the evening prior to each study. Subjects arrived at 08:30 on each study day, were cannulated and then allowed to relax for 30 minutes prior to the onset of the study protocol. Subjects were infused with either saline or $0.8 \text{ pmol.kg}^{-1}.\text{min}^{-1}$ PYY₃₋₃₆ for 90 minutes, in a double blind randomized crossover design. Two hours after the termination of the infusion, 25 subjects were offered an excess free-choice buffet meal, such that all appetites could be satisfied. Food and water were weighed pre- and postprandially and caloric intake calculated. Caloric intake following saline and PYY₃₋₃₆ were compared using a paired t test ($p < 0.001$). The number of calories ingested following administration of PYY₃₋₃₆ differed significantly from the number of calories ingested following 30 administration of saline for both the overweight group and the lean group. The overweight group showed a $28.8 \pm 4.3\%$ reduction and the lean group a $31.1 \pm 4.4\%$ reduction. However, the reduction for the overweight group did not differ

significantly from the reduction for the lean group. These findings demonstrate that infusion of PYY₃₋₃₆, matching postprandial levels, caused a marked inhibition of both appetite and food intake in both lean and overweight subjects.

Administration of PYY₃₋₃₆ at doses of 10 nmoles and 20 nmoles (in saline) 5 subcutaneously to human subjects resulted in raised plasma levels of PYY₃₋₃₆ as shown in Fig. 10.

Without being bound by theory, cells within the arcuate nucleus could detect circulating peripheral satiety signals and relay these signals to other brain regions (Butler et al., *Nature Neuroscience* 4, 605-611, 2001). This is supported by the 10 observation that leptin modifies the activity of both the POMC and NPY arcuate neurons (Cowley et al., *Nature* 411, 480-484, 2001). The results disclosed herein demonstrate, through a combination of electrophysiological and hypothalamic explant studies, that the gut hormone, PYY₃₋₃₆, can directly influence hypothalamic circuits, resulting in coordinate changes in POMC and NPY action. The results 15 presented here demonstrate that NPY neurons in the ARC are not protected by the blood/brain barrier, and thus are accessible to circulating molecules. Furthermore, PYY₃₋₃₆ administered directly into this brain region reduces food intake.

The data disclosed herein demonstrates that postprandial levels of PYY₃₋₃₆ 20 inhibit food intake in more than one mammalian species (e.g. rodents and human subjects) for up to 12 hours, thereby demonstrating a role in regulation of food intake. This role can be described as a long term role, such as over a period of several hours (e.g. at least two, three, four, eight, or twelve hours, or from about two to about fifteen hours). This is in contrast to previously characterized gut-derived 'short-term' satiety signals, e.g. cholecystokinin (Schwartz et al., *Nature* 404, 661- 25 671, 2000; Moran, *Nutrition* 16, 858- 865, 2000), the effects of which are relatively short-lived (e.g., from about 1-4 hours).

The failure of PYY₃₋₃₆ to inhibit food intake in the *Y2r*-null mice provides 30 evidence that PYY₃₋₃₆ reduces food intake via a Y2R dependent mechanism. The results disclosed herein suggest the existence of a novel gut-hypothalamic pathway in the regulation of feeding, involving postprandial PYY₃₋₃₆ acting at the arcuate Y2R. Thus, PYY, and analogs thereof, such as PYY₃₋₃₆ provide novel therapeutic agents for the treatment of obesity.

Example 7**Parenterally administered GLP-1**

5 GLP-1 i.e. GLP-1 (7-36) amide (25 nmol/kg) or saline (control) was administered parenterally to non-fasted rats prior to the onset of the dark phase. Exendin 9-39 (20 nmoles/kg) or saline (control) injected into the arcuate nucleus of non-fasted rats. Food intake one hour later was measured.

10 The results as shown in Fig. 11 demonstrate that the effect on 1-hour food intake in non-fasted rats of IP administration of GLP-1 is not influenced by the presence of concomitant exendin 9-39, a GLP-1 receptor agonist, in the arcuate nucleus of the rat. This indicates that circulating GLP-1 acts via the brainstem rather than the arcuate nucleus. These and other findings suggest that the main site for appetite inhibition by peripheral GLP-1 is the dorsal vagal complex, acting in part 15 directly through the area postrema.

Example 8**Co-administration of PYY₃₋₃₆ and GLP-1 in rats**

20 PYY₃₋₃₆ and GLP-1 i.e. GLP-1 (7-36) amide were injected (intraperitoneally) into non-fasted rats prior to the onset of the dark phase. GLP-1 and PYY₃₋₃₆ were administered separately at the following doses: GLP-1 30 μ g/kg = 1xG or 60 μ g/kg = 2xG in 500 μ l saline; PYY₃₋₃₆ (3 μ g/kg = 1xP or 6 μ g/kg = 2xP in 500 μ l saline). PYY₃₋₃₆ and GLP-1 were co-administrated IP (P+G group) at doses of GLP-1 25 30 μ g/kg and PYY₃₋₃₆ 3 μ g/kg combined.

30 The results set out in Fig. 12 show that there is a clear synergistic effect on the reduction of food intake when PYY₃₋₃₆ and GLP-1 were co-administered (P+G). However, when the procedure was carried out substituting oxyntomodulin for GLP-1, the results obtained for reduction of food intake when PYY₃₋₃₆ and

-116-

oxynotomdulin were co-administered were only the sum of the individual effects. No synergistic effect was observed.

Example 9

5 Co-administration of PYY₃₋₃₆ and GLP-1 in humans

10 healthy volunteers (4 men and 6 women) having a mean age of 24.7 (+/- 0.8) years and a mean BMI of 22.7 (+/- 0.6) kg/m² were recruited. The volunteers were asked to fast and to drink only water from 9 pm on the night preceding all 10 study days. Each volunteer attended for four study days. The volunteers arrived at 9.30 am and the venflons were inserted. A rest period from 10.00 to 10.30 followed. Intravenous infusion started at 10.30, and a buffet lunch was provided between 12.00 and 12.30. Infusion was terminated at 12.30, and the volunteers went home at 13.30.

The volunteers were given the following intra-venous infusions for 120 minutes (90 minutes pre-meal) in random order: Saline, GLP-1, PYY₃₋₃₆, PYY₃₋₃₆+GLP-1. The dose of both GLP-1 and PYY3-36 was 0.4pmol/kg/min. The study was double blinded. GLP-1 is i.e. GLP-1 (7-36) amide

The results of the food intake from the free buffet meal are given in the Table below

20 Food intake from free buffet meal (Kcal)

	Saline	G	P	P+G
mean	997	950	848	724
Sem	68.0	81.3	105.2	66.6
p v saline		0.336318995	0.069375886	3.00605E-06
p v P+G		0.003622523	0.044418337	
% reduction from saline		5%	15%	27%

25 using paired t tests for all statistical comparisons
These result demonstrate a synergistic effect when PYY3-36 and GLP-1 are
co-administered

The results obtained in Examples 4 and 5 above demonstrate that PYY₃₋₃₆ acts via the arcuate nucleus. Oxynotmodulin is also considered to act via that region

-117-

of the brain. The results obtained in Example 8 above indicates that GLP-1 acts via the brainstem. It appears from the results obtained in this Example that, when agents such as PYY₃₋₃₆ and GLP-1 that act via different areas of the brain and/or by different neurological routes are administered parenterally, they produce a

5 synergistic affect, whereas when they act by the same area and/or route such as PYY₃₋₃₆ and oxyntmodulin, they do not. The synergistic effect on food intake obtained when PYY and GLP-1 were co-administered to humans confirms the observation made in rats.

10 These results indicate the importance in vivo of the interaction of gut hormones on the regulation of appetite and food intake. The present invention demonstrates that, surprisingly, it is possible to regulate appetite and food intake more effectively by informed choice of agents.

15 It will be apparent that the precise details of the methods or compositions described may be varied or modified without departing from the spirit of the described disclosure. We claim all such modifications and variations that fall within the scope and spirit of the claims below.

CLAIMS

1. A method for decreasing calorie intake in a subject, a method for decreasing appetite in a subject, a method for decreasing food intake in a subject, a method for 5 weight control or treatment in a subject, or a method for reduction or prevention of obesity in a subject, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a therapeutically effective amount of GLP-1 or an agonist thereof to the subject.
- 10 2. A method for preventing and reducing weight gain in a subject; a method for inducing and promoting weight loss in a subject; or a method for reducing obesity in a subject as measured by the Body Mass Index, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a therapeutically effective amount of GLP-1 or an agonist thereof to the subject.
- 15 3. A method for controlling of any one or more of appetite, satiety and hunger in a subject, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a therapeutically effective amount of GLP-1 or an agonist thereof to the subject.
- 20 4. A method as claimed in claim 3 for inducing, increasing, enhancing or promoting satiety and/or sensations of satiety in a subject, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a therapeutically effective amount of GLP-1 or an agonist thereof to the 25 subject.
5. A method as claimed in claim 3 for reducing, inhibiting or suppressing hunger or sensations of hunger in a subject, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a 30 therapeutically effective amount of GLP-1 or an agonist thereof to the subject.

-119-

6. A method for maintaining desired body weight, a desired Body Mass Index, and/or a desired appearance and good health in a subject, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a therapeutically effective amount of GLP-1 or an agonist thereof to the 5 subject.
7. A method for improving lipid profile in a subject, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a therapeutically effective amount of GLP-1 or an agonist thereof to the 10 subject.
8. A method for alleviating a condition or disorder which can be alleviated by reducing nutrient availability, which comprises peripherally administering a therapeutically effective amount of PYY or an agonist thereof and a therapeutically 15 effective amount of GLP-1 or an agonist thereof to the subject.
9. A method as claimed in any one of claims 1 to 8, wherein the PYY or agonist thereof and the GLP-1 or agonist thereof are administered simultaneously or sequentially.

20

10. A method as claimed in any one of claims 1 to 9, wherein the PYY and the GLP-1 are administered via different routes.
11. A method as claimed in any one of claims 1 to 10, wherein the subject is 25 overweight.
12. A method as claimed in any one of claims 1 to 11, wherein the subject is obese.
- 30 13. A method as claimed in any one of claims 1 to 12, wherein the subject is diabetic.

-120-

14. A method as claimed in any one of claims 1 to 13, wherein peripheral administration comprises subcutaneous, intravenous, intramuscular, intranasal, transdermal or sublingual administration.
- 5 15. A method as claimed in any one of claims 1 to 14, wherein peripherally administering PYY or the agonist thereof comprises administering about 45 to about 135 pmol per kilogram body weight of the subject.
- 10 16. A method as claimed in claim 15, wherein peripherally administering PYY or the agonist thereof comprises administering about 72 pmol per kilogram body weight of the subject.
- 15 17. A method as claimed in claim 15, wherein peripherally administering PYY or the agonist thereof comprises administering about 45 to about 135 pmol per kilogram body weight of the subject at least 30 minutes prior to a meal.
- 20 18. A method as claimed in any one of claims 1 to 14, wherein peripherally administering the therapeutically effective amount of PYY or the agonist thereof comprises administering PYY or an agonist thereof to the subject in a multitude of doses, wherein each dose in the multitude of doses comprises administration of about 0.5 to about 135 pmol per kilogram of body weight at least about 30 minutes prior to a meal.
- 25 19. A method as claimed in any one of claims 1 to 18, wherein the PYY or the agonist thereof is administered in an amount sufficient to decrease calorie intake for a period of at least about 2 hours.
- 30 20. A method as claimed in any claim 19, wherein the PYY or the agonist thereof is administered in an amount sufficient to decrease calorie intake for a period of about 2 to 12 hours.
21. A method as claimed in any one of claims 1 to 14, wherein the PYY or

-121-

agonist thereof is administered peripherally at a dose of 0.1 nmoles per kg body weight of the subject or more, for example, 0.2 nmoles or more, for example, 0.4 nmoles or more, for example, 0.6 nmoles or more, for example, 0.8 nmoles or more, for example, 1.0 nmole or more, for example, 1.2 nmoles or more, for example, 1.4 nmoles or more, for example, 1.6 nmoles or more, for example, 1.8 nmoles or more, for example, 2.0 nmoles or more, for example, 2.2 nmoles or more, for example, 2.4 nmoles or more, for example, 2.6 nmoles or more, for example, 2.8 nmoles, for example, 3.0 nmoles or more, for example, up to 3.2 nmoles per kg body weight.

10 22. A method as claimed in any one of claims 1 to 14 or claim 21, wherein the PYY or agonist thereof and/or the GLP-1 or agonist thereof is administered peripherally in an amount of up to 3.0 nmoles per kg body weight, for example, up to 2.8 nmoles, for example, up to 2.6 nmoles, for example, up to 2.4 nmoles, for example, up to 2.2 nmoles, for example, up to 2.0 nmoles, for example, up to 1.8 nmoles, for example, up to 1.4 nmoles, for example, up to 1.2 nmoles, for example, up to 1.0 nmoles, for example, up to 0.8 nmoles, for example, up to 0.6 nmoles, for example, up to 0.4 nmoles, for example, up to 0.2 nmoles per kg body weight.

15 23. A method as claimed in any one of claims 1 to 14, 21 and 22, wherein the GLP-1 or agonist thereof is administered peripherally at a dose of 0.1 nmoles per kg body weight of the subject or more, for example, 0.2 nmoles or more, for example, 0.4 nmoles or more, for example, 0.6 nmoles or more, for example, 0.8 nmoles or more, for example, 1.0 nmole or more, for example, 1.2 nmoles or more, for example, 1.4 nmoles or more, for example, 1.6 nmoles or more, for example, 1.8 nmoles or more, for example, 2.0 nmoles or more, for example, 2.2 nmoles or more, for example, 2.4 nmoles or more, for example, 2.6 nmoles or more, for example, 2.8 nmoles, for example, 3.0 nmoles or more, for example, up to 3.2 nmoles per kg body weight.

25 24. A method as claimed in any one of claims 1 to 14 and 21 to 23, wherein the GLP-1 or agonist thereof is administered peripherally in an amount of up to 3.0 nmoles per kg body weight, for example, up to 2.8 nmoles, for example, up to 2.6

nmoles, for example, up to 2.4 nmoles, for example, up to 2.2 nmoles, for example, up to 2.0 nmoles, for example, up to 1.8 nmoles, for example, up to 1.4 nmoles, for example, up to 1.2 nmoles, for example, up to 1.0 nmoles, for example, up to 0.8 nmoles, for example, up to 0.6 nmoles, for example, up to 0.4 nmoles, for example, 5 up to 0.2 nmoles per kg body weight.

25. A method as claimed in any one of claims 1 to 24, wherein the PYY agonist comprises a molecule that specifically binds the Y2 receptor.
- 10 26. A method as claimed in claim , wherein the PYY agonist increases the expression of c-fos in a section of an arcuate nucleus contacted with the compound.
27. A method as claimed in any one of claims 1 to 24, wherein the PYY agonist specifically binds to a neuropeptide Y neuron and inhibits an activity of a 15 neuropeptide Y neuron.
28. A method as claimed in claim 27, wherein the PYY agonist decreases the action potential firing rate of the neuropeptide Y neuron.
- 20 29. A method as claimed in claim 27, wherein the neuropeptide Y neuron synapses with a proopiomelanocortin neuron, and wherein binding of the PYY agonist to the neuropeptide Y neuron results in an increased activity of the proopiomelanocortin neuron.
- 25 30. A method as claimed in claim 27, wherein the decreased activity of the neuropeptide Y neuron results in an increase in action potential firing on the proopiomelanocortin neuron.
- 30 31. A method as claimed in any one of claims 1 to 30, wherein the GLP-1 agonist is exendin-4 or a derivative thereof that is a GLP-1 agonist.
32. A method as claimed in any one of claims 1 to 31, further comprising

-123-

administering a therapeutically effective amount of amfepramone (diethylpropion), phentermine, mazindol, phenylpropanolamine, fenfluramine, dextfenfluramine, or fluoxetine

- 5 33. A method as claimed in any one of claims 1 to 32, wherein the subject is human.
34. Use of PYY and an agonist thereof and GLP-1 or an agonist thereof for the manufacture of a medicament or medicaments for use in a method as defined in any 10 one of claims 1 to 33.
35. Use of PYY and an agonist thereof in the manufacture of a medicament for use, in combination with GLP-1 or an agonist thereof, in a method for decreasing calorie intake in a subject, a method for decreasing appetite in a subject, a method 15 for decreasing food intake in a subject, a method for weight control or treatment in a subject, or a method for reduction or prevention of obesity in a subject.
36. Use of PYY and an agonist thereof in the manufacture of a medicament for use, in combination with GLP-1 or an agonist thereof, in a method for preventing 20 and reducing weight gain in a subject; a method for inducing and promoting weight loss in a subject; or a method for reducing obesity in a subject as measured by the Body Mass Index .
37. Use of PYY and an agonist thereof in the manufacture of a medicament for 25 use, in combination with GLP-1 or an agonist thereof, in a method for controlling of any one or more of appetite, satiety and hunger in a subject.
38. Use of PYY and an agonist thereof in the manufacture of a medicament for use, in combination with GLP-1 or an agonist thereof, in a method for inducing, 30 increasing, enhancing or promoting satiety and/or sensations of satiety in a subject.
39. Use of PYY and an agonist thereof in the manufacture of a medicament for

use, in combination with GLP-1 or an agonist thereof, in a method for reducing, inhibiting or suppressing hunger or sensations of hunger in a subject.

40. Use of PYY and an agonist thereof in the manufacture of a medicament for
5 use, in combination with GLP-1 or an agonist thereof, in a method for maintaining
desired body weight, a desired Body Mass Index, and/or a desired appearance and
good health in a subject.

41. Use of PYY and an agonist thereof in the manufacture of a medicament for
10 use, in combination with GLP-1 or an agonist thereof, in a method for improving
lipid profile in a subject

42. Use of PYY and an agonist thereof in the manufacture of a medicament for
use, in combination with GLP-1 or an agonist thereof, in a method for alleviating a
15 condition or disorder which can be alleviated by reducing nutrient availability.

43. Use as claimed in any one of claims 31 to 41, wherein the PYY or agonist
thereof and the GLP-1 or agonist thereof are administered simultaneously, or
sequentially in either order.

20

44. A pharmaceutical composition comprising PYY and an agonist thereof and
GLP-1 or an agonist thereof, in admixture or conjunction with a pharmaceutically
suitable carrier.

25 45. A pharmaceutical composition comprising PYY or an agonist thereof in
admixture with a pharmaceutically suitable carrier, in a form suitable for
subcutaneous administration

46. A pharmaceutical composition as claimed in claim 45, which comprises 10
30 nmoles or more, for example, 20 nmoles or more, for example, 30 nmoles or more,
for example, 40 nmoles or more of PYY or an agonist thereof.

-125-

47. A pharmaceutical composition as claimed in claim 46, which comprises from 20 to 60 nmoles, for example, 35 to 45 nmoles of PYY or an agonist thereof.

48. A pharmaceutical composition as claimed in claim 45, which comprises from 5 PYY or an agonist thereof in an amount suitable for administration peripherally in an amount of up to 3.0 nmoles per kg body weight, for example, up to 2.8 nmoles, for example, up to 2.6 nmoles, for example, up to 2.4 nmoles, for example, up to 2.2 nmoles, for example, up to 2.0 nmoles, for example, up to 1.8 nmoles, for example, up to 1.4 nmoles, for example, up to 1.2 nmoles, for example, up to 1.0 nmoles, for 10 example, up to 0.8 nmoles, for example, up to 0.6 nmoles, for example, up to 0.4 nmoles, for example, up to 0.2 nmoles per kg body weight.

49. A pharmaceutical composition as claimed in claim 45, which comprises from 15 PYY or an agonist thereof in an amount suitable for administration peripherally in an amount of 0.1 nmoles per kg body weight of the subject or more, for example, 0.2 nmoles or more, for example, 0.4 nmoles or more, for example, 0.6 nmoles or more, for example, 0.8 nmoles or more, for example, 1.0 nmole or more, for example, 1.2 nmoles or more, for example, 1.4 nmoles or more, for example, 1.6 nmoles or more, for example, 1.8 nmoles or more, for example, 2.0 nmoles or more, 20 for example, 2.2 nmoles or more, for example, 2.4 nmoles or more, for example, 2.6 nmoles or more, for example, 2.8 nmoles, for example, 3.0 nmoles or more, for example, up to 3.2 nmoles per kg body weight.

50. A pharmaceutical composition as claimed in claim 48 or 49, in unit dosage 25 form.

51. A method as defined in any one of claims 1 to 8, which comprises administering PYY or an agonist thereof subcutaneously at a dose of 10 nmoles or more, for example, 20 nmoles or more, for example, 30 nmoles or more, for 30 example, 40 nmoles or more to the subject of the method.

52. A method as claimed in claim 51, wherein from 20 to 60 nmoles, for

example, 35 to 45 nmoles of PYY or an agonist thereof are administered.

53. Use of PYY or an agonist thereof for the manufacture of a composition for
subcutaneous administration to a subject for treatment of the subject by a method as
5 defined in any one of claims 1 to 8.

54. Use as claimed in claim 53, wherein the treatment comprises administering
PYY or an agonist thereof subcutaneously at a dose of 10 nmoles or more, for
example, 20 nmoles or more, for example, 30 nmoles or more, for example, 40
10 nmoles or more to the subject.

55. Use as claimed in claim 54, wherein the treatment comprises administering
PYY or an agonist thereof subcutaneously at a dose of from 20 to 60 nmoles, for
example, 35 to 45 nmoles to the subject.

15 56. Use as claimed in claim 53, wherein the PYY or agonist thereof is
administered peripherally at a dose of 0.1 nmoles or more per kg body weight of the
subject, for example, 0.2 nmoles or more, for example, 0.4 nmoles or more, for
example, 0.6 nmoles or more, for example, 0.8 nmoles or more, for example, 1.0
20 nmole or more, for example, 1.2 nmoles or more, for example, 1.4 nmoles or more,
for example, 1.6 nmoles or more, for example, 1.8 nmoles or more, for example, 2.0
nmoles or more, for example, 2.2 nmoles or more, for example, 2.4 nmoles or more,
for example, 2.6 nmoles or more, for example, 2.8 nmoles, for example, 3.0 nmoles
or more, for example, up to 3.2 nmoles per kg body weight.

25 57. Use as claimed in claim 53 or claim 56, wherein the PYY or agonist thereof
is administered peripherally in an amount of up to 3.0 nmoles per kg body weight,
for example, up to 2.8 nmoles, for example, up to 2.6 nmoles, for example, up to 2.4
nmoles, for example, up to 2.2 nmoles, for example, up to 2.0 nmoles, for example,
30 up to 1.8 nmoles, for example, up to 1.4 nmoles, for example, up to 1.2 nmoles, for
example, up to 1.0 nmoles, for example, up to 0.8 nmoles, for example, up to 0.6

-127-

nmoles, for example, up to 0.4 nmoles, for example, up to 0.2 nmoles per kg body weight.

1/16




Fig. 1a




Fig. 1b




Fig. 1c

2/16




Fig. 1d

3/16




Fig. 2a




Fig. 2b

4/16




Fig. 2c

5/16




Fig. 3a




Fig. 3b




Fig. 3c




Fig. 3d

6/16




Fig. 3e




Fig. 3f

7/16




Fig. 4a




Fig. 4b




Fig. 4c

8/16

Fig. 4d




Fig. 4e




Fig. 4f

9/16




Fig. 5a




Fig. 5b




Fig. 5c




Fig. 5d

10/16



Fig. 6a





Fig. 6b



● c-Fos ● EGFP colocalisation

Fig. 6c




Fig. 6d

11/16




Fig. 7a

Fig. 7b




Fig. 7c

12/16




Fig. 8a




Fig. 8b

13/16




Fig. 8c




Fig. 8d

Fig. 8e

14/16




Fig. 9a




Fig. 9b




Fig. 9c

15/16




Fig. 10




Fig. 11




Fig. 12

SEQUENCE LISTING

<110> Imperial College Innovations LTD
Oregon Health and Science University

<120> Modification of Feeding Behavior

<130> 9248 WO/Jsvn

<150> GB 02 00507.2
<151> 10-01-2002

<150> PCT/US02/31944
<151> 24-09-2002

<160> 341

<170> PatentIn version 3.1

<210> 1

<211> 36

<212> PRT

<213> Homo sapiens

<400> 1

Tyr Pro Ile Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu
1 5 10 15

Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 2

<211> 36

<212> PRT

<213> Homo sapiens

<400> 2

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15

Met Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 3

<211> 36

<212> PRT

<213> Homo sapiens

<400> 3

Ala Ser Leu Glu Pro Glu Tyr Pro Gly Asp Asn Ala Thr Pro Glu Gln
1 5 10 15

Met Ala Gln Tyr Ala Ala Glu Leu Arg Arg Tyr Ile Asn Met Leu Thr
20 25 30

Arg Pro Arg Tyr

35

<210> 4
<211> 3
<212> PRT
<213> Homo sapiens

<400> 4

000
1

<210> 5
<211> 36
<212> PRT
<213> Rattus sp.

<400> 5

Tyr Pro Ala Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu
1 5 10 15

Leu Ser Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 6
<211> 36
<212> PRT
<213> Sus sp.

<400> 6

Tyr Pro Ala Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu
1 5 10 15

Leu Ser Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 7
<211> 36
<212> PRT
<213> Cavia porcellus

<400> 7

Tyr Pro Ser Lys Pro Glu Ala Pro Gly Ser Asp Ala Ser Pro Glu Glu
1 5 10 15

Leu Ala Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 8
<211> 36
<212> PRT
<213> Rana sp.

<400> 8

Tyr Pro Pro Lys Pro Glu Asn Pro Gly Glu Asp Ala Ser Pro Glu Glu
1 5 10 15

Met Thr Lys Tyr Leu Thr Ala Leu Arg His Tyr Ile Asn Leu Val Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 9

<211> 36

<212> PRT

<213> Raja sp.

<400> 9

Tyr Pro Pro Lys Pro Glu Asn Pro Gly Asp Asp Ala Ala Pro Glu Glu
1 5 10 15

Leu Ala Lys Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 10

<211> 36

<212> PRT

<213> Dogfish sp.

<400> 10

Tyr Pro Pro Lys Pro Glu Asn Pro Gly Glu Asp Ala Pro Pro Glu Glu
1 5 10 15

Leu Ala Lys Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 11

<211> 36

<212> PRT

<213> Lampetra sp.

<400> 11

Phe Pro Pro Lys Pro Asp Asn Pro Gly Asp Asn Ala Ser Pro Glu Gln
1 5 10 15

Met Ala Arg Tyr Lys Ala Ala Val Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 12

<211> 36

<212> PRT

<213> Petromyzontidae gen. sp.

<400> 12

Met Pro Pro Lys Pro Asp Asn Pro Ser Pro Asp Ala Ser Pro Glu Glu
1 5 10 15
Leu Ser Lys Tyr Met Leu Ala Val Arg Asn Tyr Ile Asn Leu Ile Thr
20 25 30
Arg Gln Arg Tyr
35

<210> 13

<211> 36

<212> PRT

<213> Rattus sp.

<400> 13

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15
Met Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30
Arg Gln Arg Tyr
35

<210> 14

<211> 36

<212> PRT

<213> Oryctolagus cuniculus

<400> 14

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15
Met Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30
Arg Gln Arg Tyr
35

<210> 15

<211> 36

<212> PRT

<213> Canis familiaris

<400> 15

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15
Met Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30
Arg Gln Arg Tyr
35

<210> 16

<211> 36

<212> PRT

<213> Sus sp.

<400> 16

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15

Leu Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 17
<211> 36
<212> PRT
<213> Bos taurus

<400> 17

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15

Leu Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 18
<211> 36
<212> PRT
<213> Ovis aries

<400> 18

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Asp Asp Ala Pro Ala Glu Asp
1 5 10 15

Leu Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 19
<211> 36
<212> PRT
<213> Cavia porcellus

<400> 19

Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15

Met Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 20
<211> 36
<212> PRT
<213> Avian

<400> 20

Tyr Pro Ser Lys Pro Asp Ser Pro Gly Glu Asp Ala Pro Ala Glu Asp
Page 5

1 5 10 15
Met Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30
Arg Gln Arg Tyr
35

<210> 21
<211> 36
<212> PRT
<213> Rana sp.

<400> 21
Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15

Met Ala Lys Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 22
<211> 36
<212> PRT
<213> Carassius auratus

<400> 22
Tyr Pro Thr Lys Pro Asp Asn Pro Gly Glu Gly Ala Pro Ala Glu Glu
1 5 10 15

Leu Ala Lys Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 23
<211> 36
<212> PRT
<213> Dogfish sp.

<400> 23
Tyr Pro Ser Lys Pro Asp Asn Pro Gly Glu Gly Ala Pro Ala Glu Asp
1 5 10 15

Leu Ala Lys Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 24
<211> 36
<212> PRT
<213> Lampetra sp.

<400> 24
Pro Pro Asn Lys Pro Asp Ser Pro Gly Glu Asp Ala Pro Ala Glu Asp
1 5 10 15

Leu Ala Arg Tyr Leu Ser Ala Val Arg His Tyr Ile Asn Leu Ile Thr
20 25 30

Arg Gln Arg Tyr
35

<210> 25
<211> 36
<212> PRT
<213> Ovis aries

<400> 25

Ala Pro Leu Glu Pro Val Tyr Pro Gly Asp Asn Ala Thr Pro Glu Gln
1 5 10 15

Met Ala Gln Tyr Ala Ala Asp Leu Arg Arg Tyr Ile Asn Met Leu Thr
20 25 30

Arg Pro Arg Tyr
35

<210> 26
<211> 36
<212> PRT
<213> Sus sp.

<400> 26

Ala Pro Leu Glu Pro Val Tyr Pro Gly Asp Asp Ala Thr Pro Glu Gln
1 5 10 15

Met Ala Gln Tyr Ala Ala Glu Leu Arg Arg Tyr Ile Asn Met Leu Thr
20 25 30

Arg Pro Arg Tyr
35

<210> 27
<211> 36
<212> PRT
<213> Canis familiaris

<400> 27

Ala Pro Leu Glu Pro Val Tyr Pro Gly Asp Asp Ala Thr Pro Glu Gln
1 5 10 15

Met Ala Gln Tyr Ala Ala Glu Leu Arg Arg Tyr Ile Asn Met Leu Thr
20 25 30

Arg Pro Arg Tyr
35

<210> 28
<211> 36
<212> PRT
<213> Felis catus

<400> 28

Ala Pro Leu Glu Pro Val Tyr Pro Gly Asp Asn Ala Thr Pro Glu Gln
1 5 10 15

Met Ala Gln Tyr Ala Ala Glu Leu Arg Arg Tyr Ile Asn Met Leu Thr
20 25 30

Arg Pro Arg Tyr
35

<210> 29
<211> 36
<212> PRT
<213> Bos taurus

<400> 29

Ala Pro Leu Glu Pro Glu Tyr Pro Gly Asp Asn Ala Thr Pro Glu Gln
1 5 10 15

Met Ala Gln Tyr Ala Ala Glu Leu Arg Arg Tyr Ile Asn Met Leu Thr
20 25 30

Arg Pro Arg Tyr
35

<210> 30
<211> 36
<212> PRT
<213> Rattus sp.

<400> 30

Ala Pro Leu Glu Pro Met Tyr Pro Gly Asp Tyr Ala Thr His Glu Gln
1 5 10 15

Arg Ala Gln Tyr Glu Thr Gln Leu Arg Arg Tyr Ile Asn Thr Leu Thr
20 25 30

Arg Pro Arg Tyr
35

<210> 31
<211> 36
<212> PRT
<213> Mus musculus

<400> 31

Ala Pro Leu Glu Pro Met Tyr Pro Gly Asp Tyr Ala Thr Pro Glu Gln
1 5 10 15

Met Ala Gln Tyr Glu Thr Gln Leu Arg Arg Tyr Ile Asn Thr Leu Thr
20 25 30

Arg Pro Arg Tyr
35

<210> 32
<211> 37
<212> PRT
<213> Cavia porcellus

<400> 32

Ala Pro Leu Glu Pro Val Tyr Pro Gly Asp Asn Ala Thr Pro Glu Gln
1 5 10 15

Gln Met Ala Gln Tyr Ala Ala Glu Met Arg Arg Tyr Ile Asn Met Leu
Page 8

20

25

30

Thr Arg Pro Arg Tyr
35

<210> 33
<211> 36
<212> PRT
<213> Gallus gallus
<400> 33

Gly Pro Ser Gln Pro Thr Tyr Pro Gly Asp Asp Ala Pro Val Glu Asp
1 5 10 15

Leu Ile Arg Phe Tyr Asn Asp Leu Gln Gln Tyr Leu Asn Val Val Thr
20 25 30

Arg His Arg Tyr
35

<210> 34
<211> 36
<212> PRT
<213> Alligator sp.

<400> 34

Thr Pro Leu Gln Pro Lys Tyr Pro Gly Asp Gly Ala Pro Val Glu Asp
1 5 10 15

Leu Ile Gln Phe Tyr Asn Asp Leu Gln Gln Tyr Leu Asn Val Val Thr
20 25 30

Arg Pro Arg Phe
35

<210> 35
<211> 36
<212> PRT
<213> Rana catesbeiana

<400> 35

Ala Pro Ser Glu Pro His His Pro Gly Asp Gln Ala Thr Pro Asp Gln
1 5 10 15

Leu Ala Gln Tyr Tyr Ser Asp Leu Tyr Gln Tyr Ile Thr Phe Ile Thr
20 25 30

Arg Pro Arg Phe
35

<210> 36
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 36

Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 37
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 37

Arg His Thr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 38
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 38

Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 39
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 39

Arg His Tyr Ile Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 40
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 40

Arg His Tyr Val Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 41
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 41
Arg His Tyr Leu Gln Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 42
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 42

Arg His Tyr Leu Asn Ile Val Thr Arg Gln Arg Tyr
1 5 10

<210> 43
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 43

Arg His Tyr Leu Asn Val Val Thr Arg Gln Arg Tyr
1 5 10

<210> 44
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 44

Arg His Tyr Leu Asn Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 45
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 45

Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 46
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 46

Arg His Tyr Leu Asn Leu Val Ser Arg Gln Arg Tyr
Page 11

1 5 10

<210> 47
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 47

Arg His Tyr Leu Asn Leu Val Thr Lys Gln Arg Tyr
1 5 10

<210> 48
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 48

Arg His Tyr Leu Asn Leu Val Thr Arg Asn Arg Tyr
1 5 10

<210> 49
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 49

Arg His Tyr Leu Asn Leu Val Thr Arg Gln Lys Tyr
1 5 10

<210> 50
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 50

Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Thr
1 5 10

<210> 51
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 51

Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Phe
1 5 10

<210> 52
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 52

Lys His Thr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 53
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 53

Lys His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 54
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 54

Lys His Tyr Ile Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 55
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 55

Lys His Tyr Val Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 56
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 56

Lys His Tyr Leu Gln Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 57
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation
<400> 57

Lys His Tyr Leu Asn Ile Val Thr Arg Gln Arg Tyr
1 5 10

<210> 58
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation
<400> 58

Lys His Tyr Leu Asn Val Val Thr Arg Gln Arg Tyr
1 5 10

<210> 59
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation
<400> 59

Lys His Tyr Leu Asn Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 60
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation
<400> 60

Lys His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 61
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 61

Lys His Tyr Leu Asn Leu Val Ser Arg Gln Arg Tyr
1 5 10

<210> 62

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 62

Lys His Tyr Leu Asn Leu Val Thr Lys Gln Arg Tyr
1 5 10

<210> 63

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 63

Lys His Tyr Leu Asn Leu Val Thr Arg Asn Arg Tyr
1 5 10

<210> 64

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 64

Lys His Tyr Leu Asn Leu Val Thr Arg Gln Lys Tyr
1 5 10

<210> 65

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 65

Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Thr
1 5 10

<210> 66

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 66

Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Phe
1 5 10

<210> 67

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 67

Arg His Thr Ile Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 68

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 68

Arg His Thr Val Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 69

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 69

Arg His Thr Leu Gln Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 70

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 70

Arg His Thr Leu Asn Ile Val Thr Arg Gln Arg Tyr
1 5 10

<210> 71

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 71

Arg His Thr Leu Asn Val Val Thr Arg Gln Arg Tyr
1 5 10

<210> 72

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 72

Arg His Thr Leu Asn Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 73

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 73

Arg His Thr Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 74

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 74

Arg His Thr Leu Asn Leu Val Ser Arg Gln Arg Tyr
1 5 10

<210> 75

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 75

Arg His Thr Leu Asn Leu Val Thr Lys Gln Arg Tyr
1 5 10

<210> 76

<211> 12

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 76

Arg His Thr Leu Asn Leu Val Thr Arg Asn Arg Tyr
1 5 10

<210> 77
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 77

Arg His Thr Leu Asn Leu Val Thr Arg Gln Lys Tyr
1 5 10

<210> 78
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 78

Arg His Thr Leu Asn Leu Val Thr Arg Gln Arg Thr
1 5 10

<210> 79
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 79

Arg His Thr Leu Asn Leu Val Thr Arg Gln Arg Phe
1 5 10

<210> 80
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 80

Arg His Phe Ile Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 81
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 81

Arg His Phe Val Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 82
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 82

Arg His Phe Leu Gln Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 83
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 83

Arg His Phe Leu Asn Ile Val Thr Arg Gln Arg Tyr
1 5 10

<210> 84
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 84

Arg His Phe Leu Asn Val Val Thr Arg Gln Arg Tyr
1 5 10

<210> 85
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 85

Arg His Phe Leu Asn Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 86
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 86

Arg His Phe Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 87
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 87

Arg His Phe Leu Asn Leu Val Ser Arg Gln Arg Tyr
1 5 10

<210> 88
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 88

Arg His Phe Leu Asn Leu Val Thr Lys Gln Arg Tyr
1 5 10

<210> 89
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 89

Arg His Phe Leu Asn Leu Val Thr Arg Asn Arg Tyr
1 5 10

<210> 90
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 90

Arg His Phe Leu Asn Leu Val Thr Arg Gln Lys Tyr
1 5 10

<210> 91
<211> 12
<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 91

Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Thr
1 5 10

<210> 92

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 92

Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Phe
1 5 10

<210> 93

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 93

Arg His Tyr Leu Gln Ile Val Thr Arg Gln Arg Tyr
1 5 10

<210> 94

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 94

Arg His Tyr Leu Gln Val Val Thr Arg Gln Arg Tyr
1 5 10

<210> 95

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 95

Arg His Tyr Leu Gln Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 96

<211> 12

<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 96

Arg His Tyr Leu Gln Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 97
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 97

Arg His Tyr Leu Gln Leu Val Ser Arg Gln Arg Tyr
1 5 10

<210> 98
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 98

Arg His Tyr Leu Gln Leu Val Thr Lys Gln Arg Tyr
1 5 10

<210> 99
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 99

Arg His Tyr Leu Gln Leu Val Thr Arg Asn Arg Tyr
1 5 10

<210> 100
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 100

Arg His Tyr Leu Gln Leu Val Thr Arg Gln Lys Tyr
1 5 10

<210> 101

<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 101

Arg His Tyr Leu Gln Leu Val Thr Arg Gln Arg Thr
1 5 10

<210> 102
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 102

Arg His Tyr Leu Gln Leu Val Thr Arg Gln Arg Phe
1 5 10

<210> 103
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 103

Arg His Tyr Leu Asn Ile Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 104
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 104

Arg His Tyr Leu Asn Ile Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 105
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 105

Arg His Tyr Leu Asn Ile Val Ser Arg Gln Arg Tyr
1 5 10

<210> 106

<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 106

Arg His Tyr Leu Asn Ile Val Thr Lys Gln Arg Tyr
1 5 10

<210> 107
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 107

Arg His Tyr Leu Asn Ile Val Thr Arg Asn Arg Tyr
1 5 10

<210> 108
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 108

Arg His Tyr Leu Asn Ile Val Thr Arg Gln Lys Tyr
1 5 10

<210> 109
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 109

Arg His Tyr Leu Asn Ile Val Thr Arg Gln Arg Thr
1 5 10

<210> 110
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 110

Arg His Tyr Leu Asn Ile Val Thr Arg Gln Arg Phe
1 5 10

<210> 111
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 111
Arg His Tyr Leu Asn Val Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 112
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 112

Arg His Tyr Leu Asn Val Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 113
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 113

Arg His Tyr Leu Asn Val Val Ser Arg Gln Arg Tyr
1 5 10

<210> 114
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 114

Arg His Tyr Leu Asn Val Val Thr Lys Gln Arg Tyr
1 5 10

<210> 115
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 115

Arg His Tyr Leu Asn Val Val Thr Arg Asn Arg Tyr
1 5 10

<210> 116
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 116

Arg His Tyr Leu Asn Val Val Thr Arg Gln Lys Tyr
1 5 10

<210> 117
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 117

Arg His Tyr Leu Asn Val Val Thr Arg Gln Arg Thr
1 5 10

<210> 118
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 118

Arg His Tyr Leu Asn Val Val Thr Arg Gln Arg Phe
1 5 10

<210> 119
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 119

Arg His Tyr Leu Asn Leu Ile Ser Arg Gln Arg Tyr
1 5 10

<210> 120
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 120

Arg His Tyr Leu Asn Leu Ile Thr Lys Gln Arg Tyr
1 5 10

<210> 121
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 121

Arg His Tyr Leu Asn Leu Ile Thr Arg Asn Arg Tyr
1 5 10

<210> 122
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 122

Arg His Tyr Leu Asn Leu Ile Thr Arg Gln Lys Tyr
1 5 10

<210> 123
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 123

Arg His Tyr Leu Asn Leu Ile Thr Arg Gln Arg Thr
1 5 10

<210> 124
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 124

Arg His Tyr Leu Asn Leu Ile Thr Arg Gln Arg Phe
1 5 10

<210> 125
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 125

Arg His Tyr Leu Asn Leu Leu Ser Arg Gln Arg Tyr
1 5 10

<210> 126
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 126

Arg His Tyr Leu Asn Leu Leu Thr Lys Gln Arg Tyr
1 5 10

<210> 127
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 127

Arg His Tyr Leu Asn Leu Leu Thr Arg Asn Arg Tyr
1 5 10

<210> 128
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 128

Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Lys Tyr
1 5 10

<210> 129
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 129

Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Thr
1 5 10

<210> 130
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 130

Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Phe
Page 28

1 5 10

<210> 131
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 131

Arg His Tyr Leu Asn Leu Val Ser Lys Gln Arg Tyr
1 5 10

<210> 132
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 132

Arg His Tyr Leu Asn Leu Val Ser Arg Asn Arg Tyr
1 5 10

<210> 133
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 133

Arg His Tyr Leu Asn Leu Val Ser Arg Gln Lys Tyr
1 5 10

<210> 134
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 134

Arg His Tyr Leu Asn Leu Val Ser Arg Gln Arg Thr
1 5 10

<210> 135
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 135

Arg His Tyr Leu Asn Leu Val Ser Arg Gln Arg Tyr
1 5 10

<210> 136
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 136

Arg His Tyr Leu Asn Leu Val Thr Lys Asn Arg Tyr
1 5 10

<210> 137
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 137

Arg His Tyr Leu Asn Leu Val Thr Lys Gln Lys Tyr
1 5 10

<210> 138
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 138

Arg His Tyr Leu Asn Leu Val Thr Lys Gln Arg Thr
1 5 10

<210> 139
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 139

Arg His Tyr Leu Asn Leu Val Thr Lys Gln Arg Phe
1 5 10

<210> 140
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 140

Arg His Tyr Leu Asn Leu Val Thr Arg Asn Lys Tyr
1 5 10

<210> 141
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 141

Arg His Tyr Leu Asn Leu Val Thr Arg Asn Arg Thr
1 5 10

<210> 142
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 142

Arg His Tyr Leu Asn Leu Val Thr Arg Asn Arg Phe
1 5 10

<210> 143
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 143

Arg His Tyr Leu Asn Leu Val Thr Arg Gln Lys Thr
1 5 10

<210> 144
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 144

Arg His Tyr Leu Asn Leu Val Thr Arg Gln Lys Phe
1 5 10

<210> 145
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 145

Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 146

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 146

Ile Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 147

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 147

Val Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 148

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 148

Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 149

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 149

Thr Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 150

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 150

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 151

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 151

Ser Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 152

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 152

Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 153

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 153

Thr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 154

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 154

Phe Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 155

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 155

Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg
1 5 10 15

Tyr

<210> 156

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Polypeptide variation

<400> 156

Thr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg
1 5 10 15

Tyr

<210> 157

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Polypeptide variation

<400> 157

Phe Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg
1 5 10 15

Tyr

<210> 158

<211> 18

<212> PRT

<213> Artificial sequence

<220>

<223> Polypeptide variation

<400> 158

Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln
1 5 10 15

Arg Tyr

<210> 159

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 159

Lys Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln
1 5 10 15

Arg Tyr

<210> 160
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 160

Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 161
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 161

Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 162
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 162

Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 163
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 163

Ile Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 164
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 164

Val Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
1 5 10 15
Arg Gln Arg Tyr
20

<210> 165
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 165

Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val
1 5 10 15
Thr Arg Gln Arg Tyr
20

<210> 166
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 166

Asp Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val
1 5 10 15
Thr Arg Gln Arg Tyr
20

<210> 167
<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 167

Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu
1 5 10 15
Val Thr Arg Gln Arg Tyr
20

<210> 168
<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 168

Asp Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu
1 5 10 15

Val Thr Arg Gln Arg Tyr
20

<210> 169
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 169

Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn
1 5 10 15

Leu Val Thr Arg Gln Arg Tyr
20

<210> 170
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 170

Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu
1 5 10 15

Asn Leu Val Thr Arg Gln Arg Tyr
20

<210> 171
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 171

Thr Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu
1 5 10 15

Asn Leu Val Thr Arg Gln Arg Tyr
20

<210> 172

<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 172

Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr
1 5 10 15

Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 173
<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 173

Ser Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr
1 5 10 15

Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 174
<211> 26
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 174

Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His
1 5 10 15

Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 175
<211> 26
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 175

Glu Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His
1 5 10 15

Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 176
<211> 27

<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 176

Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg
1 5 10 15

His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 177
<211> 27
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 177

Asp Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Arg
1 5 10 15

His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 178
<211> 28
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 178

Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu
1 5 10 15

Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 179
<211> 29
<212> PRT
<213> Artificial sequence

<220>
<223> Polypeptide variation

<400> 179

Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser
1 5 10 15

Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 180
<211> 30
<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 180

Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala
1 5 10 15

Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 181

<211> 30

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 181

Ser Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala
1 5 10 15

Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 182

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 182

Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr
1 5 10 15

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 183

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 183

Asp Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr
1 5 10 15

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 184

<211> 32

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 184

Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr
1 5 10 15
Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 185

<211> 33

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 185

Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg
1 5 10 15
Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg
20 25 30

Tyr

<210> 186

<211> 33

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 186

Arg Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg
1 5 10 15
Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg
20 25 30

Tyr

<210> 187

<211> 33

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 187

Gln Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg
1 5 10 15
Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg
20 25 30

Tyr

<210> 188
<211> 33
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 188

Asn Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg
1 5 10 15

Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg
20 25 30

Tyr

<210> 189
<211> 34
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 189

Leu Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn
1 5 10 15

Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln
20 25 30

Arg Tyr

<210> 190
<211> 34
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 190

Val Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn
1 5 10 15

Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln
20 25 30

Arg Tyr

<210> 191
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 191

Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 192
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 192

Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 193
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 193

Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 194
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 194

Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 195
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 195

Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg
1 5 10 15

Tyr

<210> 196
<211> 18
<212> PRT
<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 196

Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln
1 5 10 15

Arg Tyr

<210> 197

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 197

Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 198

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 198

Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 199

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 199

Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val
1 5 10 15

Thr Arg Gln Arg Tyr
20

<210> 200

<211> 22

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 200

Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu
1 5 10 15

Val Thr Arg Gln Arg Tyr
20

<210> 201
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 201

Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn
1 5 10 15

Leu Val Thr Arg Gln Arg Tyr
20

<210> 202
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 202

Ser Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn
1 5 10 15

Leu Val Thr Arg Gln Arg Tyr
20

<210> 203
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 203

Ala Ser Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr Leu
1 5 10 15

Asn Leu Val Thr Arg Gln Arg Tyr
20

<210> 204
<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation
<400> 204

Asp Ala Ser Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His Tyr
1 5 10 15

Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 205

<211> 26

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 205

Glu Asp Ala Ser Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu Lys His
1 5 10 15

Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 206

<211> 28

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 206

Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser Leu
1 5 10 15

Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 207

<211> 29

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 207

Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr Tyr Ala Ser
1 5 10 15

Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 208

<211> 29

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 208

Ala Pro Gly Glu Asp Ala Ser Glu Glu Leu Asn Arg Tyr Tyr Ala Ser

1 5 10 15

Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25

<210> 209

<211> 30

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 209

Glu Ala Pro Gly Glu Asp Ala Ser Glu Glu Leu Asn Arg Tyr Tyr Ala
1 5 10 15

Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 210

<211> 32

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 210

Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn Arg Tyr
1 5 10 15

Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 211

<211> 32

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 211

Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Glu Glu Leu Asn Arg Tyr
1 5 10 15

Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
20 25 30

<210> 212

<211> 33

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<400> 212

Ile Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Glu Glu Leu Asn Arg
1 5 10 15

Tyr Tyr Ala Ser Leu Lys His Tyr Leu Asn Leu Val Thr Arg Gln Arg
20 25 30

Tyr

<210> 213
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> ACETYLATION

<400> 213

Leu Arg His Tyr Ile Asn Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 214
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> ACETYLATION

<400> 214

Leu Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 215
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 215

Leu Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 216
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 216

Pro Ala Glu Asp Leu Ala Gln Tyr Ala Ala Glu Leu Arg His Tyr Leu
1 5 10 15

Asn Leu Leu Thr Arg Gln Arg Tyr
20

<210> 217
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 217

Leu Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 218
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 218

Met Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 219
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 219

Ala Arg Tyr Tyr Ser Ala Leu Arg His Phe Ile Asn Leu Ile Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 220
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D Ala

<220>
<221> MOD_RES
<222> (1)..(1)
<223> ACETYLATION

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 220

Xaa Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 221
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (18)..(18)
<223> AMIDATION

<400> 221

Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg Gln
1 5 10 15

Arg Tyr

<210> 222
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> Nle

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 222

xaa Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 223
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D Ser

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 223

xaa Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 224
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 224

Ala Ala Arg Tyr Ser His Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 225
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D Ile

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 225

Xaa Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 226
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES

<222> (20)..(20)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (1)..(1)
<223> ACETYLATION

<400> 226

Arg Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 227
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (18)..(18)
<223> AMIDATION

<400> 227

Gln Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg Gln
1 5 10 15

Arg Tyr

<210> 228
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 228

Ala Arg Phe Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 229
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> MeLeu

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 229

Xaa Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1. 5 10 15

Arg Gln Arg Tyr
20

<210> 230
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (1)..(1)
<223> METHYLATION

<400> 230

Leu Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1. 5 10 15

Arg Gln Arg Tyr
20

<210> 231
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> desamino

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 231

Xaa Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15
Arg Gln Arg Tyr
20

<210> 232
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (1)..(1)
<223> FORMYLATION

<400> 232

Ala Arg Tyr Tyr Ser Glu Leu Arg Arg Tyr Ile Asn Leu Ile Thr Arg
1 5 10 15
Gln Arg Tyr

<210> 233
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> Nva

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES

<222> (20)..(20)
<223> AMIDATION

<400> 233

Xaa Ala Arg Tyr Ala Ser Ala Leu Arg His Tyr Leu Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 234
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 234

Ala Arg Tyr Tyr Thr Gln Leu Arg His Tyr Ile Asn Leu Ile Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 235
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> desamino

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 235

Leu Ala Arg Tyr Tyr Ser Asn Leu Arg His Tyr Ile Asn Val Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 236
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 236

Ala Arg Tyr Tyr Asp Ser Leu Arg His Tyr Ile Asn Thr Ile Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 237
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 237

Ala Arg Tyr Tyr Ser Ala Leu Gln His Tyr Ile Asn Leu Leu Thr Arg
1 5 10 15

Pro Arg Tyr

<210> 238
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)

<223> AMIDATION

<400> 238

Leu Ala Arg Tyr Tyr Ser Ala Leu Arg Gln Tyr Arg Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Phe
20

<210> 239

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> N terminus is bonded to H

<220>

<221> MOD_RES

<222> (18)..(18)

<223> AMIDATION

<400> 239

Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln
1 5 10 15

Arg Phe

<210> 240

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> N terminus is bonded to H

<220>

<221> MOD_RES

<222> (19)..(19)

<223> AMIDATION

<400> 240

Ser Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 241

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> ACETYLATION

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 241

Ser Arg Tyr Tyr Ala Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 242
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> Nle

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 242

Xaa Ala Arg Tyr Tyr Asn Ala Leu Arg His Phe Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 243
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D isomer of Ala

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 243

Xaa Arg Tyr Glu Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg
1 5 10 15

His Arg Tyr

<210> 244
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (21)..(21)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Bz

<400> 244

Xaa Leu Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile
1 5 10 15

Thr Arg Pro Arg Phe
20

<210> 245
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 245

Ala Leu Tyr Tyr Ser Ala Leu Arg His Phe Val Asn Leu Ile Thr Arg
1 5 10 15

Gln Arg Tyr

<210> 246
<211> 19

<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D Ala

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 246

Xaa Arg Tyr Tyr Ser Ala Leu Arg His Tyr Val Asn Leu Ile Phe Arg
1 5 10 15

Gln Arg Tyr

<210> 247
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> MeSer

<220>
<221> MOD_RES
<222> (18)..(18)
<223> AMIDATION

<400> 247

Xaa Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Met Ile Thr Arg Gln
1 5 10 15

Arg Phe

<210> 248
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>

<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 248

Arg Ile Arg Tyr Tyr Ser Ala Leu Arg His Phe Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Phe
20

<210> 249
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminal is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 249

Leu Ser Arg Tyr Tyr Ser Ala Leu Arg His Phe Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 250
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Xaa is MeIle

<400> 250

Xaa Arg Tyr Tyr Ser Ala Leu Gln His Phe Ile Asn Leu Ile Thr Arg
1 5 10 15 -

Gln Arg Tyr

<210> 251
<211> 19
<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D Ser

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 251

Xaa Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg
1 5 10 15

Gln Arg Phe

<210> 252
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 252

Met Ala Arg Tyr Tyr Ser Asp Leu Arg Arg Tyr Ile Asn Leu Ile Thr
1 5 10 15

Arg Gln Arg Tyr
20

<210> 253
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>

<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 253

Ala Arg Tyr Tyr Ser Glu Leu Arg His Tyr Ile Ile Leu Ile Thr Arg
1 5 10 15
Gln Arg Tyr

<210> 254
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D Ala

<220>
<221> MOD_RES
<222> (20)..(20)
<223> AMIDATION

<400> 254

Xaa Ala Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr
1 5 10 15
Arg Gln Arg Tyr
20

<210> 255
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 255

Ala Ser Leu Arg His Trp Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 256
<211> 35
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (25)..(25)
<223> im DNP HIS; 2,2 diphenylalanine Hisitidine

<220>
<221> MOD_RES
<222> (35)..(35)

<223> AMIDATION

<400> 256

Tyr Pro Ala Lys Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu
1 5 10 15

Ser Thr Tyr Tyr Ala Ser Leu Arg Xaa Tyr Leu Asn Leu Val Thr Arg
20 25 30

Glx Arg Tyr
35

<210> 257

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (15)..(15)

<223> AMIDATION

<400> 257

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 258

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (15)..(15)

<223> AMIDATION

<400> 258

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Ala Arg Gln Arg Tyr
1 5 10 15

<210> 259

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (15)..(15)

<223> AMIDATION

<400> 259

Ala Ala Leu Arg His Tyr Leu Asn Leu Val Ala Arg Gln Arg Tyr
1 5 10 15

<210> 260
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 260

Ala Ser Leu Arg His Tyr Glu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 261
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa is Ornithine

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 261

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Xaa Arg Tyr
1 5 10 15

<210> 262
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> Xaa is p.C1.Pro; 4 chlorophenylalanine

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES

<222> (15)..(15)
<223> AMIDATION

<400> 262

Ala Ser Leu Arg Xaa Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 263
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 263

Ala Ser Leu Arg His Tyr Glu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 264
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is N Me Tyr

<400> 264

Ala Ser Leu Arg His Phe Glu Asn Leu Val Thr Arg Gln Arg Xaa
1 5 10 15

<210> 265
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa is Ornithine

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is N Me Tyr

<400> 265

Ala Ser Leu Arg His Tyr Glu Asn Leu Val Thr Arg Xaa Arg Xaa
1 5 10 15

<210> 266
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> LIPID
<222> (1)..(1)
<223> N alpha myristoyl

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 266

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 267
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N alpha naphthaleneacetyl

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 267

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 268

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MISC_FEATURE

<222> (15)..(15)

<223> Xaa is N Me Tyr

<220>

<221> MOD_RES

<222> (1)..(1)

<223> N alpha ACETYLATION

<220>

<221> MOD_RES

<222> (15)..(15)

<223> AMIDATION

<220>

<221> MISC_FEATURE

<222> (13)..(13)

<223> Xaa is Ornithine

<400> 268

Ala Ser Leu Arg His Phe Glu Asn Leu Val Thr Arg Xaa Arg Xaa
1 5 10 15

<210> 269

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (1)..(1)

<223> N alpha ACETYLATION

<220>

<221> MOD_RES

<222> (15)..(15)

<223> AMIDATION

<400> 269

Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 - 10 15

<210> 270

<211> 15

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> Xaa is 3 benzothienylalanine

<220>
<221> MOD_RES
<222> (7)..(7)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<400> 270

Ala Ser Leu Arg His Xaa Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 271
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Xaa is 4,4' biphenylalanine

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (16)..(16)
<223> AMIDATION

<400> 271

Xaa Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 272
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)

<223> AMIDATION

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> Xaa is 3 benzothienylalanine

<400> 272

Ala Ser Leu Arg His Xaa Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 273

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (1)..(1)

<223> N alpha ACETYLATION

<220>

<221> MOD_RES

<222> (15)..(15)

<223> AMIDATION

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> Xaa is 3 benzothienylalanine

<400> 273

Ala Ser Leu Arg His Xaa Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 274

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (1)..(1)

<223> N alpha ACETYLATION

<220>

<221> MOD_RES

<222> (15)..(15)

<223> AMIDATION

<400> 274

Ala Ser Leu Arg His Trp Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 275

<211> 15

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 275

Ala Ser Leu Arg His Trp Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 276
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> Xaa is 2 thiencylalanine

<400> 276

Ala Ser Leu Arg Asn Xaa Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 277
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> Xaa is tetrahydroisoquinoline

<400> 277

Ala Ser Leu Arg His Xaa Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 278
<211> 3
<212> PRT
<213> Homo sapiens

<400> 278

000
1

<210> 279
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (11)..(11)
<223> AMIDATION

<400> 279

His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 280
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (1)..(1)
<223> ACETYLATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is 2 thiencylalanine

<400> 280

Ala Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Xaa
1 5 10 15

<210> 281

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (1)..(1)

<223> N alpha ACETYLATION

<220>

<221> MOD_RES

<222> (16)..(16)

<223> AMIDATION

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> Xaa is 4 Thiazolylalanine

<400> 281

Ala Ser Leu Arg His Xaa Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 282

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (1)..(1)

<223> N alpha ACETYLATION

<220>

<221> MOD_RES

<222> (16)..(16)

<223> AMIDATION

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> Xaa is 4 Thiazolylalanine

<400> 282

Ala Ser Leu Arg His Xaa Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 283

<211> 3

<212> PRT

<213> Homo sapiens

<400> 283

000
1

<210> 284
<211> 3
<212> PRT
<213> Homo sapiens

<400> 284

000
1

<210> 285
<211> 3
<212> PRT
<213> Homo sapiens

<400> 285

000
1

<210> 286
<211> 3
<212> PRT
<213> Homo sapiens

<400> 286

000
1

<210> 287
<211> 3
<212> PRT
<213> Homo sapiens

<400> 287

000
1

<210> 288
<211> 3
<212> PRT
<213> Homo sapiens

<400> 288

000
1

<210> 289
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 289

Phe Ser Leu Arg Asn Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 290
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 290

Tyr Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 291
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 291

Ala Ser Leu Arg His Tyr Trp Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 292
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 292

Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 293
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 293

Ala Ser Leu Arg Ala Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 294
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (14)..(14)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> Xaa is 3' benzothienylalanine

<400> 294

Ala Ser Leu Arg Xaa Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 295
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 295

Ala Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 296
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 296

Ala Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Phe
1 5 10 15

<210> 297
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISCE_FEATURE
<222> (11)..(11)
<223> Xaa is D form of Trp

<220>
<221> MOD_RES
<222> (11)..(11)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (11)..(11)
<223> N alpha ACETYLATION

<400> 297

Ala Ser Leu Arg His Phe Leu Asn Leu Val Xaa Arg Gln Arg Tyr
1 5 10 15

<210> 298
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (13)..(13)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to CH3CO

<400> 298

Leu Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 299
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (13)..(13)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to CH3CO

<400> 299

Leu Arg His Tyr Ile Asn Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 300
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>

<221> MOD_RES
<222> (1)..(1)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (13)..(13)
<223> AMIDATION

<400> 300

Leu Arg His Tyr Leu Asn Leu Leu Thr Arg Gln Arg Tyr
1 5 10

<210> 301
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (13)..(13)
<223> AMIDATION

<400> 301

Leu Arg His Tyr Ile Asn Leu Ile Thr Arg Gln Arg Tyr
1 5 10

<210> 302
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE

<222> (10)..(10)
<223> Xaa is Norvaline

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> Xaa is Norleucine

<400> 302

Ala Ser Xaa Arg His Trp Xaa Asn Xaa Xaa Thr Arg Gln Xaa Xaa
1 5 10 15

<210> 303
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> Xaa is Norvaline

<400> 303

Ala Ser Xaa Arg His Trp Xaa Asn Trp Xaa Thr Arg Gln Xaa Xaa
1 5 10 15

<210> 304

<211> 15

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>

<221> MOD_RES

<222> (1)..(1)

<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> Xaa is Norvaline

<400> 304

Ala Ser Xaa Arg His Phe Xaa Asn Xaa Xaa Thr Arg Gln Xaa Xaa
1 5 10 15

<210> 305

<211> 15

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)

<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> Xaa is Norvaline

<400> 305

Ala Ser Xaa Arg His Phe Xaa Asn Trp Xaa Thr Arg Gln Xaa xaa
1 5 10 15

<210> 306
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (12)..(12)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (12)..(12)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<220>
<221> MISC_FEATURE
<222> (11)..(11)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH

<400> 306

Arg His Tyr Leu Asn Trp Val Thr Arg Gln Xaa Xaa

1 5 10

<210> 307
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (12)..(12)
<223> AMIDATION

<400> 307

Arg His Tyr Leu Asn Trp Val Thr Arg Gln Arg Tyr
1 5 10

<210> 308
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa is a psuedopeptide bond consisting of CH2 NH2

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is a psuedopeptide bond consisting of CH2 NH2

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> Xaa is Norvaline

<400> 308

Ala Ser Leu Arg His Tyr Xaa Asn Trp Xaa Thr Arg Gln Xaa Xaa
1 5 10 15

<210> 309
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH2

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa is a pseudopeptide bond consisting of CH2 NH2

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa is Norleucine

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> Xaa is Norvaline

<400> 309

Ala Ser Xaa Arg His Tyr Xaa Asn Trp Xaa Thr Arg Gln Xaa Xaa
1 5 10 15

<210> 310
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> bonded to OCH3

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<400> 310

Ile Asn Pro Ile Tyr Arg Leu Arg Tyr
1 5

<210> 311

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> DISULFID

<222> (4)..(4)

<223> Sequence is linked to identical sequence by a disulfide bond

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> N terminus is bonded to H

<220>

<221> MISC_FEATURE

<222> (9)..(9)

<223> C terminus is bonded to NH2

<400> 311

Ile Asn Pro Cys Tyr Arg Leu Arg Tyr
1 5

<210> 312

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Polypeptide variation

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> C terminus is bonded to OCH3

<220>

<221> DISULFID

<222> (1)..(1)

<223> sequence is linked to an identical sequence

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> N terminus is bonded to H

<400> 312

Cys Tyr Arg Leu Arg Tyr
1 5

<210> 313

<211> 8

<212> PRT

<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MISC_FEATURE
<222> (3)..(4)
<223> Connected by NH CH CO

<220>
<221> MISC_FEATURE
<222> (3)..(4)
<223> Identical peptide chains are connected by (CH2)4 at the CH o
f NH CH CO

<400> 313

Ile Asn Pro Tyr Arg Leu Arg Tyr
1 5

<210> 314
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminus is bonded to H

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> C terminus is bonded to OCH3

<400> 314

Tyr Arg Leu Arg Tyr Tyr Arg Leu Arg Tyr
1 5 10

<210> 315
<211> 34
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> DISULFID
<222> (18)..(22)
<223>

<400> 315

Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp Met Ala
1 5 10 15

Arg Cys Tyr Ser Ala Cys Arg His Tyr Ile Asn Leu Ile Thr Arg Gln
20 25 30

Arg Tyr

<210> 316
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 316

Arg His Tyr Leu Asn Leu Ile Gly Arg Gln Arg Tyr
1 5 10

<210> 317
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (3)..(7)
<223> ACETYLATION

<400> 317

Arg His Gly Leu Asn Leu Leu Gly Arg Gln Arg Tyr
1 5 10

<210> 318
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 318

Tyr Ile Asn Leu Ile Tyr Arg Leu Arg Tyr
1 5 10

<210> 319
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 319

His Tyr Ile Asn Leu Ile Tyr Arg Leu Arg Tyr
1 5 10

<210> 320
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 320

Arg His Tyr Ile Asn Leu Ile Tyr Arg Leu Arg Tyr
1 5 10

<210> 321
<211> 10

<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 321

Tyr Ile Asn Leu Leu Tyr Arg Gln Arg Tyr
1 5 10

<210> 322
<211> 15

<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> Xaa is 6 amino hexanoic acid

<400> 322

Tyr Pro Ser Leu Xaa Tyr Ile Asn Leu Ile Tyr Arg Leu Arg Tyr
1 5 10 15

<210> 323
<211> 9

<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 323

Ile Asn Leu Ile Tyr Arg Leu Arg Tyr
1 5

<210> 324
<211> 12

<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (12)..(12)
<223> AMIDATION

<400> 324

Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10

<210> 325
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 325

Ala Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 326
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> N terminal is bonded to H

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 326

Ala Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 327
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>

<221> MOD_RES
<222> (9)..(9)
<223> AMIDATION

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<400> 327

Ala Ser Leu Arg Thr Arg Gln Arg Tyr
1 5

<210> 328
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> Xaa is 2 thienylalanine

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 328

Ala Ser Leu Arg His Xaa Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 329
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<220>
<221> MOD_RES
<222> (1)..(1)
<223> N alpha ACETYLATION

<220>
<221> MOD_RES
<222> (15)..(15)
<223> AMIDATION

<400> 329

Tyr Ser Leu Arg His Phe Leu Asn Leu Val Thr Arg Gln Arg Tyr
1 5 10 15

<210> 330

<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 330

Asp Asp Asp Asp Tyr
1 5

<210> 331
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 331

Gly Pro Arg
1

<210> 332
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 332

Ala Gly Gly
1

<210> 333
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 333

His Pro Phe His Leu
1 5

<210> 334
<211> 34
<212> PRT
<213> Homo sapiens

<400> 334

Ile Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu Leu Asn
1 5 10 15
Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr Arg Gln
20 25 30

Arg Tyr

<210> 335
<211> 34
<212> PRT
<213> Artificial Sequence

<220>
<223> Polypeptide variation

<400> 335

Ser Lys Pro Asp Asn Pro Gly Glu Asp Ala Pro Ala Glu Asp Met Ala
1 5 10 15

Arg Tyr Tyr Ser Ala Leu Arg His Tyr Ile Asn Leu Ile Thr Arg Gln
20 25 30

Arg Tyr

<210> 336
<211> 37
<212> PRT
<213> Homo sapiens

<400> 336

His Asp Glu Phe Glu Arg His Ala Glu Gly Thr Phe The Ser Asp Val
1 5 10 15

Ser Ser Tyr Leu Glu Gly Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu
20 25 30

Val Lys Gly Arg Gly
35

<210> 337
<211> 36
<212> PRT
<213> Homo sapiens

<220>
<221> MISC_FEATURE
<222> (36)..(36)
<223> C terminus is bonded to NH2

<400> 337

His Asp Glu Phe Glu Arg His Ala Glu Gly Thr Phe The Ser Asp Val
1 5 10 15

Ser Ser Tyr Leu Glu Gly Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu
20 25 30

Val Lys Gly Arg
35

<210> 338
<211> 31
<212> PRT
<213> Homo sapiens

<400> 338

His Ala Glu Gly Thr Phe The Ser Asp Val Ser Ser Tyr Leu Glu Gly
1 5 10 15

Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
20 25 30

<210> 339
<211> 30
<212> PRT
<213> Homo sapiens

<220>
<221> MISC_FEATURE
<222> (30)..(30)
<223> C terminus is bonded to NH2
<400> 339

His Ala Glu Gly Thr Phe The Ser Asp Val Ser Ser Tyr Leu Glu Gly
1 5 10 15

Gly Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg
20 25 30

<210> 340
<211> 37
<212> PRT
<213> Homo sapiens
<400> 340

His Ser Gln Gly Thr Phe Thr Ser Asp Tyr Ser Lys Tyr Leu Asp Ser
1 5 10 15

Arg Arg Ala Gln Asp Phe Val Gln Trp Leu Met Asp Thr Lys Arg Asn
20 25 30

Lys Asn Asn Ile Ala
35

<210> 341
<211> 463
<212> PRT
<213> homo sapiens

<400> 341

Met Ala Gly Ala Pro Gly Pro Leu Arg Leu Ala Leu Leu Leu Gly
1 5 10 15

Met Val Gly Arg Ala Gly Pro Arg Pro Gln Gly Ala Thr Val Ser Leu
20 25 30

Trp Glu Thr Val Gln Lys Trp Arg Glu Tyr Arg Arg Gln Cys Gln Arg
35 40 45

Ser Leu Thr Glu Asp Pro Pro Pro Ala Thr Asp Leu Phe Cys Asn Arg
50 55 60

Thr Phe Asp Glu Tyr Ala Cys Trp Pro Asp Gly Glu Pro Gly Ser Phe
65 70 75 80

Val Asn Val Ser Cys Pro Trp Tyr Leu Pro Trp Ala Ser Ser Val Pro
85 90 95

Gln Gly His Val Tyr Arg Phe Cys Thr Ala Glu Gly Leu Trp Leu Gln
100 105 110

Lys Asp Asn Ser Ser Leu Pro Trp Arg Asp Leu Ser Glu Cys Glu Glu
115 120 125

Ser Lys Arg Gly Glu Arg Ser Ser Pro Glu Glu Gln Leu Leu Phe Leu
130 135 140

Tyr Ile Ile Tyr Thr Val Gly Tyr Ala Leu Ser Phe Ser Ala Leu Val
145 150 155 160

Ile Ala Ser Ala Ile Leu Leu Gly Phe Arg His Leu His Cys Thr Arg
165 170 175

Asn Tyr Ile His Leu Asn Leu Phe Ala Ser Phe Ile Leu Arg Ala Leu
180 185 190

Ser Val Phe Ile Lys Asp Ala Ala Leu Lys Trp Met Tyr Ser Thr Ala
195 200 205

Ala Gln Gln His Gln Trp Asp Gly Leu Leu Ser Tyr Gln Asp Ser Leu
210 215 220

Ser Cys Arg Leu Val Phe Leu Leu Met Gln Tyr Cys Val Ala Ala Asn
225 230 235 240

Tyr Tyr Trp Leu Leu Val Glu Gly Val Tyr Leu Tyr Thr Leu Leu Ala
245 250 255

Phe Ser Val Phe Ser Glu Gln Trp Ile Phe Arg Leu Tyr Val Ser Ile
260 265 270

Gly Trp Gly Val Pro Leu Leu Phe Val Val Pro Trp Gly Ile Val Lys
275 280 285

Tyr Leu Tyr Glu Asp Glu Gly Cys Trp Thr Arg Asn Ser Asn Met Asn
290 295 300

Tyr Trp Leu Ile Ile Arg Leu Pro Ile Leu Phe Ala Ile Gly Val Asn
305 310 315 320

Phe Leu Ile Phe Val Arg Val Ile Cys Ile Val Val Ser Lys Leu Lys
325 330 335

Ala Asn Leu Met Cys Lys Thr Asp Ile Lys Cys Arg Leu Ala Lys Ser
340 345 350

Thr Leu Thr Leu Ile Pro Leu Leu Gly Thr His Glu Val Ile Phe Ala
355 360 365

Phe Val Met Asp Glu His Ala Arg Gly Thr Leu Arg Phe Ile Lys Leu
370 375 380

Phe Thr Glu Leu Ser Phe Thr Ser Phe Gln Gly Leu Met Val Ala Ile
385 390 395 400

Leu Tyr Cys Phe Val Asn Asn Glu Val Gln Leu Glu Phe Arg Lys Ser
405 410 415

Trp Glu Arg Trp Arg Leu Glu His Leu His Ile Gln Arg Asp Ser Ser
420 425 430

Met Lys Pro Leu Lys Cys Pro Thr Ser Ser Leu Ser Ser Gly Ala Thr
435 440 445

Ala Gly Ser Ser Met Tyr Thr Ala Thr Cys Gln Ala Ser Cys Ser
450 455 460

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- BLACK BORDERS**
- IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- FADED TEXT OR DRAWING**
- BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- SKEWED/SLANTED IMAGES**
- COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- GRAY SCALE DOCUMENTS**
- LINES OR MARKS ON ORIGINAL DOCUMENT**
- REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.